



A CASE OF HYPOXIC ENCEPHALOPATHY WITH DELAYED EXACERBATION

General Medicine

Dr Mayuresh Chavan

Dr A Sankar Associate Professor.

Dr Umashankar Assistant Professor.

ABSTRACT

Most patients contract hypoxic encephalopathy after suffering a cardiac arrest. They usually endure severe neurological sequelae and the temporal profile of the disease progression remains unclear. This case study shows how the effects of hypoxic encephalopathy continue to progress for several years after the initial event. Up to eight years after the hypoxic insult, the patient's intellect steadily deteriorated, and brain atrophy progressed. As the hypoxic insult on the brain is only transient, the neurological disability seems not to be exacerbated for years. However, our case indicates that this disorder may have a long progression.

KEYWORDS

dementia, encephalopathy, hypoxia, MRI

INTRODUCTION

Recent medical progress has led to an increase in the number of cardiac arrest survivors. According to one estimate, up to 200 people per 1 million inhabitants are saved after cardiac arrest in developed countries. However, about half of these survivors suffer severe neurological sequelae, because the brain is the organ most vulnerable to hypoxia. Focal neurological deficit such as sensorimotor impairment is rare, but neuropsychological symptoms are common, which can be attributed to cortical as well as diffuse white matter injury.

The clinical features and their influence on the eventual mortality rate have been investigated. However the temporal profile of disease progression, and the clinical characteristics of the chronic stage have not yet been fully elucidated. Even after an intensive search, we could find only a few reports in which radiological exacerbation was demonstrated for no more than a few months. Thus whether or not the neurological disability continues to progress for years in hypoxic encephalopathy is still unclear. In this report, we describe a case of hypoxic encephalopathy with years of slow progression, both clinically and radiologically. The hypoxic insult may be only transient, but the progression of hypoxic encephalopathy may be long-lasting.

CASE REPORT

A 53-year-old woman with no remarkable disease history was referred to our hospital because of abdominal pain and liver dysfunction. Cholelithiasis was suspected, and endoscopic retrograde cholangiopancreatography (ERCP) was carried out on December 9, 1999. Papillotomy was simultaneously performed, and the stones in the duct were removed. Just after the procedure, the physician noticed that the patient was not breathing and was pulseless. She had ventricular fibrillation, and intensive resuscitation was immediately attempted. Defibrillation successfully achieved regular heart sinus rhythm. Blood pressure was maintained using catecholamines. She was intubated and mechanically ventilated, and the arterial blood gas was kept within the appropriate range thereafter. The duration of hypoxia suffered by her brain is uncertain, but was estimated to be less than 10 minutes because a nurse confirmed that the patient was alert several minutes before the physician noticed her collapse.

During resuscitation, the physician confirmed that the patient's bilateral pupils were dilated to 8 mm in diameter. Three hours later, this dilation decreased to 3 mm in diameter. The patient was comatose for the first several hours, but could open her eyes when called 12 hours after resuscitation. Normal respiration and blood pressure were gradually regained. She was extubated at 5 days and catecholamines were discontinued at 6 days.

On December 15, when her cardiopulmonary state became stable, she was examined by our neurology department. She was awake and could obey only simple requests such as hand grasping. Precise verbal communication was difficult for her. Although obeying complex requests was difficult at that time, no definite sensorimotor

neurological deficits were confirmed. A electroencephalogram revealed theta waves of 5 to 7 Hz bilaterally, with no paroxysms. Computed tomography (CT) (Figures 1A and B) showed decreased density at the right caudate head (Figure 1A, arrowheads) and diffuse low density in the bilateral cerebral white matter (Figure 1B, arrows). Marked brain edema was not observed however (Figures 1A and B).

Her mental state gradually improved. Two weeks later when she visited the neurology department for the second time, she could verbally express her distress. On the other hand, her recent memory was still severely impaired. Furthermore, motor clumsiness and intention tremor became apparent, which may only have been revealed as a result of our extensive neurological examination. Nevertheless, the patient could walk independently. She was discharged and returned home. Although dependent, she could live in her own home with no extreme difficulty. Magnetic resonance imaging (MRI) (Figures 2A–F) revealed scattered foci of high intensity lesions in the cerebral cortex of the T1-weighted image (Figures 2A and 2C, arrowheads). The right caudate head, which showed decreased density by CT in the acute stage (Figure 1A), showed high intensity in the T1-weighted image (Figure 2B, arrow). In addition, slight brain atrophy was suspected when the patient's age was taken into account. The T2-weighted image, however, showed no abnormal intensity at this time point (Figures 2D–F).

Her neurological findings seemed stable. Dementia as well as clumsiness and intention tremor continued, but no exacerbation was apparent for a while. On the Mini-Mental Status Examination, the patient had gained 18 points. Disturbance of calculation and recent memory markedly decreased the total score, but both spatial and temporal orientation had been spared any damage. Nevertheless, her family felt that her mentality began to deteriorate from that time. They insisted that the patient had become more dependent in daily activities, such as excretion and bathing. We also noticed that she needed more support in her daily activities. Her disattentive state worsened, though motor function seemed stable. Her score on the Dementia Scale at was 8 points, with marked disorientation as well as memory disturbance. MRI examined on the same day (Figures 2G–L) revealed that the cerebral sulci had become wider (Figures 2H arrowhead, 2I arrow), suggesting that the cerebral cortex had degenerated further. Dilatation of the lateral ventricles (Figure 2H, arrows) indicated that cerebral white matter had decreased in volume. In some subcortical white-matter regions, high intensity lesions were newly discernible by T2-weighted image (Figure 2L, arrow).

Her mentality continued to deteriorate thereafter. Her ability to move worsened, so that from late that year, she could no longer walk independently. Because of the difficulty in coming to hospital, her care was taken over by a physician and she did not visit our neurology department for more than 1 year. When she visited us, establishing a score on the Revised Dementia Scale was difficult because of her severe dementia: she was unable to answer our questions. Evaluating precise motor function was also difficult, but intention tremor was still

evident. CT showed marked cortical atrophy and dilatation of ventricles (Figures 1C and D). MRI (Figures 2M–R) revealed further widened cerebral sulci (Figure 2N, arrowheads) and pronounced lateral ventricle dilatation (Figures 2M–O, arrows). The third ventricle was also dilated (Figure 2N, arrow). The T2-weighted image showed scattered high-intensity foci in the subcortical white matter of the occipital (Figure 2Q, arrows) and parietal (Figure 2R, arrow) lobes. It was suggested that cortical as well as subcortical structures were affected in a delayed manner.

The patient continues to be cared for in her home, because her family strongly prefers it. A helper visits frequently, and monitors her general condition carefully.

DISCUSSION

Hypoxic encephalopathy can be caused by various disorders, but the most common etiology is cardiac arrest. Coma and seizure are commonly observed in the first several hours to days. About half of these patients die, but others take the course of gradual recovery or persistent vegetative state. The determinants of neurological outcome are not fully elucidated, but experimental studies suggest that brain temperature and blood glucose level influence the prognosis. Clinical studies show that long duration of initial unconsciousness, and the occurrence of generalized myoclonus, may be ominous indicators of a poor outcome. Of interest is that delayed resuscitation does not necessarily mean severe neurological sequelae (Roine et al 1993a); some neurons may degenerate only when post-ischemic activity is regained early (discussed below).

Apart from the above mentioned clinical and experimental studies, no reports were found of the investigation of disease progression over several years. Roine and colleagues (1993a) followed neuropsychological sequelae after hypoxic encephalopathy, and showed that patients with severe cognitive impairment gradually declined. However, we should note that their patients were studied for only up to 12 months. Some patients with mild or moderate cognitive impairment could possibly show cognitive decline several years later. We found no studies that followed neurological sequelae of hypoxic encephalopathy for more than 1 year. This may be, at least in part, because patients suffering from hypoxic encephalopathy are not followed by the same doctor for very long.

Our case showed not only clinical but also radiological exacerbation in a delayed manner (Figures 1 and 2). Early CT features of hypoxic encephalopathy include decreased density in the basal ganglia, cerebral cortex, and cerebral white matter especially at the watershed area. Our case also showed these features, though decreased density in the cerebral cortex was obscure (Figures 1A and B). A previous report showed that the CT number in the putamen correlated with good prognosis. If this rule had been applied to our case, the prognosis would have been very poor, for example, an early demise. However, our patient at one stage recovered to the extent of being able to walk. Decreased density in CT does not necessarily mean neuronal loss, but rather suggests brain edema. Severe brain edema is caused by severe energy failure, massive production of radical oxygen species, or copious release of excitatory amino acids. Our patient's brain may have suffered these biochemical insults at the acute stage, but fortunately these were resolved promptly. This may be, as discussed later, one of the causes of this case's unique temporal profile of disease progression.

Early MRI findings in our case were also compatible with previous reports, scattered high intensity spots by T1-weighted image were confirmed at the cerebral cortex and putamen (Figures 2A–C). Those in the cerebral cortex reflect cortical laminar necrosis, because some neuronal populations show selective vulnerability against hypoxia. On the other hand, the T2-weighted image at that time (Figures 2D–F) showed no definite abnormalities. A previous report demonstrated that high intensity lesions in the T2-weighted image were confirmed as early as several days after the hypoxic insult, but another showed that they appeared only several months after the insult.

Our case results were similar to those of the latter report, that is, high intensity subcortical lesions in the T2-weighted image appeared only in the chronic stage (Figures 2J–L, and 2P–R). Of interest is that subcortical high intensity lesions in the T2-weighted image should be the cause of delayed exacerbation; even though it was investigated only for several months, cases with subcortical high intensity by T2-

weighed image showed progression of brain atrophy. If we can clarify what is occurring in this T2-weighted high intensity area, we may be able to understand the mechanisms of delayed exacerbation in hypoxic encephalopathy.

The molecular mechanisms of delayed exacerbation seen in this case are difficult to explain. After transient global ischemia in the brain, the phenomenon of “delayed neuronal cell death (DND)” has gained much attention. DND of the hippocampal pyramidal cells in rodent models is well known, but some cortical as well as caudate neurons also show such degeneration. However, this kind of neuronal loss occurs only several days after transient global ischemia. Whether neuronal cell death occurs years after transient ischemic insult has not yet been investigated, because it is impossible to use rodents as model animals for such studies.

Our case suffered a sudden cardiopulmonary collapse in the hospital. Intensive resuscitation was quickly started. Cerebral blood flow with appropriate oxygen level was restored promptly. Paradoxically, this procedure might have been the cause of delayed exacerbation in this case. In an experimental study, some neurons degenerated only when the ischemic period was shorter than 10 minutes, but were spared when it was longer than 10 minutes. Sustained neuronal activity, established only when the ischemic period is quite short, could lead to delayed neuronal cell death. Although this may be a rare case, the symptoms of some patients who suffer hypoxic encephalopathy may continue to progress over a very long period.

Disclosure

The authors report no conflicts of interest in this work.





Fig 1






Fig 2

REFERENCES

1. [BRCT] Brain Resuscitation Clinical Trial II Study Group. 1991. A randomized clinical study of a calcium-entry blocker (lidoflazine) in the treatment of comatose survivors of cardiac arrest. *N Engl J Med*, 324:1225–31.
2. Hossman KA. 2006. Pathophysiology and therapy of experimental stroke. *Cell Mol Neurobiol*, 26:1057–83.
3. Jeong JW, Kim KW, Lee DY, et al. 2007. A normative study of the Revised Hasegawa Dementia Scale: comparison of demographic influences between the Revised Hasegawa Dementia Scale and the Mini-Mental Status Examination. *Dement Geriatr Cogn Disord*, 24:288–93.
4. Kjos BO, Brant-Zawadzki M, Young RG. 1983. Early CT findings of global central nervous system hypoperfusion. *Am J Roentgenol*, 141:1227–32.
5. Kirino T, Sano K. 1984. Selective vulnerability in the gerbil hippocampus following transient ischemia. *Acta Neuropathol*, 61:201–8.