



## ORAL SUBMUCOUS FIBROSIS: OVERVIEW

### Dental Science

**Dr. Mandeep Kaur Sandhu**

B.D.S, BABA Farid University Of Health Sciences, Faridkot, Punjab, India.

### ABSTRACT

Oral submucous fibrosis (OSMF) is a chronic, potentially malignant disorder of the oral cavity characterized by progressive fibrosis of the submucosal tissues. Recently, due to advances in the field of diagnostic pathology, advanced diagnostic aids are used to guide the dentist about the nature of disease and its prognosis. Management of OSMF has also seen changes. These advancements have aimed to improve treatment outcomes, relieve symptoms, and prevent the malignant transformation of the condition.

### KEYWORDS

OSMF, Arecanut, management, recent, advances

### INTRODUCTION

Oral Submucous Fibrosis (OSMF) is an Oral potentially malignant disorder. It is a collagen metabolism disorder. A number of definitions have been proposed by a number of researchers.<sup>[1]</sup> The ancient medical literature has evidence of definitions by Sushruta where it was popularly known as 'Vidari'. J.Schwartz in 1952 also fondly described it as Atropicaidiopathica (tropica) Mucosae Oris in five Indian migrant women.<sup>[2]</sup> However, it was Pindborg who coined the term OSMF and gave an extensive definition of the same.<sup>[3]</sup> The epidemiology of OSMF is dependent on the sociodemographic factors of the population.<sup>[4]</sup> OSMF is a disease of middle age group. Usually, patients are 20-40 years of age and this is progressing towards younger age group. In India, the prevalence of OSMF is 0.2-2.3% in males and 1.2-4.6% in females.<sup>[5]</sup> Males are found to be more affected due to easy habit acquisition. Regionally, it is more prevalent in the South-east Asia India, Sri Lanka, Maldives, Bangladesh, Myanmar, Taiwan and various islands in South Pacific contributing to more than half of the global prevalence. There are many other local factors besides arecanut, like capsaicin, tobacco, spicy foods and alcohol have synergistic effects. Systemic factors like iron, Vitamin B12 deficiency, anemia, genetic predisposition are also contributory.<sup>[6]</sup> These etiologic factors directly dictate the pathogenesis of OSMF. This review provides an overview of OSMF with recent advances in diagnosis and management.

### Pathogenesis

Although the pathogenesis of OSMF is one of the most researched area, its pathogenesis is still obscure. There are a number of proposed mechanisms for the same:

### OSMF "collagen metabolism disorder"

Disturbance in collagen turnover is central to pathogenesis of OSMF. Arecanut which is a causative factor for OSMF contains alkaloids and flavonoids. The alkaloids are responsible for collagen synthesis whereas flavonoids decrease the activity of collagenase-the enzyme responsible for breakdown of collagen. The net effect is increased collagen production and decreased collagen degradation.<sup>[3]</sup> In an interesting study, the qualitative properties of collagen were assessed with histopathological grading of OSMF. Picrosirius red stain was used to locate collagen fibers.

The submucosa exhibited a denser arrangement of collagen fibers, indicating the initiation of fibrosis in that region. The progressive histopathological grades showed an elevated presence of perpendicular type III fibers, implying their involvement in the fibrotic process.<sup>[7]</sup>

### Effect of arecanut alkaloids on fibroblasts

Alkaloids have direct effect on proliferation of fibroblast which in turn increase the production of collagen. When mixed with slaked lime, arecanut is converted to arecadine which stimulates fibroblasts. This phenomenon may arise from the clonal selection of a specific cell population within the modified tissues, influenced by local factors such as interleukin-1 released by inflammatory cells.<sup>[8]</sup>

### Collagen stabilization

The presence of Areca flavonoids, tannins, and catechins can lead to

increased fibrosis by promoting the formation of a more resilient and insoluble collagen structure through the inhibition of collagenase enzyme activity.<sup>[8]</sup> Type I collagen which is normally present in the oral cavity is completely replaced by type III collagen which is more resilient.<sup>[9]</sup> At a molecular level also, there is excess of alpha 1 chain than alpha 2 chains. Hence the collagen structure becomes more stable and less prone to degradation.<sup>[10]</sup>

### Nutritional deficiency

Nutritional deficiencies like iron, vitamin, protein, etc are seen frequently in OSMF patients. Hydroxyproline, an amino acid exclusively present in collagen, is integrated into collagen in its hydroxylated form, specifically as 4-hydroxyproline. This process relies on the presence of iron and ascorbic acid. The decline in iron levels might be attributed to iron utilization in the fibrotic process.<sup>[11]</sup>

### Diagnosis

The clinical presentation of OSMF is very characteristic and diagnosis is usually done by oral examination.

- Reduced mouth opening.
- Other signs such as ulcers, xerostomia, blanching of oral mucosa, etc.
- In advanced cases the upper aerodigestive tract also has strictures.<sup>[12]</sup> The fibrous bands are progressive and makes opening of mouth difficult.

Many researchers have proposed various classifications based on the clinical features observed. The classification proposed by More et al. presents the clinical and functional staging of OSMF. This system of classification is appropriate for Indian population.<sup>[13]</sup>

There are many definitive methods to diagnose OSMF. One of the methods is the gold standard histopathological examination. A solid biopsy is performed which is wedge shaped from the most representative site. The findings can be described as mucosal and submucosal changes. Mucosal changes include atrophy of the epithelium, loss of rete ridges and liquefactive degeneration of basal layer. Rarely, pigment incontinence can be seen. Submucosa is very characteristic. Diffuse fibrosis in the submucosa with a chronic inflammatory infiltrate is the hallmark of OSMF.<sup>[14]</sup>

Exfoliative cytology although of little use, is a quick chairside investigation. In a study evaluating cellular features of OSMF, the keratinised cells and intermediate buccal cells were stained green to blue in colour and the cytoplasm appeared creased.<sup>[15]</sup>

Cyclophilin A (CYPA) was identified as a biomarker and a target for gene intervention in OSMF through proteomic two-dimensional electrophoresis (2-DE).<sup>[16]</sup>

Use of Ultrasonography (USG) in OSMF gave promising results. It is a non-invasive method to grade the degree of fibrosis. The elastography score represents the relative elasticity ratio between areas that appear normal and areas that appear abnormal.<sup>[17]</sup>

Autofluorescence spectroscopy, optical coherence tomography (OCT), and Fourier transform infrared spectroscopy (FTIR) were used in patients who do not undergo biopsy for OSF diagnosis.<sup>[18]</sup>

## Management

As per the definition by Pindborg, OSMF is a chronic disease. Conservative treatments usually are of little benefit. Habit intervention remains the central point of OSMF treatment. Patient education regarding ill effects of areca nut and measures should be taken to stop its usage. Nevertheless, for the ease of understanding the treatment of OSMF can be divided as: conservative therapy, pharmacotherapy and surgical intervention.

### 1. Conservative therapy

For mild cases of OSMF i.e., mouth opening greater than 25 mm conservative therapy can be considered. Physiotherapy such as using mouth props, various splints, jaw stretching exercises have proved to be beneficial in early stages of OSMF.<sup>[17]</sup>

### 2. Pharmacotherapy

A lot of research has been done to find the medicinal treatment of OSMF. Some of these aim at reversal of the disease process while some provide symptomatic control.

Steroids have been the drug of choice in treatment of OSMF since a long time. Their anti-inflammatory property has proved to be highly beneficial. Commonly used steroids are hydrocortisone, triamcinolone, methylprednisolone, betamethasone, dexamethasone and clobetasol propionate. They can be used topically or as intralesional injections.<sup>[19,20]</sup>

Interferon gamma: This cytokine is recognized for its anti-fibrotic properties, as it promotes the production of collagenase in the oral epithelium and the underlying lamina propria, while inhibiting the proliferation of fibroblasts.<sup>[21]</sup>

Herbal extracts: Herbal medicines such as colchicine,<sup>[22]</sup> basil, turmeric,<sup>[23]</sup> aloe vera,<sup>[24]</sup> lycopene<sup>[25]</sup> are vastly researched to aid in treatment of OSMF. All of them have satisfactory results in a long term.

Iron supplements: Iron is essential for epithelial turnover. Deficiency of iron leads to atrophic epithelium. Numerous research studies have indicated a decline in haemoglobin and serum ferritin levels in OSMF, with the iron content being depleted during collagen synthesis.<sup>[26]</sup> Hence iron supplements can be beneficial in treatment of OSMF.

Fibrinolytic therapy: The fibrosis in OSMF can be broken down by the means of proteolytic enzymes such as hyaluronidase, collagenase, and chymotrypsin and have proved to be successful.<sup>[27]</sup>

### 3. Surgical treatment

Treatment of patients with advanced stages of OSMF is a challenging task. Surgical treatment of OSMF primarily aims to increase the mouth opening. The prognosis of surgical interventions have been debatable. The treatment options for OSMF essentially involve excision of the fibrous bands, excision of bands with myotomy either with or without coronoideotomy, and covering the exposed area with various materials such as skin grafts, fresh amnion, collagen membrane, buccal fat pad, local flaps, or vascularized free flaps. This is followed by proactive post-operative jaw physiotherapy, incorporating antioxidants and ensuring proper nutrition. Regular follow-ups are essential to maintain oral opening and promptly identify any potential malignant changes.<sup>[28]</sup> In a systematic review, it is observed that lasers such as Diode, KTP-532, ErCr:YSGG have proved to be promising in treatment of OSMF.

### 4. Stem Cell Therapy

Researchers have investigated the potential of stem cell-based approaches, particularly mesenchymal stem cell (MSC) transplantation, to reverse fibrosis and facilitate tissue regeneration in OSMF. MSCs possess immunomodulatory and regenerative properties, making them a prospective therapeutic option for OSMF. However, further investigation is necessary to establish their effectiveness and safety.<sup>[29]</sup>

### 5. Targeted Therapies

The management of OSMF has witnessed the emergence of molecular targeted therapies. These therapies focus on specific signaling pathways involved in fibrosis, such as the transforming growth factor-beta (TGF-β) and Janus kinase/signal transducer and activator of transcription (JAK/STAT) pathways. In preclinical studies, drugs targeting these pathways have demonstrated potential by disrupting the fibrotic process and potentially impeding the progression of the disease.<sup>[32]</sup>

## CONCLUSION

In recent years, significant strides have been made in both diagnosing and managing oral submucous fibrosis (OSMF). Cutting-edge diagnostic techniques, including optical coherence tomography (OCT) and confocal microscopy, have emerged as promising non-invasive methods for detecting OSMF-related changes in the oral mucosa at an early stage. These innovative technologies play a crucial role in facilitating timely diagnosis and monitoring disease progression. Moreover, the field of pharmacological interventions has witnessed remarkable progress, broadening the array of treatment options for OSMF. Antioxidants, immunomodulators, and collagen cross-linking agents have been extensively studied for their potential in alleviating symptoms and modulating the fibrotic process. Surgical interventions have also evolved, with techniques such as laser therapy offering minimally invasive alternatives to enhance mouth opening and foster tissue regeneration. However, further advancements and optimization of care necessitate continued research and clinical trials in order to refine these innovative approaches and benefit individuals affected by OSMF.

## REFERENCES

- More CB, Rao NR. Proposed clinical definition for oral submucous fibrosis. *J Oral Biol Craniofacial Res.* 2019;9(4):311–4.
- Shah KM. Association of Candida species with Oral submucous fibrosis and Oral leukoplakia: a case control study. *Annals of Clinical and Laboratory Research.* 2018;6(3):248.
- Rajalalitha P, Vali S. Molecular pathogenesis of oral submucous fibrosis—a collagen metabolic disorder. *Journal of oral pathology & medicine.* 2005 Jul;34(6):321–8.
- Singh AG, Roy S, Oza S, Singhavi H, Chatterjee K, Chaturvedi P. A contemporary narrative review to guide molecular epidemiology of oral submucous fibrosis. *Int J Mol Epidemiol Genet.* 2021 Aug 15;12(4):61–70.
- Rao NR, Villa A, More CB, Jayasinghe RD, Kerr AR, Johnson NW. Oral submucous fibrosis: a contemporary narrative review with a proposed inter-professional approach for an early diagnosis and clinical management. *J Otolaryngol Head Neck Surg.* 2020;49:3.
- Rajendran R. Oral submucous fibrosis: etiology, pathogenesis, and future research. *Bull World Health Organ.* 1994;72(6):985–96.
- Nishat R, Kumar H. Collagen fibers in oral submucous fibrosis - A polarizing microscopy study using two special stains. *Indian J PatholMicrobiol.* 2019 Oct-Dec;62(4):537–543.
- Meghji S, Scutt A, Harvey W, Canniff JP. An in vitro comparison of human fibroblasts from normal and oral submucous fibrosis tissue. *Arch Oral Biol*1987;32:213–5.
- Utsunomiya H, Tilakaratne WM, Oshiro K, Maruyama S, Suzuki M, Ida Yonemochi H, Cheng J, Saku T. Extracellular matrix remodeling in oral submucous fibrosis: its stage-specific modes revealed by immunohistochemistry and in situ hybridization. *Journal of oral pathology & medicine.* 2005 Sep;34(8):498–507.
- Scutt A, Meghji S, Canniff JP, Harvey W. Stabilisation of collagen by betel nut polyphenols as a mechanism in oral submucous fibrosis. *Experientia*1987;43:391–3.
- Anuradha CD, Devi CS. Serum protein, ascorbic acid & iron & tissue collagen in oral submucous fibrosis—a preliminary study. *The Indian Journal of Medical Research.* 1993 Jun 1;98:147–51.
- Xu H, Lyu F yuan, Song J yuan, Xu Y ming, Jiang E hui, Shang ZJ, et al. Research Achievements of Oral Submucous Fibrosis: Progress and Prospect. *BioMed Res Int.* 2021 Mar 18;2021:6631856.
- More CB, Gavli N, Chen Y, Rao NR. A novel clinical protocol for therapeutic intervention in oral submucous fibrosis: an evidence based approach. *Journal of oral and maxillofacial pathology: JOMFP.* 2018 Sep;22(3):382.
- Isaac U, Issac JS, Ahmed Khoso N. Histopathologic features of oral submucous fibrosis: a study of 35 biopsy specimens. *Oral Surg Oral Med Oral Pathol Oral Radiol Endodontology.* 2008 Oct;106(4):556–60.
- Dhanvantri. An exfoliative cytology study to assess oral mucosa of oral submucous fibrosis patients in Chennai, Tamil Nadu: A preliminary study [Internet]. [cited 2023 Jun 9].
- Keshav, R.; Narayanappa, U. Expression of proliferating cell nuclear antigen (PCNA) in oral submucous fibrosis: An immunohistochemical study. *J. Clin. Diagn. Res.* 2015, 9, ZC20.
- Mukul SK, Kumar S, Pandey S, Mokhtar EA, Kumar A. Ultrasound elastography as a potential diagnostic aid in oral submucous fibrosis. *Natl J Maxillofac Surg.* 2019;10(2):129–33.
- Vedswari, C.P.; Jayachandran, S.; Ganesan, S. In vivo autofluorescence characteristics of pre- and post-treated oral submucous fibrosis: A pilot study. *Indian J. Dent. Res.* 2009, 20, 261–267.
- Chole RH, Gondivkar SM, Gadail AR, et al. Review of drug treatment of oral submucous fibrosis. *Oral Oncol*2012;48:393–8.
- Borle RM, Borle SR. Management of oral submucous fibrosis: a conservative approach. *J Oral MaxillofacSurg Off J Am Assoc Oral Maxillofac Surg.* 1991 Aug;49(8):788–91.
- Haue MF, Meghji S, Nazir R, Harris M. Interferon gamma may reverse oral submucous fibrosis. *J Oral Pathol Med*2001;30:12–21.
- Krishnamoorthy B, Khan M. Management of oral submucous fibrosis by two different drug regimens: A comparative study. *Dent Res J (Isfahan)*2013; 10(4):327–32.
- Srivastava A, Agarwal R, Chaturvedi T, Chandra A, Singh O. Clinical evaluation of the role of tulsi and turmeric in the management of oral submucous fibrosis: A pilot, prospective observational study. *J Ayurveda Integr Med.* 2015 Apr 16;6:45–9.
- A Sarwar A Iqbal KY Giri Efficacy of aloe vera gel as an adjuvant treatment of oral submucous fibrosis. *Oral Surg Oral Med Oral Pathol Oral Radiol*2013;116:71724.
- A Kumar ABagewadi V Keluskar M Singh Efficacy of lycopene in the management of oral submucous fibrosis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endodontology* 2007;103:2071310.1016/j.tripleo.2006.07.011.
- Nagaraj T, Santosh HN. Estimation of serum hepcidin in oral submucous fibrosis before and after supplementation with oral iron: A randomized control clinical trial. *Journal of Oral and Maxillofacial Pathology: JOMFP.* 2018 Sep;22(3):303.
- Kerr AR, Warnakulasuriya S, Mighell AJ, Dietrich T, Nasser M, Rimal J, et al. A systematic review of medical interventions for oral submucous fibrosis and future research opportunities. *Oral Dis*2011; 17 Suppl 1:42–57.
- Sadat A, ArefinMds, Nusrath M, Rita S. Management of Oral Sub-Mucous Fibrosis: An update. *2020 May 12;38:135–44.*
- Gn S, Arora M, Lakhanpal M. Stem cell therapy: A novel treatment approach for oral mucosal lesions. *J Pharm Bioallied Sci.* 2015 Feb 24;7:2–8.