



A PROSPECTIVE OBSERVATIONAL COMPARATIVE STUDY BETWEEN TOTAL LAPAROSCOPIC HYSTERECTOMY AND VAGINAL HYSTERECTOMY

Health Science

Dr Annesha Dutta* Department of Obstetrics & Gynaecology, District Hospital Howrah, West Bengal
*Corresponding Author

Prof Murari Mohan Koley Department of Obstetrics & Gynaecology, District Hospital Howrah, West Bengal

Dr Kavitha Narayanaswamy Department of Obstetrics & Gynaecology, Guntoor Medical College, Andhra Pradesh

ABSTRACT

Total laparoscopic hysterectomy (TLH) is now emerging as a safe procedure even in patients suitable for vaginal hysterectomy (VH) due to its advantages like better visualisation, less post-operative pain and shorter hospital stay. This study was done to compare the duration of surgery, intra-operative and postoperative complications, hospital stay and post-operative analgesia requirement in TLH and VH. **Methods:** The Prospective observational comparative study between women undergoing TLH and VH was carried out between September 2020 to September 2021. Patients with suspected genital malignancy and uterine prolapse were excluded. Baseline characteristics like age, BMI, parity, indication for hysterectomy, uterine size and previous pelvic surgeries were noted. Intra-operative and post-operative parameters like duration of surgery, complications, post-operative analgesic dosage and hospital stay were compared between the two surgeries and the results were analysed using Chi square test and independent t test. **Results:** The mean time taken to perform TLH was significantly longer, i.e. 150.89 minutes compared with VH, i.e. 70.44 minutes (p value 0.002). But the duration of stay in the hospital was shorter for the women undergoing TLH, mean duration being 2.75 days as opposed to 5.34 days in women undergoing VH (p value 0.003). Also, women undergoing VH required more analgesic doses (mean 2.89) than those undergoing TLH (mean 2.00). When we studied the rate of complications in both the groups, we found no statistically significant difference. **Conclusions:** TLH was as safe as VH and had advantages like shorter hospital stay and reduced analgesia dose.

KEYWORDS

Total laparoscopic hysterectomy, Vaginal hysterectomy, Hospital stay, Post-operative analgesia

INTRODUCTION

Hysterectomy is one of the most frequently performed surgical procedures in gynaecology(1).It can be performed vaginally, abdominally, or with laparoscopic or robotic assistance. Factors that may influence the route of hysterectomy include the indication for surgery, size of the uterus, presence or absence of associated pelvic pathology, surgeon's training and preference and patient's choice.(2) Evidence demonstrates that, in general, vaginal hysterectomy is associated with better outcomes and views, allows performance of concomitant surgery, and is suitable for larger uteri and those with non – descent uterus, which may prove difficult to remove vaginally.(3) This study was done to compare laparoscopic hysterectomy with vaginal hysterectomy in women with benign disorders and to see if laparoscopic hysterectomy had advantages over vaginal hysterectomy.

Methods

This prospective observational comparative study was carried out during the period between September 2020 to September 2021.Thirty patients who underwent total laparoscopic hysterectomy (TLH) were included in the TLH group and twenty patients who underwent vaginal hysterectomy (VH) were included in the VH group. The choice of surgery was made by the patients after discussion with their consultants.

Exclusion criteria were suspicion of malignancy and a vaginal prolapse higher than first degree. All the patients were evaluated for fitness for surgery. The vaginal hysterectomy was performed following Heaney's technique. The laparoscopic technique was always a total laparoscopic hysterectomy. For each patient we recorded the baseline characteristics, including age; parity; previous caesarean section; other previous pelvic surgery; body mass index (BMI); uterine size and the indication for hysterectomy. Intra-operative parameters including conversion to laparotomy, time of surgery, complications – injury to bowel, bladder, ureter and postoperative parameters including hospital stay, infections, vault haematoma, DVT/pulmonary embolism, vesicovaginal fistula and haemorrhage, if any, were recorded. Analgesic doses on the day of surgery were also recorded. Prophylactic antibiotic was given to all patients at the beginning of the surgery and repeated 12 hours later. The data were analyzed using independent T test and chi square test. A p value of <0.05 was considered to be statistically significant.

RESULTS

The mean age of women undergoing TLH was 48.45 years and for those in VH group it was 50.11 years and the difference was statistically significant. Women undergoing TLH had a higher BMI,

i.e., 24.91 kg/m2 compared with women undergoing VH, i.e., 27.67 kg/m2 and this difference was statistically significant. However, neither age nor BMI had any impact on the parameters compared in this study. The difference in the size of the uterus between the two groups was also statistically significant. In the TLH group, the mean size of the uterus was 14.28 weeks whereas in VH group it was 10.22 weeks (Table 1)

Table 1: Baseline Characteristics: Age, BMI, and Uterine Size

GROUP	TLH			VH			P Value
	Mean	Standard Deviation	Median	Mean	Standard deviation	Median	
Age (Years)	48.45	8.52	49.89	50.11	7.80	51.18	0.005
BMI (kg/m2)	24.91	6.37	23.11	27.67	5.67	27.45	0.018
Uterine Size (weeks)	14.28	6.84	12.01	10.22	5.37	10.04	0.003

There was no statistically significant difference in parity between the groups. There were more number of women with previous caesarean in TLH group compared to VH group but the difference was not statistically significant. Women in both the groups had undergone previous pelvic surgeries other than caesarean section (Table 2). The mean time taken to perform TLH was significantly longer, i.e. 150 minutes compared with VH, i.e. 80 minutes, with the p value being <0.0001.

Table 2: Baseline Characteristics: Parity, Previous Caesarean and Presence of Previous Pelvic Surgery.

GROUP		TLH(25)		VH(20)		P Value
		N	%	N	%	
Para	Nulliparous	10	40%	8	40%	0.76
	Multiparous	15	60%	12	60%	0.81
Previous LSCS	Yes	8	32%	4	20%	0.23
	No	17	68%	16	80%	0.45
Presence of previous pelvic Surgery	Yes	12	48%	6	30%	0.89
	No	13	52%	14	70%	0.67

But the duration of stay in the hospital was shorter for the women undergoing TLH, mean duration being 2.75 days as opposed to 5.34 days in women undergoing VH. This difference was statistically significant with p value being 0.003. Also women undergoing VH required more analgesic doses (mean 2.89) than those undergoing TLH (mean 1.45) and this difference was also statistically significant (Table 3).

Table 3: Intra-operative and Post-operative Parameters like Operation Time, Hospital Stay and Analgesia Dose Required.

GROUP	TLH			VH			P Value
	Mean	Standard Deviation	Median	Mean	Standard Deviation	Median	
Operation Time (Hours)	150.89	50.34	137.89	70.44	30.44	65.34	0.002
Hospital Stay (days)	2.75	1.67	3.07	5.34	2.43	4.45	0.003
Analgesia Dosage	2.00	1.45	2.18	2.89	1.45	2.81	0.0003

When we studied the rate of complications in both the groups, we found no statistically significant differences. Two woman undergoing VH had bladder injury which was recognised intraoperatively and repaired. Two woman had vault haematoma in post-operative period and one woman had vesico-vaginal fistula which were also managed conservatively. Three woman in TLH group came to us with vesicovaginal fistula on post-operative day 10, 14 and 16 and those were repaired by laparoscopy. Conversion to laparotomy was not done in any case in both the groups.

Table 4: Intra-operative and Post-operative Complications and Conversion to Laparotomy.

GROUP		TLH(25)		VH(20)		P Value
		n	%	n	%	
Intraoperative complications	Nil	25	100	18	90	0.58
	Visceral Injury	0	0	2	10	0.88
Conversion to laparotomy	Nil	25	100	20	100	0.49
Post operative complications	Infection	0	0	0	0	Not applicable
	Vault hematoma	0	0	2	10	Not Applicable
	Vesicovaginal fistula	3	12	1	5	Not applicable

When the time taken for surgery was calculated, we found an increase in the operative time for TLH compared with VH. This may be due to the increase in the size of the uterus in TLH group compared with VH group. Other studies comparing TLH with VH have also shown an increase in operative time in the TLH group compared to vaginal route. However, women in TLH group stayed in the hospital for a significantly lesser duration of time compared to those undergoing VH. In the study done by V DaCosta et al they found that although not significantly different, patients in TLH group tended to spend on an average a shorter duration of time in hospital.

In the study done by Mathew Morton et al,² they found that the incidence of intraoperative complications in both the TLH and VH groups appeared to be similar, especially in patients with a uterine mass of less than 300 g. Niel Johnson et al,¹⁵ in their meta-analysis of randomised control trials of methods of hysterectomy found no significant differences in urinary tract injury for laparoscopic versus vaginal hysterectomy. Also, no significant differences were noted in long term complications like fistula formation. Even in our study the complications were rare and there was no significant difference in the rate of complications between the two groups. However, the sample size was not sufficient to analyse the incidence of complications.

There was no conversion to laparotomy in our study in both the TLH group and VH group. In the study done by Matthew Morton et al, one case in TLH group was converted to laparotomy. In the study done by V DaCosta et al,³ none of the cases were converted to abdominal route.

DISCUSSION

In contemporary gynecological practice, it is recommended that,

where possible, hysterectomy for benign indications should be approached vaginally⁽⁴⁾. But the studies which said that VH is superior to TLH, like the eVALuate study and the Cochrane review, had their own limitations. In the eVALuate study, the trial comparing vaginal hysterectomy with laparoscopic hysterectomy was underpowered and was inconclusive on the rate of major complications but it stated that vaginal hysterectomy took less time⁽⁵⁾.

In the Cochrane review, no advantages of laparoscopic hysterectomy over vaginal hysterectomy could be found. Laparoscopic hysterectomies had a greater risk of damaging the bladder or ureter. Vaginal hysterectomy should be performed whenever possible. Where vaginal hysterectomy is not possible, both a laparoscopic approach and abdominal hysterectomy have their pros and cons and these should be incorporated in the decision-making process. However, the evidence in this review has to be interpreted with caution as adverse event rates were low, resulting in low power for these comparisons⁽⁶⁾. From a technical point of view, laparoscopic approach benefits include high-definition imaging and vision amplification, allowing for an easier access to the uterine vessels, ureter, rectum and vagina⁽⁷⁾. Despite the initial controversy regarding laparoscopic approach-related complication rate, most recent studies are consensual and present TLH as a safe and first-line technique in multiple surgical scenarios, either benign or malignant^(8, 9, 10).

In our study we found that the TLH group had lower BMI and larger sized uteri which was statistically significant. There was a statistically significant difference in the number of analgesic dosages used in our study, with lesser number of analgesics given in TLH group compared with VH group. This reflects less post-operative pain in TLH group compared with VH group. The results of various other studies comparing TLH and VH are similar^(11, 12). Raxita Patel et al¹¹ in their study found a statistically significant difference in pain scores between TLH and VH with patients in TLH having significantly lower pain scores. TLH may be performed in a wide variety of patients, including those with a large uterus, which would ordinarily require an abdominal approach for hysterectomy. It is also preferable for nulliparous patients or for those with limited vaginal access and the resulting challenges of VH. It facilitates removal of adnexa, either normal or pathological. Also, urinary bladder can be clearly visualised and adhesions can be identified and divided in cases of previous caesarean section.^(13,14)

CONCLUSION

From a technical point of view, laparoscopic approach benefits include high-definition imaging and vision amplification, allowing for an easier access to the uterine vessels, ureter, rectum and vagina. Despite the initial controversy regarding laparoscopic approach-related complication rate, most recent studies are consensual and present TLH as a safe and first-line technique in multiple surgical scenarios, either benign or malignant.

The limitations of this study are lack of randomisation and decreased sample size to study the rate of complications in the two groups. The financial implications of TLH vs. VH, as well as long-term differences in post-operative quality of life, urinary incontinence, and sexual function, require further study with larger sample sizes.

Funding: No funding sources

Conflict of Interest: None declared Ethical approval: Not required

REFERENCES

1. Cristina Nogueira-Silva, Samuel Santos-Ribeiro, Sônia Barata, Conceição Alho, Filipa Osorio, Carlos Calhaz-Jorge Total Laparoscopic hysterectomy: Retrospective analysis of 262 cases Acta Med Port 2014 Jan-Feb;27(1):73-81.
2. Morton M, Cheung VY, Rosenthal DM. Total laparoscopic versus vaginal hysterectomy: a retrospective comparison. J Obstet Gynaecol Can 2008;30:1039-44.
3. V Dacosta, A McIntosh et al. Total laparoscopic hysterectomy versus vaginal hysterectomy at the university hospital of the West Indies: a 5 year retrospective study. West Indian Medical Journal. Vol 61; No 9 Mona dez.2012.
4. Falcone T, Walters M. Hysterectomy for benign disease. Obstet Gynecol 2008;111:753-67.
5. Garry R, Fountain J, Mason S, Have J, Napp V, Abbott J et al. The eVALuate study: two parallel randomised trials, one comparing laparoscopic with abdominal hysterectomy, the other comparing laparoscopic with vaginal hysterectomy. BMJ 2004;328:129.
6. Aarts JWM, Nieboer TE, Johnson N, Tavender E, Garry R, Mol BJ, Kluivers KB . Surgical approach to hysterectomy for benign gynaecological diseases. Cochrane review 1 September 2015.
7. Donnez O, Jadoul P, Squiffet J, Donnez J. A series of 3190 laparoscopic hysterectomies for benign disease from 1990 to 2006: evaluation of complications compared with vaginal and abdominal procedures. BJOG. 2009; 116:492-500.
8. Donnez O, Donnez J. A series of 400 laparoscopic hysterectomies for benign disease: a single centre, single surgeon prospective study of complications confirming previous retrospective study. BJOG.2010;117:752-5.
9. Karaman Y, Bingol B, Günenç Z. Prevention of complications in laparoscopic

- hysterectomy: experience with 1120 cases performed by a single surgeon. *J Minim Invasive Gynecol.* 2007;14:78-84.
10. Bojahr B, Raatz D, Schonleber G, Abri C, Ohlinger R. Perioperative complication rate in 1706 patients after a standardized laparoscopic supracervical hysterectomy technique. *J Minim Invasive Gynecol.* 2006;13:183-9.
 11. Schindlbeck C, Klausner K, Dian D, Janni W, Friese K. Comparison of total laparoscopic, vaginal and abdominal hysterectomy. *Arch Gynecol Obstet* 2008;277:331-7.
 12. Kim HB, Song JE, Kim GH, Cho HY, Lee KY. Comparison of clinical effects between total vaginal hysterectomy and total laparoscopic hysterectomy on large uteruses over 300 grams. *J Obstet Gynaecol Res* 2010;36: 656-60.
 13. Candiani M, Izzo S, Bulfoni A, Riparini J, Ronzoni S, Marconi A. Laparoscopic vs. vaginal hysterectomy for benign pathology. *Am J Obstet Gynecol* 2009;200:368.e1-7.
 14. Rasha G, Colin A. Walsh, Stewart R. Walsh, Emmanuel Karantanis. Vaginal hysterectomy versus total laparoscopic hysterectomy for benign disease: a metaanalysis of randomized controlled trials. *Am J Obstet Gynecol.* 2011;204(5):388.e1-388.e8