



THE DIAGNOSTIC VALUE OF THE RELATIONSHIP BETWEEN LIPID PROFILE, THE DIFFERENT GRADES OF CORONARY CALCIFICATION AND STENOSIS IN PATIENTS WITH POSSIBLE CAD

Biochemistry & Radiology

Dr. Subodh Kumar Goel Assistant Professor, Department of Biochemistry, Rama Medical College & Research Centre, Hapur

Dr. Vandana Goel* Associate Professor, Department of Radiology, NCR Institute of Medical Sciences, Meerut *Corresponding Author

ABSTRACT

Background: Coronary calcification is an important risk factor for adverse outcomes in the general population (1, 2). Coronary calcification causes reduction in the vascular compliance (3). Prediction of the severity of CAD is valuable due to the increased risk for cardiovascular events CT scan of the coronary arteries allows assessment of the degree of coronary calcification as well as the degree of stenosis and the type of plaque (calcific or non-calcific) (6). **Objectives:** To investigate the relationship between the lipid profile parameters and the different grades of coronary calcification, using MSCT and correlate them with the significance of coronary stenosis during coronary angiography in patients suspected to have coronary artery disease. **Subjects and Methods:** Our study included 300 adult symptomatic patients, in the period between January 2017 and February 2018. Lipid profile for each patient was registered, calcium score was calculated for each patient, including areas with a density above 130 HU using (Agatston score), we divided the study population into 2 categories (one group having calcium score above 400 and the other group having calcium score below 400) for a better correlation. After calculating the calcium score, coronary catheterization was done for the assessment of the significance of coronary stenosis, significant lesions were defined as greater than 70% reduction in the luminal diameter, the time interval between catheterization and CT scan was not more than one week. The correlation between the lipid profile parameters and the degree of coronary calcification as well as the significance of coronary stenosis was established. **Results:** Serum cholesterol, LDL and triglycerides had a strong positive correlation with the total calcium score, whereas HDL had a strong negative correlation with the total calcium score (p value <0.001). Serum cholesterol, LDL and Triglycerides were also significantly higher in the group of patients having significant lesions (p value <0.001), whereas HDL level was significantly lower in the group of patients having significant stenosis (p value <0.001). Age was significantly higher in the group of high calcium score (agatston score >400), the number of hypertensive and diabetic patients was higher in the group of patients with high calcium score (agatston score >400), the number of patients with a positive family history of ischemic heart diseases was higher in the group of high calcium score (agatston score >400). **Conclusion:** dyslipidemia was found to be an important predictor for coronary calcification (detected by CT scan) as well as the presence of significant coronary artery stenosis (detected by coronary angiography)

KEYWORDS

lipid profile, CT scan

INTRODUCTION

Coronary calcification is an important risk factor for adverse events in the general population. Many risk factors have been identified, contributing to the initiation and progression of coronary calcification.

Trials were done to control coronary calcification with medical therapy, but failed. Drug-eluting stents and devices for plaque modification have modestly improved to overcome calcification in the coronaries, but the adverse event rates are still high.

Innovative pharmacologic and device-based approaches are needed to improve the poor prognosis of patients with extensive coronary calcification.

Coronary calcification reduces the vascular compliance, vasomotor reaction, with impaired myocardial perfusion and is associated with worse outcomes in the general population⁽⁵⁾.

Methods for Detection of Coronary Calcification

1- MDCT: is the only noninvasive test with high sensitivity and specificity for calcium detection, capable of quantifying calcium score⁽¹⁰⁾. In large-scale observational studies, CT-scan- based calcium scores added prognostic value for predicting cardiac death and myocardial infarction⁽¹¹⁾.

Noninvasive assessment of calcium score is reasonable for cardiovascular risk assessment in asymptomatic patients with intermediate pretest probability for CAD⁽¹²⁾.

A calcium score of >0 suggests some underlying atherosclerosis, whereas scores ≥ 400 should prompt risk factor modification and further diagnostic evaluation⁽¹⁵⁾. Asymptomatic persons without traditional risk factors but with a documented calcium score ≥ 400 might have a worse cardiovascular prognosis than those with ≥ 3 risk factors but no CT-detected coronary calcification⁽¹⁵⁾.

2- Coronary Angiography: has a low to moderate sensitivity compared with gray-scale intravascular ultrasound (IVUS) and CT for detection of coronary calcification, but is very specific. Angiographic detection of coronary calcification is often classified into 3 groups:

none/mild, moderate, and severe. Severe calcification is most commonly defined as radio-opacities seen without cardiac motion before contrast injection, usually affecting both sides of the arterial lumen, and moderate calcification as radio-opacities noted only during the cardiac cycle before contrast injection⁽⁴⁸⁾.

3- Intravascular Ultrasound: is more accurate than cine-angiography for calcium detection, with sensitivity of 90% to 100% and specificity of 99% to 100%⁽¹⁷⁾. The signature of calcified plaque on grayscale IVUS is a bright echo with acoustic shadowing⁽¹⁸⁾, and the extent of calcification can be graded by several metrics. The arc of calcium is classified as none, 1 quadrant (0° to 90°), 2 quadrants (91° to 180°), 3 quadrants (181° to 270°), or 4 quadrants (271° to 360°). Calcium location is defined as superficial if present in the intimal-luminal interface, deep if within the medial- adventitial border or closer to the adventitia than the lumen, or both superficial and deep. Deposits can be assessed relative to the thickest plaque accumulation as concordant (center of calcium arc $\leq 45^\circ$ of thickest plaque accumulation), perpendicular (center of calcium arc 45° to 135° from thickest plaque accumulation), or opposite (center of arc of calcium $\geq 135^\circ$ from thickest plaque accumulation). Finally, the calcium length can be measured⁽¹⁹⁾. However, because ultrasound does not penetrate calcium, calcium thickness cannot be determined, and volume cannot be calculated.

Radiofrequency analysis of the IVUS signal allows for in vivo characterization of coronary plaque components, including fibrotic plaque, fibro-fatty plaque, necrotic core, and dense calcium⁽²⁰⁾.

1- Optical Coherence Tomography (OCT): provides higher-resolution imaging (10 to 20 μm) than grayscale IVUS (150 to 200 μm). Optical coherence tomography detects calcium as low- intensity, low-attenuation areas with sharp borders. The sensitivity (95% to 100%) and specificity (97% to 100%) of OCT for coronary calcification⁽²¹⁾. Moreover, because light penetrates calcium, OCT can in many cases assess calcium thickness and measure calcium volume⁽²¹⁾.

Methods for Ssessment of the Degree of Coronary Stenosis.

1- Coronary Angiography

A coronary angiogram shows the geometry of the lumen of the vessels,

but not the vessel wall and plaque⁽²²⁾. However, it provides limited information about the surface of the atherosclerotic plaque either smooth or irregular and calcifications of the vessel wall. Several studies described that the majority of culprit lesions in patients with unstable angina had a characteristic angiographic appearance. These lesions have been termed Ambrose type II lesions. However, these characteristics describe the plaques that have already ruptured, rather than those at risk to rupture.⁽²³⁾

The extent of fluoroscopic calcifications is a marker of the overall atherosclerotic disease burden and, therefore, has a prognostic value.⁽²⁴⁾

2-CT Angiography:

Contrast-enhanced CT scans of the coronary arteries allow the visualization of non-calcified atherosclerotic plaque in lesions that are not associated with a significant stenosis. Detection, quantification, and characterization of coronary plaque by CT are currently the subjects of intense evaluation, helping in the risk stratification of the studied population. CT measurements can further characterize the different types of non-calcified plaque, because lipid-rich plaque is, on average, associated with lower Hounsfield attenuation numbers than fibrous plaques (14 to 72 HU for lipid-rich plaques and 90 to 121 HU for fibrous plaques)⁽²⁵⁾

Whereas, coronary calcification can be reliably detected by CT at a relatively low radiation dose, and coronary calcium has clearly been shown to be associated with an increased risk of future coronary artery disease events in asymptomatic individuals, available clinical data are rare for the detection of non-calcified plaque by contrast-enhanced CT.

The Predictive value of MDCT for the presence of significantly obstructive coronary artery lesions is around 80% in most comparative studies⁽²⁹⁾. MDCT does not render "projections" but "sections" of the coronary arteries with any angulation and, in a retrospective way, after the whole cardiac volume has been acquired.

Images obtained from MDCT may include cross-sectional views of the vessels. Which are not available in angiographic studies, this allows a more detailed analysis of the lesions⁽³⁰⁾.

3-Intravascular Ultrasound (IVUS):

Intravascular ultrasound provides imaging of the vessel wall, plaque size, and plaque morphology.⁽³²⁾

During cardiac catheterization, a miniature ultrasound catheter is placed distal to the target lesion site, and then withdrawn during continuous imaging, resulting in a series of cross-section images. The vessel wall of each cross-section can be described by the signal characteristics on a continuum from echo-dense (bright echo signal) to echo-lucent (faint echo signal). The measurement of lumen area, intima-media area, and external elastic membrane (EEM) allows assessment of the state of arterial remodeling and plaque burden. Several IVUS studies have demonstrated the effectiveness of ultrasound imaging in predicting the composition of the atherosclerotic plaque when compared with histology.⁽³³⁾

Advances in data acquisition and analysis in intravascular ultrasound information allowed a more detailed description of vessel wall and plaque morphology.⁽³⁴⁾ In particular, radio-frequency analysis has been used for the characterization of plaque structures such as the lipid pool and fibrous cap.⁽³⁴⁾

Subjects and Methods

Non-contrast enhanced CT which was done 24hrs to 1 week before coronary angiography depending upon the patient risk profile and clinical presentation

All participants underwent measurement of coronary calcium using Multi-Detector CT (MDCT) at baseline using Agatston score. Tissue densities ≥ 130 Hounsfield units were set as the attenuation level for a calcified plaque.

We used a dual source scanner (Somatom Definition Flash) using two X-ray sources which would yield better visualization of the coronary tree, ECG triggering was set to 75% of the RR interval.

Agatston score was used to calculate the sum of calcification in all coronaries as well as the calcium score in each vessel separately after reconstructing slices of 3 mm in thickness, beginning 1 cm below the

carina and progressing caudally to include the proximal coronary arteries. The coronary calcification was calculated based on the area and density for each of the major coronary arteries: LMT, LAD, LCX, and RCA

The images were evaluated by four experienced observers, who had no knowledge of the clinical data. Two contiguous pixels with an attenuation coefficient >130 Hounsfield units were defined as a calcium deposit. The calcification was scored on a vessel by vessel and a total score was obtained for all vessels

All studies performed were scored independently by the observers to assess inter-observer variations

The collected data were analyzed on dedicated software platforms.

Routine laboratory investigations with special concern to:

Fasting Blood sugar level, Fasting plasma lipid profile including (Total Cholesterol, Triglycerides, HDL, LDL) to assess the interaction between the cardiovascular risk factors and the degree of coronary calcification as well as the significance of coronary stenosis.

Coronary angiography was performed in different projections after discussing the risk-benefit ratio of the procedure and obtaining signed informed consent, all patients were brought to the cardiac catheterization laboratory. Mild conscious sedation was commonly used, and Oxygen saturation was constantly monitored. The vascular access site was cleansed with an antibacterial solution and covered with surgical drapes.

A percutaneous radial artery approach was occasionally used in patients with severe peripheral vascular disease or when the groin approach was not possible or technically difficult.

Coronary angiograms were obtained with the use of pre shaped (usually Judkins, but occasionally Amplatz, multipurpose or other) 6 French diameter diagnostic catheters. After engaging the ostium of the coronary artery it was confirmed that the tip of the catheter was free, and not against the wall of the artery, under a sub-intimal plaque. This was accomplished by confirming that the pressure was not damping and with the use of a test injection using a small amount of contrast material⁽³⁵⁾.

Various views or projections of the coronary angiograms were obtained by rotating the x-ray video camera around the patient.

The cinegrams were interpreted by 2 experienced cardiologists who had no knowledge about the CT findings

The vessels were subdivided anatomically as described for CT and the condition of each vessel was described by one of the following categories, mild ($<50\%$ luminal stenosis), borderline (50-70% luminal stenosis) and significant ($>70\%$ luminal stenosis), diseased vessels were defined as those having luminal stenosis $>50\%$, and significantly stenotic vessels were defined as those having $>70\%$ luminal narrowing

The Analysis of Data Included:

- Initial assessment of the lipid profile parameters and quantification of the amount of calcification by agatston score using a non-contrast enhanced CT scan.
- The included subjects were then categorized into two groups for better data analysis, the first group was for those with calcium score less than 400 (low calcium score) and the other group was for those with calcium score more than 400 (high calcium score group).
- The severity or degree of stenosis was measured angiographically by comparing the area of narrowing to an adjacent normal segment, and as a percentage reduction and calculated in the projection which demonstrates the most severe narrowing⁽³⁵⁾.
- The patient was considered to have significant stenosis, if ≥ 1 plaque is causing $>70\%$ luminal narrowing. Analysis of the data obtained by correlating the lipid profile parameters, calcium score and the presence of significant stenosis, using specific statistical programs.

Data were analyzed using Statistical Program for Social Science (SPSS). Quantitative data were expressed as mean $\pm$ standard deviation (SD). Qualitative data were expressed as frequency and percentage.

The Following Tests were used:

- Independent-samples t-test of significance was used when comparing between two means.
- Chi-square (X2) test of significance was used in order to compare proportions between two qualitative parameters.
- Pearson's correlation coefficient (r) test was used for correlating data.
- The confidence interval was set to 95% and the margin of error accepted was set to 5%. So, the p-value was considered significant as the following:
 - Probability (P-value)
 - P-value <0.05 was considered significant.
 - P-value <0.001 was considered as highly significant.
 - P-value >0.05 was considered insignificant.

RESULTS

Patient Demographics

The clinical characteristics of patients are summarized in (Table 1), the total number of the studied population was 300 patients, the mean age of the study group was (55.40±9.59), 86.7% were males and 13.3% were females, 59% were hypertensive, 47.7% were diabetic, 48.7% were smokers, 5.7% had a family history of ischemic heart diseases.

Table (1): Baseline Characteristic data of the study group.

Characteristic data	Total (N=300)
gender	
Female	40 (13.3%)
Male	260 (86.7%)
Age (years)	31-78 (55.40±9.59)
HTN	177 (59.0%)
DM	143 (47.7%)
Smoking	146 (48.7%)
Family history of ischemic heart disease	17 (5.7%)

The mean age of the study group was (55.40±9.59), 86.7% were males and 13.3% were females, 177 (59%) patients were hypertensive, 143 (47.7%) patients were diabetic, 146 (48.7%) patients were smokers, 17 (5.7%) patients had a family history of ischemic heart disease

Lipid profile parameters of the studied population were summarized in (table 2), the mean total cholesterol was 242.92 ± 43.25, mean LDL cholesterol level was 153.12 ± 31.01, mean HDL cholesterol was 50.91±17.04, the mean serum triglycerides level was 197.88±51.18.

Table (2): Lipid profile parameters of the patients included in the study.

Lipid Profile	Total (N=300)
S. Cholesterol	155-350 (242.92±43.25)
LDL	89-230 (153.12±31.01)
HDL	26-159 (50.91±17.04)
S. Triglycerides	90-340 (197.88±51.18)

LDL=Low density lipoprotein, HDL = High density lipoprotein, SD=Standard deviation.

The clinical data of the patients are also characterized in both groups of calcium score (the high calcium score group and the low calcium score group), summarized in (table 3). Age was significantly correlated with the total calcium score (fig1) and the higher age group of patients was markedly distributed in the group of high calcium score (p value <0.001)(fig2), the number of hypertensive and diabetic patients was significantly higher in the group of higher calcium score, family history of ischemic heart diseases was strongly positive in the group of higher calcium score (p value 0.011) (fig3). Sex difference and smoking had no significant difference between both calcium score groups.

Table (3): Comparison between the effect of demographic data on Calcium score in both groups <400 and >400 according to baseline data.

Characteristic data	Ca score <400 (N=200)	Ca score >400 (N=100)	t/x2#	p-value
Age (years)			67.103	<0.001
Mean±SD	52.50±8.42	61.21±9.19		
Range	31-75	40-78		
Sex			0.014#	0.904
Female	27 (13.50%)	13 (13.00%)		
Male	173 (86.50%)	87 (87.00%)		

HTN	103 (51.50%)	74 (74.00%)	13.952#	<0.001
DM	84 (42.0%)	59 (59.00%)	7.057#	0.008
Smoking	96 (48.00%)	50 (50.00%)	0.107#	0.744
Family history	6 (3.00%)	11 (11.00%)	6.555#	0.011

Age was significantly higher in the calcium score group >400 (p value <0.001), the number of hypertensive and diabetic patients was significantly higher in the calcium score group >400, family history of ischemic heart diseases was strongly positive in the calcium score group >400 (p value 0.011). Sex difference and smoking had no significant difference in their effect between both groups.




Figure (1): Scatter plot between Ca score and age (years) showing a significant positive correlation between age and the total calcium score.




Figure (2): Bar chart showing the difference between both calcium score groups as regarding the age of the included patients, a significant distribution of the advanced age patients in the calcium score group >400.



Figure (3): Bar chart showing the distribution of the different risk factors between both calcium score groups.

Calcium score distribution of the study group was summarized in (table 4); calcium score in the study group was ranging from 45-4000.

200 cases had low calcium score (calcium score <400) and 100 cases had high calcium score (calcium score >400)

Table (4): Ca Score Distribution of the Study Group.

Ca score	Total (N=300)
<400	200 (66.7%)
>400	100 (33.3%)
Range [Mean±SD]	45-4000 (447.16±406.80)

This table shows that 200 cases were distributed in the subgroup <400 Calcium score and 100 cases were distributed in the subgroup >400 Calcium score, calcium score in the study group was ranging from 45-400. The relationship between the different lipid profile parameters and the total calcium score was summarized in (table 5), there was a significant positive correlation between the total cholesterol level and the total calcium score (fig4), a significant positive correlation was detected between LDL and the total calcium score (fig5), a significant negative correlation between HDL and the total calcium score (fig 6), a significant positive correlation was detected between triglycerides and

the total calcium score (fig7)

Table (5): Comparison Between Both Calcium Score Groups (<400 and >400) According to Lipid Profile Parameters

Lipid Profile	Ca score <400 (N=200)	Ca score >400 (N=100)	t-test	p-value
S. Cholesterol			108.595	<0.001
Mean±SD	227.15±38.33	274.48±34.45		
Range	155-350	190-340		
LDL			126.286	<0.001
Mean±SD	141.18±27.26	177.01±23.38		
Range	89-210	120-230		
HDL			94.725	<0.001
Mean±SD	56.82±17.63	39.09±6.39		
Range	39-159	26-61		
S. Triglycerides			216.974	<0.001
Mean±SD	174.43±35.37	244.78±45.41		
Range	90-320	145-340		

Total cholesterol, LDL cholesterol, and serum triglycerides were significantly higher in the group of high calcium score (> 400) than the group of lower calcium score (< 400) (P value < 0.001). HDL was significantly lower in the group of high calcium score (>400), P value < 0.001

LDL = Low density lipoprotein, HDL = High density lipoprotein




Figure (4):Scatter plot between Ca score and S. cholesterol showing a significant positive correlation.




Figure (5): Scatter plot between Ca score and LDL showing a significant positive correlation




Figure (6): Scatter plot between Ca score and HDL showing a significant negative correlation




Figure (7): Scatter plot between Ca score and S. Triglycerides showing a significant positive correlation.

The relationship between the different lipid profile parameters and the significance of stenosis was summarized in (table 6, fig8), Serum cholesterol, LDL and Triglycerides were significantly higher in the group of patients having significant coronary stenosis, whereas HDL level was significantly lower in that group.

Table (6): Comparison between non-significant stenosis and significant stenosis according to the lipid profile parameters.

Lipid Profile	Non significant stenosis (N=156)	Significant stenosis (N=144)	t-test	p-value
S. Cholesterol			274.264	<0.001
Mean±SD	214.21±31.47	274.03±31.04		
Range	155-350	189-340		
LDL			593.689	<0.001
Mean±SD	128.85±17.11	179.42±18.83		
Range	89-180	127-230		
HDL			84.325	<0.001
Mean±SD	58.58±19.59	42.59±7.57		
Range	33-159	26-60		
S. Triglycerides			262.848	<0.001
Mean±SD	164.28±27.95	234.28±45.42		
Range	90-284	159-340		

DISCUSSION

High-resolution, contrast enhanced computed tomography of the coronary arteries allows the evaluation not only of the degree of coronary stenosis, but also evaluates coronary plaques, the degree and pattern of calcification.⁽³⁶⁾

The main purpose of this study was to explore the relationship between the lipid profile parameters and the degree of coronary calcification as well as the severity of coronary stenosis.

There was a significant relation between the different lipid profile parameters and the degree of calcification , as a positive correlation was found between LDL , total cholesterol , serum triglycerides and the degree of calcification (P < 0.001) ,Whereas HDL cholesterol had a significant negative correlation with the degree of calcification (P < 0.001),which was supported by the results of Richard A. Kronmal et al,that studied the relationship between dyslipidemia and incident coronary calcification⁽⁴¹⁾.

A significant positive correlation was found between different risk factors including age, hypertension, diabetes, family history of IHD and the presence of as well as the degree coronary calcification; however smoking had a non-significant effect on coronary calcification. In agreement with the previous finding , studies done by Mazzone T et al , N.D. Wong et al , H. Eisenberg, et al who found a significant relation between individual risk factors including age, hypertension, dyslipidemia, diabetes and the presence as well as the degree of coronary calcification^(42,43,44)

We found also that the total cholesterol, LDL, and TG levels were significantly higher in the group of patients who had significant stenosis than in the group of patients having non-significant stenosis (P < 0.001), whereas HDL cholesterol was significantly lower in the group of patients having significant stenosis. This is supported by the results of Shipilova et al.that found a positive linear correlation between the severity of stenosis on one hand, and the total cholesterol,

LDL cholesterol and serum triglycerides on the other hand, negative linear correlation between HDL cholesterol and severity of stenosis was also found.⁽⁴⁵⁾ The study done by Ladeia AM, et al found that Lower HDL-C was associated with more severe and intensive CAD⁽⁴⁶⁾ supporting our findings about the negative correlation between HDL and the severity of CAD

In contrast to our results, a study done by M F Reardon et al. on 65 men and 42 women and noted that only LDL and TGs have the only positive correlation with the stenosis severity, whereas no relation could be detected between HDL and the severity of stenosis, total cholesterol did not account also for changes in stenosis severity in women.⁽⁴⁷⁾

CONCLUSIONS

Dyslipidemia has a strong correlation with the degree of coronary calcification as well as the degree of stenosis, as revealed by multi-detector computed tomography and coronary angiography.

Other risk factors including age, hypertension, and diabetes are strongly correlated with the degree of coronary calcification as well as the degree of stenosis.

MSCT is the best modality for detecting and calculating the degree of coronary artery calcification.

REFERENCES

- 1- S.S. Kalra, C.M. Shanahan Vascular calcification and hypertension: cause and effect *Ann Med*, 44 (Suppl 1) (2012), pp. S85-S92
- 2- R. Vliegenthart, M. Oudkerk, A. Hofman, et al. Coronary calcification improves cardiovascular risk prediction in the elderly *Circulation*, 112 (2005), pp. 572-577
- 3- L. Wang, M. Jerosch-Herold, D.R. Jacobs Jr., et al. Coronary artery calcification and myocardial perfusion in asymptomatic adults: the MESA (Multi-Ethnic Study of Atherosclerosis)
- 4- R. Vliegenthart, M. Oudkerk, A. Hofman, et al. Coronary calcification improves cardiovascular risk prediction in the elderly *Circulation*, 112 (2005), pp. 572-577
- 5- M. Vavuranakis, K. Toutouzas, C. Stefanadis, et al. Stent deployment in calcified lesions: can we overcome calcific restraint with high-pressure balloon inflations? *Catheter Cardiovasc Interv*, 52 (2001), pp. 164-172
- 6- Achenbach S, Moselewski F, Ropers D, et al., Detection of calcified and non calcified coronary atherosclerotic plaque by contrast-enhanced submillimeter multidetector spiral computed tomography: A segment based comparison with intravascular ultrasound, *Circulation* 2004; 109:14-17.
- 7- N.D. Wong, D. Kouwabunpat, A.N. Vo, et al. Coronary calcium and atherosclerosis by ultrafast computed tomography in asymptomatic men and women: relation to age and risk factors *Am Heart J*, 127
- 8- R.L. McClelland, H. Chung, R. Detrano, W. Post, R.A. Kronmal Distribution of coronary artery calcium by race, gender, and age: results from the Multi-Ethnic Study of Atherosclerosis (MESA) *Circulation*, 113 (2006), pp. 30-37
- 9- P. Greenland, L. LaBree, S.P. Azen, T.M. Doherty, R.C. Detrano Coronary artery calcium score combined with Framingham score for risk prediction in asymptomatic individuals *JAMA*, 291 (2004), pp. 210-215
- 10- S.R. Tanenbaum, G.T. Kondos, K.E. Veselik, et al. Detection of calcific deposits in coronary arteries by ultrafast computed tomography and correlation with angiography *Am J Cardiol*, 63 (1989), pp. 870-872
- 11- S.E. Elias-Smale, R.V. Proenca, M.T. Koller, et al. Coronary calcium score improves classification of coronary heart disease risk in the elderly: the Rotterdam study *J Am Coll Cardiol*, 56 (2010), pp. 1407-1414
- 12- P. Greenland, J.S. Alpert, G.A. Beller, et al. 2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines *J Am Coll Cardiol*, 56 (2010), pp. e50-103
- 13- T.S. Polonsky, R.L. McClelland, N.W. Jorgensen, et al. Coronary artery calcium score and risk classification for coronary heart disease prediction *JAMA*, 303 (2010), pp. 1610-1616
- 14- A.S. Agatston, W.R. Janowitz, F.J. Hildner, et al. Quantification of coronary artery calcium using ultrafast computed tomography *J Am Coll Cardiol*, 15 (1990), pp. 827-832
- 15- K. Nasir, J. Rubin, M.J. Blaha, et al. Interplay of coronary artery calcification and traditional risk factors for the prediction of all-cause mortality in asymptomatic individuals *Circ Cardiovasc Imaging*, 5 (2012), pp. 467-473
- 16- G.S. Mintz, J.J. Popma, A.D. Pichard, et al. Patterns of calcification in coronary artery disease. A statistical analysis of intravascular ultrasound and coronary angiography in 1155 lesions *Circulation*, 91 (1995), pp. 1959-1965
- 17- G.J. Friedrich, N.Y. Moes, V.A. Muhlberger, et al. Detection of intraluminal calcium by intracoronary ultrasound depends on the histologic pattern *Am Heart J*, 128 (1994), pp. 435-441
- 18- G.S. Mintz, S.E. Nissen, W.D. Anderson, et al. American College of Cardiology Clinical Expert Consensus Document on Standards for Acquisition, Measurement and Reporting of Intravascular Ultrasound Studies (IVUS) *J Am Coll Cardiol*, 37 (2001), pp. 1478-1492
- 19- G.S. Mintz, P. Douek, A.D. Pichard, et al. Target lesion calcification in coronary artery disease: an intravascular ultrasound study *J Am Coll Cardiol*, 20 (1992), pp. 1149-1155
- 20- T.G. Kwon, Y.H. Seo, C.S. Lee, et al. Discrepancy of calcium detection between gray scale intravascular ultrasound and spectral analysis of radiofrequency data *Int J Cardiol*, 167 (2013), pp. 2611-2616
- 21- H. Yabushita, B.E. Bouma, S.L. Houser, et al. Characterization of human atherosclerosis by optical coherence tomography *Circulation*, 106 (2002), pp. 1640-1645
- 22- Topol EJ, Nissen SE. Our preoccupation with coronary luminology: The dissociation between clinical and angiographic findings in ischemic heart disease, *Circulation* 1995;92:2333-2342.
- 23- Ambrose JA, Winters SL, Arora RR, et al., Angiographic evolution of coronary artery morphology in unstable angina, *J Am Coll Cardiol* 1986;7:472-478.
- 24- Margolis JR, Chen JT, Kong Y, et al., The diagnostic and prognostic significance of coronary artery calcification, *Radiology* 1980;137:609-616.
- 25- Leber AW, Becker CR, Knez A, et al., Accuracy of 64-slice computed tomography to classify and quantify plaque volumes in the proximal coronary system: A comparative study using intravascular ultrasound, *J Am Coll Cardiol* 2006; 47:672-677
- 26- Nieman K, Oudkerk M, Rensing BJ, et al., Coronary angiography with multi-slice computed tomography, *Lancet* 2001; 357: 599-603.
- 27- Leber AW, Knez A, von ZF, et al., Quantification of obstructive and nonobstructive coronary lesions by 64-slice computed tomography: a comparative study with quantitative coronary angiography and intravascular ultrasound, *J Am Coll Cardiol* 2005; 46: 147-154.
- 28- Nakanishi T, Kayashima Y, Inoue R, et al., Pitfalls in 16-detector row CT of the coronary arteries, *Radiographics* 2005; 25: 425-438
- 29- Kuettner A, Kopp AF, Schroeder S, et al., Diagnostic accuracy of multidetector computed tomography coronary angiography in patients with angiographically proven coronary artery disease, *J Am Coll Cardiol* 2004; 43: 831-839.
- 30- Caussin C, Ohanessian A, Lancelin B, et al., Coronary plaque burden detected by multislice computed tomography after acute myocardial infarction with near-normal coronary arteries by angiography, *Am J Cardiol* 2003; 92: 849-852.
- 31- Toussaint JF, LaMuraglia GM, Southern JF, et al., Magnetic resonance images lipid, fibrous, calcified, hemorrhagic, and thrombotic components of human atherosclerosis in vivo, *Circulation* 1996;94:932-938.
- 32- Nissen SE, Gurley JC, Grines CL, et al., Intravascular ultrasound assessment of lumen size and wall morphology in normal subjects and patients with coronary artery disease, *Circulation* 1991;84:1087-1099.
- 33- Hodgson JMc, Reddy KG, Suneja R, et al., Intracoronary ultrasound imaging: Correlation of plaque morphology with angiography, clinical syndrome and procedural results in patients undergoing coronary angioplasty, *J Am Coll Cardiol* 1993;21:35-44.
- 34- Moore MP, Spencer T, Salter DM, et al., Characterization of coronary atherosclerotic morphology by spectral analysis of radiofrequency signal: In vitro intravascular ultrasound study with histological and radiological validation, *Heart* 1998;79:459-467.
- 35- Scanlon PJ, Faxon DP, Audet AM, et al. ACC/AHA guidelines for coronary angiography. A report of the American College of Cardiology/American Heart Association Task Force on practice guidelines *J Am Coll Cardiol* 1999 May;33(6):1756-824
- 36- Schroeder S, Kopp AF, Baumbach A, et al., Non invasive detection and evaluation of atherosclerotic coronary plaques with multislice computed tomography, *J Am Coll Cardiol* 2001; 37:1430-1435
- 37- Boaz D, Rosen, Veronica Fernandes, Robyn L, et al. Relationship between Baseline Coronary Calcium Score and Demonstration of Coronary Artery Stenoses During Follow up in the Multi-Ethnic Study of Atherosclerosis (MESA)
- 38- Budoff MJ, Georgiou D, Brody A, Agatston AS, et al. Ultrafast computed tomography as a diagnostic modality in the detection of coronary artery disease: a multicenter study