



## INTESTINAL PARASITIC INFECTIONS AMONG PATIENTS ATTENDING A TERTIARY CARE HOSPITAL IN BHAGALPUR

### Microbiolog

**Dr. Kaushal Kumar Gupta**

Tutor, Department Of Microbiology, Jawahar Lal Nehru Medical College And Hospital, Bhagalpur, Bihar, India

**Dr. Farhin Nishat\***

Tutor, Department Of Microbiology, Jawahar Lal Nehru Medical College And Hospital, Bhagalpur, Bihar, India \*Corresponding Author

**Dr. Amit Kumar**

Professor And Head, Department Of Microbiology, Jawahar Lal Nehru Medical College And Hospital, Bhagalpur , Bihar, India

### ABSTRACT

**Introduction:** Intestinal parasitic infections are among the most common infections in the world and are responsible for considerable morbidity and mortality. "There is increased prevalence of these infections in rural areas of developing countries." **Aim:** The present study was done to determine the intestinal parasitic infection among patients attending to tertiary care hospital. **Materials And Methods:** Faecal samples from 300 patients were collected and screened using conventional saline and iodine wet mount and examined by direct microscopy. Samples were further concentrated by formal-ether sedimentation technique. **Results:** In the present study the occurrence of intestinal parasitic infection was found to be 14.60%. The common parasite detected was *A. lumbricoides* (47.94%) followed by hookworm (30.13%). However, *E. histolytica* and *G. lamblia* were only 1.36%, 5.74% respectively. Multiple infection was detected only in 0.80% of infected cases. **Conclusion:** Soil transmitted helminthic infections are more common than protozoal infections. Hence there is a need for health programmes to be held regularly that will involve health education regarding personal hygiene, hand washing, importance of sanitary procedures and periodic deworming.

### KEYWORDS

Intestinal parasites, Occurrence, Ascaris.

#### INTRODUCTION:

Human intestinal parasitic infections have a worldwide distribution, with the greatest incidence and intensity occurring in developing countries<sup>1</sup>. The high prevalence in rural area of developing countries is probably due to poverty, poor sanitation, inadequate personal hygiene and lack of clean water supply. Helminths and protozoa are the two groups of pathogens causing intestinal illness with both direct and indirect method of transmission among people<sup>2</sup>. Helminthic infections are the main cause of loss of appetite, iron deficiency anaemia, gastrointestinal symptoms, perianal pruritus, gastrointestinal or biliary obstruction<sup>3</sup>. They also cause malnutrition in children<sup>4</sup> which leads to impairments in physical, intellectual and cognitive functions<sup>4,5,6</sup>.

In India overall prevalence rate of intestinal parasitic infection ranges from 12.5% to 66% with varying prevalence rate for individual parasite. About 50% of the urban population and 68% of the rural population in India is affected<sup>7</sup>. Intestinal parasitic infections are found in every age group and in both sex<sup>8</sup>. The purpose of this study was to determine the intestinal parasitic infection in the patients who are residents of Bhagalpur attending to the hospital.

#### MATERIALS AND METHODS:

The present study was carried out in the Department of Microbiology, in Jawahar Lal Nehru Medical College and Hospital. Three hundred patients of both sexes aged between 1 to 50 years with or without symptoms suggestive of parasitic infections were included in the study. The study was conducted between MARCH 2024 and September 2024. Informed oral consent was obtained from the patients. Questionnaire was administered on each subject so as to collect sociodemographic data including age, sex, type of toilet used, family size, source of drinking water and method of water treatment. Each patient was provided with a wide mouthed clean, dry, screw-capped container for collection of stool sample. The stool samples were examined within one hour of collection. Macroscopic examination was done to look for consistency, presence or absence of blood and mucus, adult intestinal worms and proglottids of *Taenia* species. The microscopic examination was done by normal saline and iodine mount directly from the stool. The samples found negative by microscopy were further concentrated by formal- ether sedimentation technique<sup>9</sup>. The mounts were screened under the microscope for the presence of helminthic ova, larva, trophozoites and cysts of protozoa. The modified Ziehl-Neelsen stain was used for the identification of coccidian parasites<sup>10</sup>.

#### RESULTS:

The overall occurrence of intestinal parasitic infections was 14.30% (73/500). Out of which 43 cases were detected by direct microscopy,

15 more cases were found positive after formal-ether concentration technique. Single infection was detected in 13.80% and dual infection was detected only in 0.80% of infected cases. *Ascaris lumbricoides* (48.8%) was the most common intestinal parasite, followed by hookworm (30.2%), *Trichuris trichiura* (9.3%), *Giardia lamblia* (6.9%), *Enterobius vermicularis* (4.6%), while *Entamoeba histolytica* (2.3%) [Table 1]. *A. lumbricoides* was found in single infection as well as in association with other helminthes in mixed infection [Table 2]. *Strongyloides stercoralis* infections were not found. No coccidian parasites were found.

Infection rate was highest (46.51%) in the age group 11-15 years, followed by 6-10 years (30.2%). The distribution of intestinal parasitic infections in various age groups is shown in Table 3. Among the 43 positive cases, 33 were outpatients and remaining 10 were inpatients. Out of 300 patients, 178 were males and 122 were females. Infected case distribution among males and females was 16.10% and 11.4% respectively [Table 4]. The sociodemographic data of the study group is given in Table 5. Among the positive cases, 37 (86.04%) used well water of which 34 (79.06%) boiled water before drinking. 61 positive cases did not use footwear regularly, 27 of them also gave history of defecation in open fields.

**Table 1. Distribution Of Intestinal Parasites (Single Infection).**

Parasites	Total number of cases (percentage)
<i>Ascaris lumbricoides</i>	20 (48.8%)
Hookworm	13 (30.2%)
<i>Trichuris trichiura</i>	4(9.3%)
<i>Giardia lamblia</i>	3 (6.9%)
<i>Enterobius vermicularis</i>	2 (4.6%)
<i>Entamoeba histolytica</i>	1 (2.3%)
Total	43(100%)

**Table 2. Occurrence Of Dual Infection.**

Parasites	Total number of cases (percentage)
<i>A. lumbricoides</i> + Hookworm	2 (2.73%)
<i>A. lumbricoides</i> + <i>T. trichiura</i>	1 (1.36%)
Total	3(4.09%)

**Table 3. Distribution Of Parasites In Different Age Groups.**

Parasites	Age in years									Total
	<5	6-10	11-15	16-20	21-25	26-30	31-35	36-40	>40	
<i>A. lumbricoides</i>	2	3	9	2	0	1	1	1	1	20

Hookworm	2	2	5	0	1	0	1	1	1	13
T. trichiura	1	2	1	0	0	0	0	0	0	4
E. vermicularis	1	1	0	0	0	0	0	0	0	2
G. lamblia	1	1	1	0	0	0	0	0	0	3
E. histolytica	0	1	0	0	0	0	0	0	0	1
S. stercoralis	0	0	0	0	0	0	0	0	0	0
Total	7	10	16	2	1	1	2	2	2	43

Table 4. Sex Wise Distribution Of Parasitic Infection.

Parasites	Males	Females
A. lumbricoides	14(51.02%)	6(41.66 %)
Hookworm	9 (26.53%)	4(37.5 %)
T. trichiura	2 (10.20%)	2 (8.33 %)
G. lamblia	2 (6.12%)	1 (4.16 %)
E. vermicularis	1 (4.08%)	1 (4.16 %)
E. histolytica	1 (2.04%)	0
S. stercoralis	0	0
Total	29 (16.2%)	14 (11.4%)

Table 5. Sociodemographic Data Of The Study Group

Parameter	Positive cases- 73 (%)	Negative cases- 427 (%)
<b>Source of drinking water</b>		
Well water	37(86.04%)	208(81.03%)
River water	5(11.63%)	-
Bore water	1 (2.33%)	49 (18.97%)
<b>Method of water treatment</b>		
Boiling	34(79.1%)	215 (83.66%)
Chlorination	4(9.3%)	26 (10.12%)
No treatment	5(11.6%)	16 (6.22%)
Usage of toilet	16 (37.20%)	195 (75.88%)
Open defecation	27 (62.80%)	62 (24.12%)
Usage of footwear	7 (16.27%)	200(77.82%)
No footwear used	36 (83.73%)	57 (22.18%)
Deworming done	6 (13.95%)	39 (15.1%)
Deworming not done	37 (86.05%)	218 (84.9%)

## DISCUSSION:

Intestinal parasitic infections are endemic worldwide and remain a major public health concern in many tropical and subtropical countries. The prevalence rates of intestinal parasitic infections and type of parasite exhibit wide variation from country to country, between geographical areas, communities and even seasonal variations also occurs<sup>11</sup>. Study on the occurrence of infection by various intestinal parasites is important to formulate appropriate intervention measures.

The present study showed 14.3% overall occurrence of infection, which is nearly comparable to the study of Baragundi et al (12%)<sup>8</sup>. However, our results are considerably lower than that of earlier studies<sup>1,4,12</sup>. In our study, A. lumbricoides (48.8%) was found to be the dominant intestinal pathogen, followed by hookworm (30.2%) and T. trichiura (9.3%). The possible reason for this, is the infective stage of A. lumbricoides, the embryonated egg have enormous capacity for withstanding the environmental extremes. Furthermore, Ascaris eggs are coated with a mucopolysaccharide that renders them adhesive to a wide variety of environmental surfaces<sup>13</sup>. They readily adhere to raw fruits and vegetables when fertilised with contaminated night soil or washed with contaminated water. They have also been found adherent to eating and cooking utensils and under fingernails. The results showed occurrence rate of helminthic infection are higher than those reported in other studies<sup>14</sup>.

In contrast to helminthic infection, the occurrence of protozoan infection in this study was significantly low (9.2%), but higher in other studies<sup>12,15</sup>. This may be due to people having access to clean drinking water, well water being the main source of water. Water is prime source of spread of infection along with contamination of food in protozoal infection<sup>16</sup>. Taenia and H. nana infections were not seen in our study, probably due to restricted consumption of pork and beef and H. nana infection significantly occurs by the presence of rodents and beetle in house. Contrary to our study some studies have shown high prevalence of H. nana<sup>7</sup>. All other parasites showed low occurrence when compared to other studies. This could be due to variation in geographical distribution of different parasite<sup>11,17</sup>.

We found that in our study dual infection was 1%, this coincides with the findings of most of the previous studies<sup>7</sup>. This is a clear indication of large number of various species of parasites in the locality. It was noted that occurrence of intestinal parasitic infection was highest among the age group 11-15 years (46.51%), followed by 6-10 years (30.2%). This finding is in agreement with previous studies<sup>12</sup>. 26.67% of the children were malnourished. Even though, gender is not a significant risk factor for intestinal parasitic infection<sup>18</sup>, in this study males were more infected than females. This finding is endorsed by earlier studies<sup>19</sup>. A. lumbricoides was the most common helminthic parasite in both male (50.12%) and female (41.66%), followed by hookworm (26.53% in male; 37.5% in female). This may be due to the contamination of soil by human faeces, use of raw sewage for agricultural purposes; use of waste water irrigated vegetables and contaminated imported vegetables<sup>20</sup>. 83.56% of the study population do not use footwear, and still 63.01% defecate in open fields which increase the susceptibility to helminthic infections. Mostly children are affected as they come in contact with contaminated water, food, faeces and other source of infection through play and other unhygienic behaviour<sup>20</sup>. No immunocompromised patients found in our study which may be the reason we found no coccidian parasites.

## CONCLUSION:

Human parasitic infection is a global problem of enormous proportion, responsible for mild but chronic morbidity. This region of study has high rainfall, high humidity, dense shade, poor sanitation and human host, which helps to complete the life cycle of the helminths. Hence there is a need to improve sanitation facilities. Regular health education should be conducted regarding personal hygiene, hand washing using soap, wearing shoes, periodic deworming and food hygiene. These multiple interventions will result in lowering infection rate but would require further investigation to verify the effect of these interventions.

## REFERENCES:

- Naish S, McCarthy J, William GM. Prevalence, intensity and risk factors of soil transmitted helminth infections in South Indian fishing village. Acta Tropica 2004; 91:177-87.
- Cleaveland S, Laurensen MK, Taylor LH. Diseases of human and their domestic mammals: pathogen characteristic host range and the risk of emergence. Phil Trans R Soc Land B 2001; 356:991-9.
- Samuel L, Stanley Jr. Amebiasis and infection with free living amebae. In: Longo DL, Fauci AS, Kasper DL, editors. Harrison's principles of internal medicine. 18th ed. USA: McGraw Hill Medical 2012; 1683-8.
- Srihari N, Kumudini TS, Mariraj J, Krishna S. The prevalence of intestinal parasitic infections in a tertiary care hospital- a retrospective study. J Pharm Biomed Sci 2011; 12(08):e1-4.
- Jardim- Botelho A, Raff S, Rodrigues, et al. Hook worm, A. lumbricoides infection and polyparasitism associated with poor cognitive performance in brazilian school children. Trop Med Int Health 2008; 13(8):994-4.
- Bethony J, Brooker S, Albonico M, et al. Soil transmitted helminth infections: Ascariasis, trichuriasis, hookworm. Lancet 2006; 367(9521):152-3.
- Ragunathan L, Kalivaradhan SK, Ramadas S, Nagaraj M, Ramesh K. Helminthic infection in school children in Pudukcherry, South India. J Microbiol Immunol Infect 2010; 43(3):228-32.
- Baragundi MC, Sonth SB, Solahamwar S, Patil CS. The prevalence of parasitic infections in patients attending tertiary care hospital. Nat J Bas Med Sci 2011; 2(1):31-4.
- Chatterjee KD, Parasitology: 12th ed. India: Chatterjee medical publisher, Calcutta 1995:211.
- Ortolani EL. Standardisation of the modified Ziehl -neelsen's technique to stain oocysts of Cryptosporidium species. Braz J Vet Parasitol 2000; 9(1):29-31.
- Tappe KH, Mohammadzadeh H, Khashaveh S, Rezapour B, Barazesh A. Prevalence of intestinal parasitic infection among primary school attending students in Barandoozchay region of urmia, West Azerbaijan province, Iran in 2008. Afr J Microbiol Res 2011; 5(7):788-91.
- Mbuh JV, Ntonifor HN, Ojong JT. The incidence, intensity and host morbidity of human parasitic protozoan infections in gastrointestinal disorder outpatients in Bura subdivision Cameroon. J infect Dev Ctries 2010; 4(1):38-43.
- Cromptein DWT. Biology of Ascaris lumbricoides. In: Cromptein DWT, Neshem MC, Pawlowski ZN editors. Ascaris and its prevention and control. London: Taylor & Francis 1989; 9-44.
- Choubasia SL, Choubasia L. Intestinal helminthic infections in tribal population of southern Rajasthan, India. J Parasit Dis 2006; 30(2):163-7.
- Bisht D, Verma AK, Bharadwaj HD. Intestinal parasitic infestation among children in a semi-urban Indian population. Trop Parasitol 2011; 104-07.
- Ravdin JI, Stauffer WM. Entamoeba histolytica (Amoebiasis). In: Mandell GL, Bennett JE, Dolin R, editors. Principles and Practice of infectious diseases. 6th ed. Pennsylvania: Elsevier Churchill Livingstone 2005; 3097-107.
- WHO Guidelines for evaluation of soil transmitted helminthiasis and schistosomiasis at community level. Schistosomiasis and intestinal parasite unit, WHO, Geneva WHO/CTD/SIP/98.1, 1998.
- Wani SA, Ahamad F, Zargar SA, Fomda BA, Ahmad Z, Ahmad P. Helminthic infestation in children of Kupwara district: A prospective study. Indian J Med Microbiol 2007; 25(4):398-400.
- Wani SA, Ahamad F, Zargar SA, Amin A, Ahmad Z, Ahmad P. Intestinal helminthiasis in children of Gurez valley of Jammu & Kashmir state, India. J Glob Infect Dis 2010; 2(2):91-4.
- Brooker S, Bundy DAP. Soil transmitted helminths ( Geohelminths). In: Cook GC, Zumla AI, editors. Manson's Tropical diseases. 22nd ed. China: Saunders Elsevier 2009; 151544.