



AUDIT OF DOCUMENTATION PRACTICES IN CERVICAL PAP TEST REQUISITION FORMS RECEIVED IN A TERTIARY CARE CYTOPATHOLOGY LABORATORY

Pathology

Dr. Bijoya Debnath*	Assistant Professor, Department of Pathology, Shija Academy of Health Sciences, Imphal, Manipur *Corresponding Author
Dr. Phuritshabam Pinky	Associate Professor, Department of Pathology, Shija Academy of Health Sciences, Imphal, Manipur
Dr. Wangol Kiyam	Associate Professor, Department of Pathology, Shija Academy of Health Sciences, Imphal, Manipur

ABSTRACT

Background: Pap smear cytology remains a cornerstone of cervical cancer screening, and its diagnostic accuracy depends not only on specimen adequacy and slide interpretation but also on the completeness of clinical information provided in laboratory requisition forms. Critical details such as patient demographics, menstrual history, prior cytology results, and relevant clinical findings help cytopathologists correctly interpret cellular changes. **Objective:** To assess the importance of completely and accurately filled Pap smear requisition forms for proper diagnostic evaluation. **Materials and method:** A retrospective cross-sectional study where all the original laboratory requisition forms received for Pap tests at the cytopathology unit of Shija Academy of Health Sciences, Imphal, Manipur, over a period of 10 months were retrieved, and completeness of clinical details and various other parameters were assessed along with the clarity of the documentation. **Results:** All Pap test LRFs consistently included patient details—name, age, gender, and unique identification number. However, clinical details and clinician-related information were largely absent in a significant proportion. **Conclusion:** Completely filled in Pap test LRFs are crucial for accurate cytologic diagnosis and effective patient management.

KEYWORDS

Cytopathology Laboratory, Pap Test, Laboratory Requisition Form

INTRODUCTION:

The necessity for close cooperation between clinicians and pathologists was clearly articulated as early as 1908 by Stow, who emphasized in his refined prose the importance of “the correlation of clinician, pathologist, and layman” [1]. Despite its importance, limited data exist regarding the completeness of Papanicolaou test requisition forms in resource-limited tertiary care settings. Concerns persist regarding the sufficiency of clinical information on test request forms, as some physicians fail to provide these details either because they are unaware of their significance or because of limited time [2]. To strengthen clinico-pathological communication, pathologists have a duty to inform clinicians of their critical role that complete and relevant clinical details play in achieving the most accurate interpretation of laboratory specimens [3]. However, a comprehensive review of the available literature revealed only a few reports addressing the adequacy of clinical data provided on Pap test forms [4, 5].

The accreditation criteria for medical laboratories established by the National Accreditation Board for Testing and Clinical Laboratories (NABL) specify that gynecological cytology request forms should include details such as menstrual phase, hormonal status, and prior surgeries. However, NABL does not define specific sample rejection criteria for Pap tests [6]. The Bethesda System (2014) for Reporting Cervical Cytology also emphasizes that cervical smear findings should be interpreted in the context of the clinical information provided on the request form [7].

This study aims to evaluate the completeness of Pap test laboratory requisition forms and identify the information most frequently omitted and highlight the importance of thoroughly completing these requisition forms.

Materials and method:

This is a retrospective cross-sectional study carried out in the Cytopathology Laboratory, Department of Pathology, Shija Academy of Health Sciences. A total of 204 laboratory requisition forms submitted for Pap test examinations from January 2025 to October 2025 were retrieved from the laboratory records for analysis.

The LRFs were reviewed for the presence of specific parameters, which were categorized into patient details, clinician details, and clinical information. Patient details included name, age, and unique patient identification number, while clinician details comprised the referring department and unit, physician's name, signature, and contact number.

The clinical information included signs and symptoms, examination findings, provisional diagnosis, menstrual cycle history, last menstrual period (LMP), and prior investigations or surgery.

In addition to evaluating the completeness of the forms, the clarity and readability of the recorded information were assessed, including handwriting legibility and the use of abbreviations.

Data were entered into Microsoft Excel and analyzed using descriptive statistics to determine the frequency and percentage of Pap test LRFs with complete and missing information for various parameters, while ensuring patient confidentiality throughout the study.

RESULTS:

A retrospective study of 204 Pap test LRFs was analyzed for a period of 10 months. Complete documentation of patient identifiers—name, age, gender, and unique identification number—was observed in all Pap test laboratory requisition forms. In contrast, while the referring department was invariably noted, the departmental unit was not specified in any of the forms. While the referring consultant's name was missing in only 1.47% of cases, clinician signatures were absent in 42.15% of the forms, and contact numbers were not documented in any forms (Table 2).

The LMP was documented in only 64 out of 204 forms (31.4%) [Table 1]. Many of the forms did not include clinical symptoms (50.9%), and only 88 forms (43.2%) mentioned the clinical examination findings (per vaginum and per speculum) [Table 2].

In 53.9% of the forms, no clinical diagnosis was provided, whereas the remaining forms (46.1%) documented diagnoses including amenorrhea (5%), pyometra (1%), postmenopausal bleeding (10.7%), candidiasis (5.3%), UV prolapse with secondary cystocele (1%), cervix hypertrophy with erosion (3.4%), chronic cervicitis (10%), pelvic inflammatory disease (6.3%), dyspareunia (2.4%), and post hysterectomy bleeding from the vault (1%).

Documentation of other relevant investigations was present in only 16 forms (7.9%). The most frequently missing parameters were LMP (68.6%), clinical diagnosis (53.9%), and relevant previous investigations (92.1%) [Table 2].

Legible and easily interpretable handwriting was observed in only 196 forms (96%) of the forms, while 16 forms (7.8%) contained unacceptable abbreviations or non-standard short forms [Table 3].

Figures and tables:**Table 1. Parameters on Pap test LRFs (n=204)**

Information	Parameter assessed	No of forms with incomplete data	No of forms with complete data
Patient	Name	0	204
	Age	0	204
	Gender	0	204
	Patient identification no	0	204
Clinician	Department	0	204
	Department unit	204	0
	Clinician name	3	201
	Clinician signature	86	118
Clinical	Clinician contacts no	204	0
	LMP	140	64
	Clinical symptoms	104	100
	Clinical examination	116	88
	Clinical diagnosis	110	94
Overall percentage	Relevant investigation	188	16
		40.4%	59.6%

Table 2. Percentage distribution of the data variables in Pap test LRF's

Name of the variable	Percentage of variables absent	Percentage of variables present
Name	0%	100%
Age	0%	100%
Gender	0%	100%
Patient identification no	0%	100%
Department	0%	100%
Department unit	100%	0%
Clinician name	1.4%	98.6%
Clinician signature	42.1%	57.9%
Clinician contacts no	100%	0%
LMP	68.6%	31.4%
Clinical symptoms	50.9%	49.1%
Clinical examination findings	56.8%	43.2%
Clinical diagnosis	53.9%	46.1%
Relevant investigation	92.1%	7.9%

Table 3. Clarity of data on Pap test LRFs (n=204)

Parameters	Number of LRFs	Percentage of LRFs
Non understandable handwriting	8/204	4%
Use of abbreviations	16/204	7.8%

DISCUSSION:

An enduring tenet of pathology is that accurate interpretation of cytological specimens relies not only on microscopic findings but also on the availability of relevant clinical information [8]. Since laboratory investigations influence approximately 70% of clinical diagnoses, they serve as a key determinant of both treatment outcomes and healthcare costs [9]. Inadequacies in laboratory requisition forms represent a major source of pre-analytic errors, predominantly arising from incomplete clinical and patient information [10].

In our present study, 204 laboratory requisition forms were evaluated across multiple parameters [Table 1]. Overall, 59.6% of LRFs were filled out completely, and 40.4% were incomplete. Only the patient's name, age, gender, patient identifier, and department in all the LRFs evaluated have 100% completion, indicating good compliance with basic documentation requirements.

The clinician's name and department were documented in 98.6% and 100% of cases, respectively, in our study, which is comparable to the findings of Sanskrithi G et al. [11], which documented 96.9% and 99.4%, respectively. The most frequently missing clinician-related detail was the departmental unit and clinician contact no., 100% each, followed by clinician signature, which was missing in 42.1% in our study [Table 2]. Lack of contact details affects the timely communication of critical results to the clinicians in addition to the precious time of the laboratory.

In the present study, the last menstrual period was not documented in

68.6% of LRFs, indicating a substantial deficiency in clinical information. Although this proportion is lower than that reported in a government tertiary hospital-based study in Tirupati, India (95.5%), the omission rate remains unacceptably high [11]. Evidence indicates that the timing of the last menstrual period can impact the accuracy of Pap smear findings, especially among women with persistent cervical intraepithelial neoplasia [12]. Nevertheless, failure to record LMP can significantly affect the interpretation of cytological findings, particularly gynecological specimens, thereby impacting diagnostic accuracy.

In our study, clinical diagnosis was mentioned in only 46.1% of the forms, which is closely aligning with the 41.6% reported in another study by Shivali Sehgal [13]. Clinical symptoms and clinical examination findings were documented in 49.1% and 43.2% of LRFs, respectively, in our study [Table 2], reflecting suboptimal recording of essential clinical details. This finding may be attributed to the fact that the majority of Pap smears were submitted for routine screening; however, inadequate documentation may also be due to factors such as time constraints in busy clinical settings, lack of awareness regarding the importance of detailed clinical information, or lack of structured requisition formats. It remains essential to document any clinical symptoms, as well as findings from per vaginum and per speculum examinations on the requisition form, even in the absence of abnormal findings.

It is also important to document additional clinical details in LRFs, including hormone therapy, birth control pill usage, pregnancy status, previous Pap findings, hysterectomy, IUD, abnormal bleeding, history of radiation or any malignancy, and any relevant investigation details [14]. The use of birth control pills mimics the effects of estrogen and progesterone, leading to an increase in the thickness of the cervical mucus plug [15]. Such information assists cytopathologists in forming accurate interpretations, thereby reducing the likelihood of false-negative or false-positive results. In our study none of the LRFs documented the use of hormonal therapy.

A study demonstrated that an educational intervention to healthcare providers with particular emphasis on providing relevant clinical details on the LRF, resulted in a significant improvement in quality indicators in the biochemistry laboratory [16]. Laboratories can adopt strategies suited to their resources to address the lack of information, with computerized entry and integration with the laboratory information system being an effective approach. Dogether et al. [17] recommended the use of electronic-based LRFs, as they found electronic forms to be more accurate, clearer, more complete, and easier to understand than paper-based forms.

CONCLUSION:

This study evaluated laboratory requisition forms with a focus on identifying specific errors and assessing the completeness of clinical information. The findings highlight that a substantial proportion of forms lacked essential information, particularly LMP (68.6%), clinical symptoms (50.9%), clinical examination findings (56.8%), provisional diagnosis (53.9%), and radiological findings (92.1%). These gaps hinder diagnostic accuracy and laboratory efficiency. Addressing these issues requires implementation of standardized forms, staff training on proper form completion, and the adoption of electronic requisition systems to ensure completeness, reduce errors, and improve overall patient care.

REFERENCES:

- Stow B. Correlation of clinician, pathologist and layman. JAMA. 1908;L1(3):191-4.
- Nakhleh RH, Gephardt G, Zarbo RJ. Necessity of clinical information in surgical pathology. Arch Pathol Lab Med. 1999;123:615-9.
- Nakhleh RE. Quality in surgical pathology communication and reporting. Arch Pathol Lab Med. 2011;135:1394-7.
- Naryshkin S, Schultz BL. Assessment of quality of data provided on pap test requisitions: Implications for quality of care and patient safety. Cytojournal. 2009;6:11.
- Costa DB, Carvalho AR, Chaves MA, Plewka J, Turkiewicz M. Patient safety by analyzing the information not provided in the requisition orders of cervical cytology test. J Bras Patol Med Lab. 2018;54:401-6.
- National Accreditation Board for Testing and Calibration Laboratories. Specific Criteria for Accreditation of Medical Laboratories (NABL 112) Issue No. 3. National Accreditation Board for Testing and Calibration Laboratories. 2012.
- Nayar R, Wilbur DC. The Bethesda System for Reporting Cervical Cytology: Definitions, Criteria, and Explanatory Notes. 3rd ed. New York: Springer; 2015.
- Rosai J, editor. Rosai and Ackerman's Surgical Pathology. 10th ed. Philadelphia: Mosby, Elsevier; 2011. pp. 1-20.
- Plebani M. Towards quality specifications in extra-analytical phases of laboratory activity. Clin Chem Lab Med. 2004;42:576-7.
- Witanikit V. Errors in laboratory requests in chulalongkorn hospital. Maharat Nakhon Ratchasima Hosp Med Bull. 2013;24:83-90.
- Sanskrithi G. Analysis Of Laboratory Requisition Forms Received In A Cytopathology

- Department of a Government Tertiary hospital- A cross sectional observational study. *Afr. J. Biomed. Res.* Vol. 28(1s) (January 2025); 1531 – 1541
12. Okodo M et al. Effects of the menstrual cycle on cytological false-negatives in women with persistent cervical intraepithelial neoplasia. *Cytopathology*. 2022 Sep; 33(5):591-9.
 13. Sehgal S, Jetley S, Jairajpuri ZS, Khan S. Critical Analysis of Laboratory Requisition Forms Received in a Cytopathology Laboratory of a Tertiary Care Centre: An Audit and Review of Literature. *J Cytol.* 2022 Jul-Sep; 39(3): 116–120.
 14. Herbert A, Bergeron C, Wiener H, Schenck U, Klinkhamer P, Bulten J and Arbyn M. European guidelines for quality assurance in cervical cancer screening. *Cytopathology*. 2007; 18(4):67-78
 15. National family planning guidelines for service providers. Nairobi: Division of Reproductive health, January 2010, pp. 69-160.
 16. Agarwal R, Chaturvedi S, Chhillar N, Goyal R, Pant I, Tripathi CB. Role on intervention on laboratory performance: Evaluation of quality indicators in a tertiary care hospital. *Indian J Clin Biochem.* 2012;27:61–8.
 17. Dogether MA, Muallem Y A, Househ M, Saddik B, Khalifa M. The impact of automating laboratory request forms on the quality of healthcare services. *J Infect Public Health.* 2016;9:749–56.