



## Spectrophotometric determination of Anionic Detergents in the river Sai at Jaunpur

\* Shafqat Alauddin \*\* Shailendra Yadav  
\*\*\* Tamheed Firdaus

\* Environmental Research Lab, Shibli National College, Azamgarh-276001, U.P., India

\*\* Environmental Research Lab, Shibli National College, Azamgarh-276001, U.P., India

\*\*\* Environmental Research Lab, Shibli National College, Azamgarh-276001, U.P., India

### ABSTRACT

Detergents includes some surfactants like LAS (linear alkyl sulphonate), TPBS (Tetra Propylene derived alkyl Benzene Sulphonate), ABS (branched chain alkyl benzene sulphonate) etc. LAS is most widely used anionic detergents and it have biodegradable capacity, while ABS like surfactants are resistant to biodegradability. The present study specially aims to study spectrophotometrically the contamination level of water of river Sai by the presence of anionic detergents flowing from the sub urban areas of Jaunpur district. In the river Sai direct wastes like municipal garbage, sewage as well as industrial wastes is dumped without treatment. River Sai is continuously been contaminated by the sewage and other types of wastes. This study revealed that there is gradual increase in the annual total concentration level of anionic detergents as we proceed from January to December along the years of study 2009, 2010 and 2011.

**Keywords :** anionic detergent, river Sai, spectrophotometer.

### INTRODUCTION-

In continuation of our study<sup>1,2</sup> on water quality of different rivers, we have now chosen river Sai of Jaunpur. Jaunpur is situated southeastern part of Uttar Pradesh and lies 25.730N latitude and 82.680E longitude embracing an area of nearly 4,038 km<sup>2</sup> (1,559 sq mile) with the total population 4, 476,072 as per census of 2011 having density of 1,108 km<sup>2</sup> (2,870 sq mile). A preliminary survey of the river Sai at Jaunpur showed that there were several points of interest. It flows from the suburban areas of the Jaunpur district and merges with River Gomti at the trimuhani (Oodpur) of the Jaunpur. It provides important source of water for the purpose of irrigation and drinking to the people of Jaunpur.

Detergents are used as a cleaning agent and components of detergents are surfactant (10-30%), builder (15%) and other ingredient like sodium silicate, amines, sodium sulphate. Surfactants are largely water soluble and forms foam in the waste water. Surfactants are classified as anionic, cationic, non-ionic and amphoteric as per their nature in water. Surfactants are used in the production of synthetic detergents, which are used every day life and industries viz. textile, cosmetics, paper, pharmaceuticals. Today's sewage contains appreciable amount of detergent along with other pollutants. River which flows near the suburban areas are polluted with such sewage which contains appreciable amount of detergent.

People of Jaunpur district use water of river Sai for irrigation, drinking and other day today needs. Intake of water having large quantity of detergent causes serious problems to human beings as well as to animals. It causes increase in the level of cholesterol, dilated mucous membranes, vomiting, lack of appetite, diarrhea and may cause corrosive injuries in the mouth and GI tract and may cause serious damage to aquatic life even death of fish fauna. Anionic detergents are found in shampoos, liquid detergents and utensil cleaners. They find their way to river water without treatment with other household sewage and industrial wastes. As per BIS the maximum contaminant level for anionic detergent is 1.00 mgL<sup>-1</sup> for river water.

**SAMPLING-** This study was done in the years 2009, 2010 and 2011. Three sampling stations were earmarked which are given below in table -1

**TABLE-1**

S. No.	Sampling Stations	Classification
1.	Teji Bazaar Bridge	Entry Point
2.	Ram Dayal Ganj Bridge	Mid Point
3.	Oodpur	Exit Point (outside the city population)

Water samples were randomly collected at the above mentioned sites from January to December during all the three study years' in the poly ethylene bottles. Samples were taken at a distance of 2 meters inside the river from the bank and depth of about 0.20 meter. The samples are taken between 15th day to 20th day of the each month.

**MATERIALS-** Materials used are of analytical grade provided by the Fisher Scientific. Following methods are used in the present study-

1. Methylene Blue Solution
2. 6 M HCl
3. Chloroform
4. Distilled Water

**EXPERIMENTAL-** A series of solutions of commercially available sodium dodecyl benzene sulphonate as T1=0.5 mgL<sup>-1</sup>, T2=1.0 mgL<sup>-1</sup>, T3=1.5 mgL<sup>-1</sup>, T4=2.0 mgL<sup>-1</sup> T5=2.5 mgL<sup>-1</sup> and T6=3.0 mgL<sup>-1</sup> by the dilution of stock solution.

The chemical reaction takes place between ionic dyes stuff and the detergents is as follows-



Where **MB<sup>+</sup>** indicate the cation of methylene blue. The reaction product can be extracted by chloroform where as the original dyestuff is insoluble in this medium and the intensity

of the cation in the chloroform layer is proportional to the concentration of the detergent. Absorbance is measured at 650 nm using 1 cm cell. A calibration curve (Fig-1) is drawn.

**RESULTS AND DISCUSSION-** The results obtained from the analysis of different parameters and for different years of study were recorded in table (2) and (3).

**TABLE-2**  
**DETAILS OF CONCENTRATION OF ANIONIC DETERGENTS**

S.No.	Month	Years' of study								
		2009			2010			2011		
		Concentration of anionic detergents in mgL-1								
		1	2	3	1	2	3	1	2	3
Entry Point	Mid Point	Exit Point	Entry Point	Mid Point	Exit Point	Entry Point	Mid Point	Exit Point		
1.	January	0.63	0.66	0.72	0.64	0.67	0.75	0.79	0.82	0.91
2.	February	0.67	0.70	0.73	0.69	0.74	0.80	0.70	0.75	0.83
3.	March	0.75	0.81	0.92	0.78	0.86	0.97	0.79	0.89	0.99
4.	April	0.79	0.84	0.93	0.84	0.90	1.00	0.83	0.90	1.00
5.	May	0.90	0.98	1.07	0.93	1.06	1.11	1.00	1.06	1.19
6.	June	0.95	1.06	1.08	0.98	1.03	1.13	1.05	1.13	1.21
7.	July	1.04	1.11	1.17	1.05	1.10	1.19	1.10	1.21	1.27
8.	August	1.03	1.07	1.13	1.06	1.11	1.17	1.12	1.18	1.25
9.	September	0.91	0.96	1.04	1.17	1.22	1.33	1.03	1.06	1.20
10.	October	0.89	0.93	1.03	1.07	1.16	1.22	1.01	1.04	1.18
11.	November	0.82	0.86	0.91	0.86	0.89	0.96	0.88	0.86	0.99
12.	December	0.71	0.77	0.80	0.72	0.76	0.83	0.83	0.86	0.93

**TABLE-3**  
**DIFFERENCE OF CONCENTRATION BETWEEN ENTRY AND EXIT POINTS**

S.No.	Month	Years' of study		
		2009	2010	2011
		Concentration of anionic detergents in mgL-1		
1.	January	0.09	0.11	0.12
2.	February	0.06	0.11	0.13
3.	March	0.17	0.19	0.20
4.	April	0.14	0.16	0.18
5.	May	0.17	0.18	0.19
6.	June	0.13	0.15	0.16
7.	July	0.13	0.14	0.17
8.	August	0.10	0.11	0.13
9.	September	0.13	0.16	0.17
10.	October	0.15	0.16	0.17
11.	November	0.09	0.10	0.11
12.	December	0.09	0.11	0.13
Total Annual increase of anionic detergent		1.45	1.68	1.86

From the tables it was found that there is a slight increase in the concentration of anionic detergents during summer and rainy

season i.e. in the month of March, April, May, June, July and August and decrease during the winter season i.e. in the month of September, October, November and December in every year of study, and this trend continue in the month of January in the subsequent year. From February the gradual increase in the concentration of anionic detergent till the month of August.

The increase shown at the entry level every year is due to dumping of the sewage of suburban areas, because it is river of sub urban area of Jaunpur district.

Data in table (3) revealed that the cumulative concentration of anionic detergent is increasing in gradual manner, if move across the years of study.

**CONCLUSION-** It has been observed during this study that there is gradual increase in concentrations of anionic detergent in the river Sai during subsequent years. The total concentration of anionic detergent increasing rapidly and the level of anionic detergent crossed the MCL during the years.

The concentration of anionic detergent is increasing due to uncontrolled use of synthetic detergents containing LAS. The concentrations of anionic detergent is increasing with every passing year and it is high time to control the dumping of untreated sewage and industrial wastes containing anionic detergent to the river water to save the river Sai from pollution.

## REFERENCES

1. Alauddin Shafiqat, Kumar Amresh, Yadav Shailendra and Firdaus Tamheed, Spectrophotometric determination of Iron, Nitrate and Phosphate in the river Gomati of Jaunpur city, International Journal of Scientific Research, 2012, Vol. 1(6), 7-8 | 2. Harrison R.M., Understanding Our Environment and Pollution, Cambridge Royal Society, 1992. | 3. Phillip M., Environmental Chemistry, 2nd Edition, 1996, 46-49 | 4. Skoog and West, Fundamentals of Analytical Chemistry, 2nd Ed., Chapter 29. | 5. Vogel, A Textbook of Quantitative Inorganic Analysis, 3rd Ed., p. 294, 310 and 787 | 6. APHA, AWWA, AND WEF, Standard methods for the Examination of Water and Wastewater, 19th Ed., Washington, DC, American Public Health Association, 1995. | 7. Richards M., Effect of detergent use on water quality in Kathmandu, Nepal, Massachusetts Institute of Technology, 2003 | 8. Medson G., Peterson and Simorsen F., Environmental and health assessment of substances in household detergents and cosmetic detergents products, 2003, Danish Environmental Protection Agency. |