

## Research Paper

## Veterinary



## Study on Comparative Efficacy of Polyherbal Preparation and Synthetic Methionine, Choline, Lysine and Biotin on Performance of Rhode Island Red Layers

\* Kiran Kumari \*\* Tiwari SP \*\*\* Ravikanth K  
\*\*\*\* Thakur A

\*, \*\* Department of Animal Nutrition, College of Veterinary Sciences, Anjora, Durg, (Chhattisgarh) India

\*\*\*, \*\*\*\* R & D Team Ayurvet Ltd, India

### ABSTRACT

An experiment was conducted for 20 weeks to evaluate the comparative efficacy of dietary supplementation of herbal (AV/CAP/18) and synthetic methionine, choline, lysine and biotin on the egg production and egg quality traits. 90 Rhode Island Red commercial layers of 15 weeks of age were randomly divided in 3 groups of 30 birds each. All birds were offered standard basal diet so as to meet their all basic nutritional requirements. Group I (T<sub>0</sub>) served as control group while Group II (T<sub>1</sub>) supplemented with combination of synthetic choline (60%) 500g/ton+ methionine 1kg/ton+ lysine 1 kg/ton + biotin 150 mg/ton and III (T<sub>2</sub>) with AV/CAP/18 @ 1 kg/ton (supplied by M/S Ayurvet Ltd. Baddi. (HP). India). At the end of 35th week, significantly higher egg production, egg weight, egg mass, yolk weight and albumen weight were recorded in T<sub>1</sub> followed by T<sub>2</sub> and lowest in T<sub>0</sub>. Mean feed conversion ratio was also better in T<sub>1</sub> (1.89) and AV/CAP/18 fed T<sub>2</sub> (1.88) than control T<sub>0</sub> (2.63). Lowest mortality percentage (%) was recorded in T<sub>2</sub> (1.78%) than T<sub>1</sub> (1.81%) and T<sub>0</sub> (3.51%). On the basis of the results obtained it was concluded that herbal combined amino acid supplement AV/CAP/18 has efficacy comparable to its synthetic counterparts. Their successful incorporation in balanced layer ration may lead to improved egg production and quality traits.

**Keywords :** AV/CAP/18, amino acids, polyherbal, egg production, egg quality traits

### 1. INTRODUCTION

Protein and amino acids have an overwhelming importance in poultry feed (Dersjant and Peisker, 2011). The scientific feeding of poultry is not in fact based on the raw protein of the diet but on the amino acids content of the diet, their balance and intake. (NRC1994). Appropriate concentration of energy and amino acids in feed are needed in order to increase the productivity of laying hens (Chan *et al.*, 2007). Available layer hybrids have nutritional requirements that may be difficult to meet unless extra amino acids can be supplemented in the diet (Berg, 2001 and Odelros, 2000). Imbalance in amino acids composition in layer diet which is needed for egg protein synthesis will consequently result in lowered egg output alongwith poor plumage condition, feather pecking, cannibalism and increased overall mortality (Ambrosen and Peterson, 1997). Synthetic methionine is metabolized into highly toxic compounds such as methylpropionate, thereby, adversely altering the performance of poultry birds (Bender, 1975). Synthetic methionine is listed among the prohibited synthetic substances and its usage has been questioned in organic farming practices (Anonymous, 1999). Therefore, the present study was undertaken to determine the comparative efficacy of combination of synthetic methionine + choline + lysine and biotin with polyherbal feed supplement AV/CAP/18 (supplied by M/S Ayurvet Limited, Baddi, (HP) India) incorporation on egg production and egg quality traits in RIR layers. AV/CAP/18 is a polyherbal product comprises of herbs like *Trigonella foenum graecum*, *Allium sativum*, *Mucuna pruriens* & many others in a fixed concentration.

### 2. MATERIALS AND METHODS

A study was conducted at Department of Animal Nutrition, College of Veterinary Sciences, Anjora, Durg, (Chhattisgarh) India. Total 90 RIR layer birds of 15 weeks age were procured and maintained under identical conditions. Vaccination was done as per the recommended schedule. All the birds were

fed standard basal diet formulated as per the recommendations of NRC, 1994. Further birds were randomly divided into 3 groups of 30 birds with 3 replicates of 10 birds in each group.

Group I (T<sub>0</sub>): control without any supplementation

Group II (T<sub>1</sub>): supplemented with combination of synthetic choline (60%) 500g/ton+ methionine 1kg/ton+ lysine 1 kg/ton + biotin 150 mg/ton

Group III (T<sub>2</sub>): supplemented with AV/CAP/18 @ 1 kg/ton (supplied by M/S Ayurvet Ltd. Baddi. (HP) India)

Individual body weights were recorded on weekly basis in order to determine the body weight changes. The number of eggs laid, egg weights and feed consumption (g/hen) were recorded daily and accordingly, the egg production (%), egg mass (g/hen/day) and feed conversion (g feed/g egg weight) were calculated. To measure egg quality, 50 eggs in each group (total 150 eggs) were sampled during time interval of 20-35<sup>th</sup> weeks. First the whole egg was weighed (g), next it was broken and egg contents were separated carefully into yolk, albumen and shell and then internal traits were measured. Tripod micrometer was used to measure thick albumen height (El-Safty and Mahrose, 2009). Also albumen weight (g), yolk weight (g), yolk to albumen ratio alongwith the egg component % was determined individually. Haugh unit was measured using the formula: HU = 100 log (H - 1.7w<sup>0.37</sup> + 7.6), where HU=Haugh unit, H=Observed height of the albumen in mm W=Weight of egg (g). Yolk colour score was measured with Roche Yolk Colour Fan (Vuilleumier, 1969). The eggshell was dried and weighed subsequently to the nearest 0.01g (Ghazalah *et al.*, 2011). Egg-specific gravity (unit), Mortality (%) was also recorded.

**Statistical analysis:**

Mean, standard deviation (SD), analysis of variance (ANOVA), least significant differences (LSD) and co-efficient of correlation values (r) were computed using the SPSS (version 11.0 for Windows).

**3. RESULTS AND DISCUSSION**

At the end of 35<sup>th</sup> week average body weight attained was highest in T<sub>1</sub> (1.74) followed by T<sub>2</sub> (1.73) and T<sub>0</sub> (1.49). T<sub>1</sub> performed better (98 g/hen/day) than T<sub>2</sub> (99 g/hen/day) and control group birds (118 g/hen/day) in terms of feed intake (Table 2). The feed conversion (g feed/g egg weight) efficiency was also significantly better (P<0.05) in T<sub>1</sub> (1.89) and T<sub>2</sub> (1.88) as compared to T<sub>0</sub> (2.63). Similar finding were also noted by the Harms and Russell, (1996), Keshavarz and Jackson, (1992). There is less consumption of feed in treated groups as compared to control, as demonstrated earlier that mild deficiencies of protein, lysine and methionine in layers (Picard *et al.*, 1993) lead to small increase in feed intake. Similar results were obtained by Abdel (2006) in case of feed intake, FCR and body weight gain when *Trigonella foenum graecum* was added to broiler feed as growth promoter and Iyayi and Taiwo (2003) in case of *Mucuna pruriens*. The difference in feed intake were in agreement with findings of Wu *et al.*, (2005), who reported that with increasing dietary energy, hens can adjust feed intake so that same amount of dietary energy is used to produce 1 g of egg. The improvement in weight gain may be due to the action of allicin (an antibiotic substance found in garlic), which inhibits the growth of pathogenic bacteria and aflatoxin producing fungi. The mode of action of these feed additives is not completely clear. They have antimicrobial, antiviral, antioxidant and many other biological activities (Cross *et al.* 2002). They act as a digestibility enhancers, stimulating the secretion of endogenous digestive enzymes (William and Rosa, 2001).

Particular	Daily nutrient intake per hen						
Group	Energy: Lys (MJ/g)	Energy (MJ)	Protein (g)	TSAAs (g)	Lys (g)	Available P (g)	Ca (g)
T <sub>0</sub>	1.18	1.26	19.95 <sup>a</sup>	0.804	1.072	0.445	4.243
T <sub>1</sub>	1.17	1.28	19.55 <sup>b</sup>	0.817	1.088	0.446	4.251
T <sub>2</sub>	1.17	1.28	19.55 <sup>b</sup>	0.817	1.088	0.446	4.251

**Table 1. Effect of herbal methionine, lysine, choline and biotin on nutrient intake per hen daily in RIR birds (from 20 to 35 wk of age)**

<sup>a-b</sup> Means within a column and under each main effect with no common superscripts differ significantly (P < 0.05).

Egg production and components traits revealed that both the treatment groups T<sub>1</sub> and T<sub>2</sub> differed significantly (P<0.05) with control group (T<sub>0</sub>) which was supplied basal diet without methionine, choline, lysine and biotin. However both the treatment groups were having identical values in terms of egg production and component traits. Egg production % in group T<sub>1</sub> (90.22%) and T<sub>2</sub> (90.20%) was significantly higher (P<0.05) than T<sub>0</sub> (75.05%). Also the egg weight observed in T<sub>1</sub> (56.75g) and T<sub>2</sub> (56.68g) was significantly higher (P<0.05) than 48.84 g (T<sub>0</sub>). Group T<sub>1</sub> (52.25) and T<sub>2</sub> (52.18) also had significantly higher egg mass (g egg/hen per d) as compared to T<sub>0</sub> (44.89). (Table 2)

Yolk weight in T<sub>1</sub> (15.34 g) and T<sub>2</sub> (15.33 g) were significantly higher (P<0.05) than T<sub>0</sub> (15.04 g). Albumen weight was also significantly higher (P<0.05) in T<sub>2</sub> (38.19 g) and T<sub>1</sub> (38.13 g) than T<sub>0</sub> (36.51 g).

**Table 2. Effect of herbal methionine, lysine, choline and**

**biotin on performance in RIR birds (from 20 to 35 wk of age)**

Particular	Feed intake (g/hen per day)	Egg production (%)	Egg weight (g)	Egg mass (g egg/hen per day)	Feed conversion (g of feed/g of egg)	Egg-specific gravity (unit)	BW (kg)	Mortality (%)
Group								
T <sub>0</sub>	118	75.05	48.84	44.89	2.63	1.0870	1.49	3.51
T <sub>1</sub>	99 <sup>c</sup>	90.22 <sup>b</sup>	56.75 <sup>a</sup>	52.25 <sup>a</sup>	1.89 <sup>c</sup>	1.0870 <sup>c</sup>	1.74 <sup>a</sup>	1.81
T <sub>2</sub>	98 <sup>c</sup>	90.20 <sup>b</sup>	56.68 <sup>a</sup>	52.18 <sup>a</sup>	1.88 <sup>c</sup>	1.0869 <sup>c</sup>	1.73 <sup>a</sup>	1.78

<sup>a-c</sup> Means within a column and under each main effect with no common superscripts differ significantly (P < 0.05).

Shell weight of eggs in group T<sub>2</sub> which was fed with polyherbal preparation AV/CAP/18 was higher (5.40 g) than T<sub>1</sub> (5.36) and T<sub>0</sub> (4.71 g). Thus, better shell strength and ultimately egg quality was obtained with herbal product supplementation.

The Haugh unit (HU) being a measure of freshness of eggs, HU unit below 60 is considered as poor quality egg. Thus HU of eggs from hens fed with the herbal (69.99) and synthetic (70.07) nutrient (methionine, choline, lysine and biotin) was significantly lower than that of hens fed with the basal diets (74.87). This may be because of increased weight of eggs. Egg-specific gravity remained unaffected in almost all groups.

In the present investigation egg component weight and egg component percentage of the AV/CAP/18 supplemented group as well as the synthetic polyamino acid supplemented group was significantly higher (p<0.05) from control (Table 3).

**Table 3. Effect of herbal methionine, lysine, choline and biotin on egg components in RIR birds (from 20 to 35 wk)**

Particular	Egg component weight (g)			Egg component (%)			
	Yolk	Albumen	Shell	Yolk	Albumen	Shell	Yolk: albumen
Group							
T <sub>0</sub>	15.04	36.51	4.71	25.76	64.85	9.03	0.397
T <sub>1</sub>	15.34 <sup>a</sup>	38.13 <sup>b</sup>	5.36 <sup>c</sup>	26.07	64.86	9.07	0.402
T <sub>2</sub>	15.33 <sup>a</sup>	38.19 <sup>b</sup>	5.40 <sup>c</sup>	26.07	64.85	9.08	0.402
P - value							
Treatment group	0.05	0.05	0.05	NS	NS	NS	NS

<sup>a-c</sup> Means within a column and under each main effect with no common superscripts differ significantly (P < 0.05).

The increase in egg production and component traits in case of treated groups is supported by the findings that Low intake of amino acids lead to reduced metabolism (Pourreza and Smith, 1988), ultimately resulting in lowered protein synthesis for body growth and egg production performance (Harms and Russell, 1996; Keshavarz and Jackson, 1992). The increase in egg weight and egg mass is supported by the findings of Shafer *et al.*, (1998), Liu *et al.*, (2005) and Zewei *et al.*, (2011) who reported that egg weight increased with increasing level of amino acid intake by laying hens. Egg mass was found to be having positive correlation with that of amino acid content in the layer diet Keshavarz, (1992). The increased egg mass was mainly contributed from the increased egg weight. The increase in albumen may be due to limited availability of amino acids having adverse effects on albumen solids (Shafer *et al.*, 1998) and egg size (Keshavarz and Jackson, 1992; Hsu *et al.*, 1998). Similar results were obtained by Hassan and Ragab (2009) in case of supplementation of *Trigonella foenum graecum* on productive performance and egg quality

of layers and by Iyayi and Taiwo (2003) in case of *Mucuna pruriens*.

Lowest mortality percentage was noted in  $T_2$  (1.78%),  $T_1$  (1.81%) and  $T_0$  (3.51%) (Table 2). Low level of mortality percentage is seen in treatment groups as adequate dietary provision of all amino acids is necessary for sustaining normal immunocompetence and protecting the host from a variety of diseases in all species (Peng Li *et al.*, 2007). Imbalance in amino acids composition in layer diet will consequently result in lowered egg output alongwith poor plumage condition, feather pecking, cannibalism and increased overall mortality (Ambrosen and Peterson, 1997).

#### 4. CONCLUSION:

The data from the trial indicates that polyherbal preparation could successfully replace the synthetic methionine, choline, lysine and biotin amino acids for optimum production performance of laying Hens as performance of layer birds supplemented with AV/CAP/18 can well be compared with birds supplemented with synthetic amino acids.

#### ACKNOWLEDGEMENTS

The Author is thankful to Dean College of Veterinary and Animal Sciences, Durg, Raipur (Chhattisgarh) for providing all the necessary facilities for the successful conduct of the trial and Ayurved Ltd. Baddi for providing necessary samples.

#### REFERENCES

- Abdel-Azem, F. (2006). Effect of using fenugreek and fennel seeds as natural feed additives on | performance of broiler chicks. Egyptian Journal of Nutrition and Feeds. 9: 277-297. | Ambrosen, T., Petersen, V.E. (1997). The influence of proteins levels in diet on cannibalism and plumage quality in layers. Poultry Science. 76: 559-563. | Anonymous, (1999). Statistica v5.5. Statsoft, Inc. Tulsa, OK, USA. | Bender, D.A. (1975). Amino acid metabolism. John Wiley and Sons. Ltd. 1st Ed., New York, | USA, pp: 112-142. | Berg, C. (2001). Health and Welfare in Organic Poultry Production Acta vet. Scand. 95, 37-45. | Chen, C., Sander, J.E. and Dale, N.M. (2003). The effect of dietary lysine deficiency on the immune response to Newcastle disease vaccination in chickens. Avian Dis., 47: 1346- 1351. | Cross, D.E., Acamovic, T., Deans, S.G. and McDewitt, R.M. (2002). The effects of dietary | inclusion of herbs and their volatile oils on the performance of growing chickens. British Poultry | Science. 43:33- 35 | Dersjant-Li Y. and Peisker, M. (2011). A review on recent findings on amino acids requirements | in poultry studies. Iranian journal of applies animal science. 1 (2): 73-79. | El-Safty, S. and Mahrose, M. (2009). Evaluation of some phenotypic, physiological and egg quality traits of African black neck ostrich under arid desert conditions of Libya International journal of poultry science. 8 (6). | Ghazalah, A.A., Abd- Elsamee, M.O. and Eman, S.M. (2011). Use of distillers dried grains with soluble (DDGS) as replacement for soybean meal in laying hen diets. International Journal of poultry science. 10 (7): 505-513. | Harms, R.H. and Russell, G.B. (1996). A re-evaluation of the methionine requirement of the | Commercial layer. Journal of Applied Animal Research. 9:141-151. | Hassan, H.A. and Ragab, M.S. (2009). The productive performance and egg quality of laying hens under the effect of dietary fenugreek seeds (ungerminated and germinated) and different | dietary protein levels. Poult. Dept., Fac. of Agric., Fay. Univ., Fayoum. Egypt. | Hsu, J.C., Lin, C.Y. and Chiou, W.S.P. (1998). Effects of ambient temperature and methionine | supplementation of a low protein diet on the performance of laying hens. Animal Feed Science | Technology, 74:289-299. | Iyayi, E.A. and Taiwo, V.O. (2003). The effect of diets incorporating *Mucuna* (*Mucuna pruriens*) seed meal on the performance of laying hens and broilers. Journal of Tropical and Subtropical Agro ecosystems. 1 (2-3): 239 - 246. | Keshavarz, K. and Jackson, M.E. (1992). Performance of growing pullets and laying hens fed low-protein, amino acid-supplemented diets. Poultry Science. 71:905-918. | Liu, Z., Bateman, A., Bryant, M.M., Zinner, B. and Roland, D.A. (2005). Performance comparisons between DL-methionine and DL methionine hydroxy analogue in layers on an unequalmolar basis. Journal of Applied Poultry Research. 14: 569-575. | National research council, (1994). Nutritional requirements in methionine + cystine for white-egg laying hens during the first cycle of production. International Journal of Poultry Science. 4: 965-968. | Odelros, A. (2000). Oro bland ekologiska äggproducenter (Worries among organic egg producers). Fjäderfå, 5: 22-28. | Peng, Li, Yu-Long, Yin, Defa, Li, Sung, W.K. and Guoyao, Wu. (2007). Amino acids and immune function. British Journal of Nutrition. 98: 237-252. | Picard, M., Sauveur, B., Fenardji, F., Angulo, I. and Mongin, P. (1993). Suitable technico-economic adjustments for poultryfeeding in hot countries (in French). INRA Production Animal.6:87-103 | Pourreza, J. and Smith, W.K. (1988). Performance of laying hens fed on low sulphur amino acids | diets supplemented with choline and methionine. British Poultry Science. 29:605-611. | Shafer, D.J., Carey, J.B., Prochaska, J.F. and Sams, A.R. (1998). Dietary methionine intake effects on egg component yield, composition, functionality and texture profile analysis. Poultry Science. 77: 1056-1062. | Vuilleumier, J.P. (1969). The roche yolk colour fan in instrument for measuring of yolk colour. Poultry Science., 48:767-779. | William, P. and Rosa, R. (2001). The use of essential oils and their compounds in poultry nutrition. World Poultry. 17: 14-15. | Wu, G., Bryant, M.M., Voitle, R.A. and Roland, D.A. (2005). Effect of dietary energy on performance and egg composition of Bovans White and Dekalb hens during phase 1. Poultry Science. 84: 1610-15 | Zewell, H.S., Abdalah, A.A., Ahmed, M.H. and Ahmed, R.S. (2011). Effect of different levels of | protein and methionine on performance of laying hens and environmental pollution. Egypt Poultry Science. 31: 621-639. | ]