



An article on imaging of posterior fossa mass lesions in adults

Dr.C.Nellaiappan

Associate professor of radiodiagnosis department, Tirunelveli medical college hospital, tirunelveli, Tamilnadu

Dr.S.Nancydora

Associate professor of radiodiagnosis department, Tirunelveli medical college hospital, tirunelveli, Tamilnadu

ABSTRACT

Posterior fossa mass lesions in adults are multiple. Cerebellopontine angle lesions are common. cerebellar parenchymal lesions can also occur. In this article we discuss about differential diagnosis of posterior fossa mass lesions in adults. cerebellar gangliocytoma, meningioma are discussed in detail.

KEYWORDS

computerised tomography, magnetic resonance imaging, cerebellopontine angle, gangliocytoma, meningioma

INTRODUCTION

Radiologic assessment of posterior fossa tumors may be achieved with several imaging techniques. computed tomography (CT), MRI are the standard techniques. Nuclear scan, PET scan, angiography are sometimes used. The radiographic features of calcification, are usually well demonstrated on CT scan. MRI is particularly useful in distinguishing intra axial from extra axial lesions. The location and extent of masses can be defined using CT and MRI. In this article, we discuss two mass lesions in the posterior fossa in adults.

Case 1:

38 years female admitted in our hospital with headache and ataxia. CT scan was taken. It shows a hyper dense mass lesion in the posterior fossa on left side causing fourth ventricle compression and mild hydrocephalus (fig1). On contrast administration, the lesion shows good enhancement (fig2). MRI – Brain was taken for further evaluation. T1 hypo intense, T2 hyper intense extra axial lesion without internal auditory canal component is seen (fig3,4,5). On contrast, the lesion shows very good enhancement without cystic components (fig6). Radiological differential diagnosis includes meningioma, acoustic nerve/ facial nerve schwannoma in atypical location. The patient underwent surgical removal. HPE is suggestive of fibroblastic meningioma.

Case 2:

43 year old female admitted with headache and ataxia. CT scan was taken. It shows mixed dense mass lesion with predominant hypo dense areas in the midline of cerebellum causing compression of fourth ventricle and minimal hydrocephalus (fig7). MRI – Brain was taken for the future evaluation. It shows T1 mixed intense, T2 hyper intense lesion, not suppressed in FLAIR lesion in the midline cerebellum compressing fourth ventricle (fig8,9,10). MRI FLAIR also shows prominent folia like pattern within the lesion (fig10). On contrast no significant enhancement of mass seen (fig11,12). The radiological differential diagnosis include ependymoma, cerebellar astrocytoma, rarely dysplastic gangliocytoma. The patient underwent surgical removal of mass. HPE is suggestive of dysplastic gangliocytoma.

DISCUSSION

Posterior fossa tumour has a very different differential in an adult as opposed to a child.

cerebellar metastases (most common) especially from lung cancer and breast cancer and also

from melanoma, thyroid malignancies, and renal cell cancer. Haemangioblastoma is most common primary parenchymal cerebellar tumour. astrocytomas and medulloblastomas are rare in the posterior fossa of adults (<1% all tumours).

A. James Barkovich. Philadelphia, PA : Lippincott Williams & Wilkins (2005) found that brain tumor distribution varies with age as follows⁽¹⁾

0 to 3 years of age	supratentorial > infratentorial
4 to 10 years of age	infratentorial > supratentorial
10 to early adult hood	infratentorial = supratentorial
adults	supratentorial > infratentorial

James G. Smirniotopoulos, MD, Nancy Chang Yue, MD, Lisabetbj Rushing (1993) have shown the percentage of masses of cerebellopontine angle as follows

Masses of the CPA	Percentage Mass of Cases ⁽²⁾
Acoustic (vestibular) schwannoma	80-90
Meningioma (petrous/tentorial)	10-15
Epidermoid inclusion cyst	5-9
Arachnoid cyst	<1
Lipoma (lipomatous hamartoma)	<1
Aneurysm	<1
Ependymoma	<1
Choroid plexus papilloma	<1
Metastasis	<1

Classification of posterior fossa meningioma "Sekhar and Wright"⁽³⁾

TYPE	LOCATION	ANATOMICAL EXTENSION
I	Cerebellar convexity, latero tentorial	Tentorium, transverse and sigmoid sinus
II	CP angle	Petrous ridge, IAC
III	Jugular foramen	Cerebello medullary angle, internal jugular vein, extra cranial
IV	Petro clival	Upper 2/3 rd clivus, cavernous sinus, meckel's cave, petrous ridge
V	Foramen magnum	Lower 1/3 rd clivus and C ₁ , C ₂ area
VI	unclassified	Entire clivus, mid and lower clivus and other types

Takashi M.D.; Ono, Yasuhiro M.D.; Matsumoto, Kengo M.D.; Ohmoto, Takashi M.D. (2001) found that in patients with meningiomas, Tumor infiltration into adjacent brain parenchyma and a

pial-cortical blood supply are critical factors for the development of peritumoral brain edema⁽⁴⁾. In practicality, a balance between good functional outcome and extent of resection is important for posterior cranial fossa meningiomas in proximity to critical structures as shown by Vijayakumar Javalkar, M.D., Anirban Deep Banerjee, M.D., and Anil Nanda, M.D., F.A.C.S in 2012⁽⁵⁾

Andrea Giorgianni, Carlo Pellegrino, Alessandro De Benedictis(2013) found that the dysplastic gangliocytoma is hypointense on unenhanced computed tomographic (CT) images. In such cases, the only diagnostic clue may be the mass effect, which manifests as compression of the fourth ventricle, effacement of the cerebellopontine angle cistern and hydrocephalus⁽⁶⁾.

MRI reveals the characteristic appearance of Lhermitte-Duclos-Disease⁽⁷⁾

MR imaging reveals a cerebellar mass with a typical striated, corduroy, or tiger-striped folial pattern that consists of alternating bands on both T1- and T2-weighted images. The bands are hyper- and isointense relative to gray matter on T2-weighted images and iso- and hypointense on T1-weighted images. Calcification is an uncommon finding, but it has been reported. Atul B.

Shinagare, MD, Nirupama K. Patil, MD, and S. Z. Sorte, MD,(2009) have shown that most dysplastic gangliocytomas do not enhance⁽⁸⁾. Recognition of the disease is of particular importance, as the frequent but under-reported coexistence with Cowden syndrome. clinical and apparative investigation to detect or exclude concomitant malignancies should be prompted⁽⁹⁾. Joachim Klischa, Freimut Juenglinga, Joachim Spreera, Donatus Kocha(2001) found that Although the exact pathophysiologic explanation for the signal characteristics of LDD in diffusion-weighted imaging/perfusion-weighted imaging, 1H MRS, FDG-PET, and 201-Tl SPECT remains unknown, the pathophysiology of this very rare entity may be answered by functional imaging studies of the lesion in future⁽¹⁰⁾

CONCLUSION

Posterior fossa mass lesions are different in adults than children. MRI is the best modality for evaluating posterior fossa lesions highlighted by above cases. Meningioma can occur anywhere in posterior fossa. Dysplastic cerebellar gangliocytoma is a rare entity. Non enhancing solid mass lesion inside cerebellar parenchyma between age of 30 to 50 should raise the possibility of this disorder in the radiologist mind.



Fig2-CT shows good enhancement of mass



Fig3-MRI T₁ W shows hypointense mass of posterior fossa



Fig1-CT shows hyperdense mass in left posterior fossa




Fig4-MRI T₂W shows hyperintense mass lesion




Fig5-MRI FLAIR shows hyperintense mass



Fig8-MRI T₁W shows hypointense mass




Fig6-MRI contrast shows good enhancement




Fig9-MRI T₂W shows hyperintense mass



Fig7-CT shows hypodense mass in midcerebellum




Fig10 MRI FLAIR -hyperintensity with prominent folia like pattern




Fig11 MRI axial contrast-no significant enhancement




Fig12-MRI coronal contrast-no significant enhancement

REFERENCES

1. Pediatric neuroimaging. A. James Barkovich. Philadelphia, PA : Lippincott Williams & Wilkins, c2005. ISBN:0781757665
2. Cerebellopontine Angle Masses:Radiologic-Pathologic Correlation. James G. Smirniotopoulos, MD, Nancy Chang Yue, Lisabetbj Rushing,Radiographics 1993; 13:1131-1 147
3. Posterior fossa meningioma "our experience" in 64 cases,Vernon Velho, VivekAganwal, Rahul Mally, and D. A. Palande, Asian J Neurosurg. 2012 Jul-Sep; 7(3): 116–124.
4. Peritumoral Brain Edema in Intracranial Meningiomas: Effects of Radiological and Histological Factors,Tamiya, Takashi M.D.; Ono, Yasuhiro M.D.; Matsumoto, Kengo M.D.; Ohmoto, Takashi M.D.,Neurosurgery:November 2001 - Volume 49 - Issue 5 - pp 1046-1052
5. Posterior Cranial Fossa Meningiomas* VijayakumarJavalkar, M.D., Anirban Deep Banerjee, M.D., and Anil Nanda, M.D., F.A.C.S. J NeurolSurg B Skull Base. 2012 Feb; 73(1): 1–10.
6. Lhermitte-Duclos Disease A Case Report Andrea Giorgianni, Carlo Pellegrino, Alessandro De Benedictis, Anna Mercuri, Fabio Baruzzi, Renzo Minotto, Antonio Tabano, and Sergio Balbi3 Neuroradiol J. 2013 Dec; 26(6): 655–660.
7. Assessing a Dysplastic Cerebellar Gangliocytoma (Lhermitte-Duclos Disease) with 7T MR Imaging ChristophMoenninghoff, MD, Oliver Kraff, MSc, Marc Schlamann, MD, Mark E. Ladd, PhD, ZazaKatsarava, MD,andElke R. Gizewski, Korean J Radiol. 2010 Mar-Apr; 11(2): 244–248.
8. Dysplastic Cerebellar Gangliocytoma (Lhermitte-Duclos Disease) AtulB.Shinagare, MD, Nirupama K. Patil, MD,and S. Z. Sorte, MD,rsna radiology April 2009,Volume 251, Issue 1
9. Lhermitte-Duclos disease (dysplastic cerebellar gangliocytoma): a malformation, hamartoma or neoplasm?D. A. NowakandH. A. Trost,ActaNeurologicaScandinavica,Volume 105, Issue 3, pages 137–145, March 2002
10. Lhermitte-Duclos Disease: Assessment with MR Imaging, Positron Emission Tomography, Single-photon Emission CT, and MR Spectroscopy,JoachimKlisch, FreimutJuengling, Joachim Spreera, DonatusKocha, TorstenThiela, Martin Bücherta, Sebastian Arnolda, Friedrich Feuerhakea and Martin Schumachera,AJNR 2001 22: 824-830