

# **Original Research Paper**

YOGA

# A Review of Yogic Practices for Leading Risk Factors of Chronic Diseases

G.S. Thangapa	ndiyan	Research scholar, Dept.of Yoga, Karpagam University, Coimbatore Tamilnadu, India.
Dr. A. Mahabo	objan	Professor & Head, Dept.of Physical Education, Bharathidasan University, Tiruchirappalli, Tamilnadu, India.
Yoga, a form of slow to recogniz that take full ac determine the e and high choles Yogic technique studies examine and its related c	Yoga, a form of physical activity is rapidly gaining in popularity and has many health benefits. Yet healthcare providers have been slow to recognize yogic practices and their ability to improve health conditions, and a few Yogic techniques have been developed that take full advantage of its benefits. The purpose of this article is to review published studies using yogic practices and to determine the effect of Yogic techniques on common risk factors of chronic diseases (obesity, high glucose level, hypertension and high cholesterol). A systematic search yielded 50 articles published between 1980 and April 2015. The studies found that Yogic techniques are generally effective in reducing obesity, high glucose level, hypertension and high cholesterol, but only some studies examined long-term adherence. Additionally, not enough studies included diverse populations at high risk for diabetes and its related common health problems.	

KEYWORDS	Yoga – Chronic disease – Risk factors – Obesity, High glucose level, Hypertension and High cholesterol
----------	--

# Introduction

Chronic diseases such as diabetes, heart disease and stroke are leading to causes of death in the world. Common conditions for these chronic diseases are obesity, high glucose level, hypertension and high cholesterol. These conditions are commonly associated with each other. According to a prospective group study focusing on male (men), who had positive relationships between body mass index (BMI) and hypertension occurrence. Men with a higher BMI have a higher likelihood to have diabetes and high cholesterol. Another study with men and women showed that obesity is linked to type 2 diabetes. Even among patients newly diagnosed with type 2 diabetes, the Hypertension in Diabetes Study found that 40% had hypertension, and that hypertension considerably increased mortality in people with type 2 diabetes.

Another recent study found that patients wish a 79% rate of hypertension received outpatient care for type 2 diabetes for at least 2 years. Individuals with impaired glucose tolerance also are more likely to have risk factors for cardiovascular disease such as elevated levels of triglycerides and low-density lipoprotein (LDL) and low levels of high-density lipoprotein (HDL). Promising in this regard is appropriate physical activity because it can reduce body weight, BP, glucose level and cholesterol. Yoga, a form of physical activity consisting of various postures (Asana) and breathing and meditation techniques (Pranayama), has been shown to have therapeutic benefits for individuals with a wide range of health conditions, including hypertension and diabetes. Yoga also appears to be effective in reducing stress and improving exercise tolerance as it is related to cardiovascular response. Yoga also is well received as a therapeutic intervention; for example, participants in a yoga intervention for insomnia found that it was easily learned and performed. Despite its popularity and positive physiologic effects, however, yoga has not been widely recognized in efforts to prevent and treat major chronic health conditions. The purpose of this article is to review published studies using yoga practice to determine the effects of yoga on common health problems, such as obesity, hypertension, high glucose level and high cholesterol.

# Methods

Articles were retrieved from electronic databases (CINAHL, Ovid MEDLINE and PsychInfo) using yoga as a keyword. This initial retrieval, performed in April 2007, yielded about 3000 articles, of

which 861 have been published since 1980. A search within those 861 articles, using the keywords obesity, blood pressure, glucose and cholesterol, identified 56 research articles. After the author had read the full text of the 56 articles and identified those that met the purpose of the study, 24 articles were excluded, mainly because they described outcome variables and characteristics of yoga that were irrelevant. Poor quality was not a reason for exclusion, but studies were excluded if they focused only on meditation or relaxation (Pranayama) or if they were case studies. This process resulted in a final total of 32 articles to be reviewed. The review comprised studies involving yoga interventions and using an experimental or quasi-experimental design. In addition, observational studies were included because they often supply important information beyond the results of clinical trials. Studies that used yoga as a control, not as an intervention, were included as long as they provided evidence of the effectiveness of yoga on variables of interest (Obesity, BP, Glucose and Cholesterol).

# Findings

1) 12 true experimental studies described the randomization method.

2) 18 described quasi-experimental studies.

3) 2 described observational studies.

4) Even though risk factors for chronic health conditions were used as keywords in the search strategy,

5) Many of the studies used healthy adult samples; only half of the 32 studies actually focused on subjects with diabetes ,

6) Hypertension or Cardiovascular disease. The studies were conducted in the US, India and other countries.

# **Content of Yogic Practices**

Other than duration and frequency, most articles did not describe the yoga sessions in detail. Only a few articles provided details of the yoga sequences used. Some listed the names of postures and breathing techniques. Of the few articles that identified the type of yoga studied, the most common was Hatha yoga, a popular form of yoga in the Western world. The Compendium of Physical Activities, a coding scheme that classifies physical activity based on energy expenditure, does not define energy expenditure while performing various types of yoga. However, Hatha yoga is stretching, which has a metabolic equivalent (MET) of 2.5 (1 MET equals the amount of energy used for resting). A recent study found that when young adult women (19 to 40-years old) performed a 30-min session of Hatha yoga, the MET was 2.17. However, if the session was made more active using Sun Salutations, a flowing series of physical postures, the MET increased to 3.74 (SD= 0.70).

### **Frequency and Duration of Yoga Sessions**

The total dose of yoga training, which depends on both the duration and frequency of yoga sessions, also needs to be considered in evaluating and comparing yoga studies. The most common duration and frequency of yoga sessions in the studies were 30-60 min per session and sessions meeting daily for 4-10 weeks, but many studies used meeting sessions 2-3 times per week for 8–12 weeks. Some Yogic practices met more frequently than others but for a shorter time span. For example, in a study by Damodaran et al. persons with essential hypertension received 1 h of yoga training daily for 3 months, which is counted as 84 h of training, whereas Ray et al. Healthy young adults who received 1 h of yoga training three times a week for 10 months, for a total of 120 h. Some findings can be useful in trying to determine effective durations of yoga sessions. For example, in one study, 1 h daily yoga practice was associated with significant reductions in body weight and cholesterol levels after 4 weeks, and those significant effects lasted for 14 weeks. In a study of 20 patients with essential hypertension, daily 30-min sessions of yoga led to a decrease in BP by the fifth day.

#### **Yogic Practices**

Any persistent benefits from yoga would rely on long-term adherence, which was examined in only in a few of the reviewed studies. In some studies, after yoga training, subjects were asked to continue their programs with daily home practice. One of these compared adherence to yoga practice after a 10-week yoga intervention between white and black American adults (mean age: 69 years for whites and 70 years for blacks). Although the drop out rate did not differ significantly (16% for whites, 22% for blacks), the black participants did not engage in yoga at home as frequently as whites. A different study found greater compliance with subsequent home practice among participants in a yoga class that met three times per week than in those who attended a weekly yoga class (86% versus 65%, P< 0.05). However, found difference was found in compliance with home practice between groups engaged in yoga or aerobic exercise.

# Weight Decreased

According to a retrospective observational study of 15,550 adults aged 53–57 years, regular yoga practice for 4 or more years was significantly associated with weight loss by obese participants. Several intervention studies also showed that yoga practice was effective in reducing body weight. After a 4-day residential yoga practice followed by 14 weeks of 1 h daily home practice, one study found a significant loss in mean body weight from 72.26 to 70.48 kg among subjects with risk factors for coronary artery disease (CAD). Other studies found that yoga was associated with significant weight loss by subjects with CAD and subjects without CAD . Manchanda et al. Show a 7% loss of body weight among adult men with CAD after 1 year of yoga practice, and in a study by Schmidt and colleagues, healthy adults lost an average of 5.7 kg after 3 months of yoga practice. All obese adults studied by Yogendra et al. reached a normal weight within 1 year after initiating yoga-based lifestyle modifications. However, this article did not show the effect size of this change.

#### **Blood Glucose Level Lowered**

Of the six studies that examined blood glucose, all found that yoga was effective in reducing blood glucose level. For example, after 4 months of yoga practice, fasting glucose fell significantly (from 144 to 119 mg dl1, P< 0.005) in a group of adults with type 2 diabetes, Another sample of 24 adults with type 2 diabetes had significantly decreased fasting glucose (from 190.1 to 141.5 mg dl1, P< 0.001) after 40 consecutive days of yoga practice, Adults with normal blood glucose levels also had significantly lower glucose levels after 3–4 h of yoga practice for 8 days (P < 0.001).

# **Bood Pressure Decreased**

This review found ample evidence that yoga was effective in reducing BP. In a group of low-income elderly people, effects on systolic BP did not differ between a yoga class and an aerobic exercise class, both held three times a week for 10 weeks. However, eight other studies found that yoga practice was effective in lowering BP in healthy samples, regardless of the type of yoga. Yoga practice also significantly improved BP among people with hypertension, cardiovascular disease or type 2 diabetes. For example, in 13 patients, aged 41–60 years, with essential hypertension, BP dropped significantly during the third week of a 4-week yoga program (1 h per day, 6 days per week), and it fell further after the program. For example, systolic BP dropped from 141.7 to 127.9 mmHg by the third week and to 120.7 mmHg by the fourth week.

# **Cholesterol Level Improved**

The practice of yoga was associated with significant decreases in cholesterol among subjects with cardiovascular disease, hypertension or type 2 diabetes. One study examined a regimen involving 4 days of a yoga program at a residential course, followed by 1 year of yoga practice at home. In both men with angina and asymptomatic participants with CAD risk factors, all lipid variables except HDL decreased beginning the fourth week of yoga practice (e.g. total cholesterol fell from 206.6 to 193.6 mg dl1), and the level of total cholesterol continued falling to 176.06 mg dl1 at 14 weeks. A study of subjects at risk for cardiovascular disease and diabetes found significant improvements (P < 0.01) in total cholesterol, triglycerides, LDL, HDL and very-LDL (VLDL, defined as total cholesterol minus LDL minus HDL) after shortterm intensive yoga practice (3–4 h per day for 8 days). Notably, for subjects whose baseline total cholesterol was 200 mg dl1 or higher, the reduction in triglycerides (from 151.5 48.9 to 132.7 50.5 mg dl1, P < 0.001) and VLDL (from 36.7 13.8 to 30.2 14.6 mg dl1, P < 0.001) was significantly greater than in subjects with lower baseline total cholesterol (triglycerides falling from 113.6 46.5 to 110.5 38.1 mg dl1, P> 0.05; VLDL from 23.7 12.8 to 23.2 12.5 mg dl1 , P > 0.05). Finally, a study of healthy adults over 40 years old found that 5 years of yoga practice reduced age-related deterioration in cardiovascular functions. Although the article describing this observational study did not detail the type of yoga performed nor the frequency or intensity of the yoga sessions, the data showed a long-term change indicating the effectiveness of yoga on cardiovascular functioning.

Four studies used aerobic training as an intervention and yoga as a control. One of these, a study of healthy active people aged 65 years and older, found that aerobic exercise produced no significant reduction in weight or BP, whereas 4 weeks of yoga practice did lead to some reduction in weight or BP (for example, systolic BP decreased from 146 to 139 mmHg). DiPietro et al. found no change in glucose and insulin responses in their yoga control group. However, the yoga control group was monitored to ensure that pulse rates did not exceed 90 beats per min during yoga practice. Therefore, this restriction should be considered in interpreting this result.

#### Discussion

The reviews showed that yoga had beneficial effects on body weight, BP, blood glucose level and cholesterol level. Nonetheless, several shortcomings in research on this topic need to be addressed, in order for clinical programs to capitalize on these health benefits.

The articles describing interventions, only a few provided details regarding the Yogic practices or the names and sequence of yoga postures. Asana and Pranayama provide different types of health benefits, and therapeutic Yogic practices can involve various combinations of these two components. An optimal comparison of study results thus requires knowing the combination of Asana and Pranayama used. The sequence of yoga postures can be inferred from the list of posture names, but more straightforward information is essential for future studies seeking to replicate or generalize the results. A related issue that remains to be resolved is

how to standardize yoga exercises for research purposes. The optimal duration and intensity required to maximize the effectiveness of yoga need to be determined, as does the need for a booster to provide long-term effects. Because the MET of yoga is low, increasing the frequency may increase the benefits. However, these factors cannot be judged from the review studies; many articles did not clearly identify the dosage of the yoga program studied, and they focused on the short-term health benefits of yoga. Only a few studies included follow-up data beyond 6 months. It also remains to be determined whether more intensive training, that is, a greater dosage, improves the likelihood of adopting and maintaining active behavior in the long run.

The samples in the reviewed studies pose additional dilemmas. It was not a surprise that a large portion of the studies were conducted in India, where the philosophy and practice of yoga originated. However, this focus on one geographical region, where yoga is particularly ingrained in the culture, limits the generalizability of results. A few studies have addressed variables of interest specific to minorities in the US (Blacks, Hispanics and Asians), which are populations believed to be more vulnerable to type 2 diabetes and physical inactivity.

Yoga has beneficial effects on various health conditions. A large portion of the reviewed studies analyzed the effects of yoga with healthy samples. There is evidence that yoga practice was more effective in lowering triglycerides among people with higher cholesterol than those with a lower cholesterol level. Therefore, it is important to consider participants' health conditions because there are possible differences in the effects of yoga by their health severity.

#### Conclusions

Analyses of yoga intervention should be designed and conducted to identify programs best suited for diverse populations and for specific populations with high-risk factors for chronic health conditions. Such studies could guide the development of more practical and effective interventions.

#### REFERENCES

- American Diabetes Association. The prevention or delay of type 2 diabetes. Diabetes Care 2003;26:S62–9.
- Gelber RP, Gaziano JM, Manson JE, Buring JE, Sesso HD. A prospective study of body mass index and the risk of developing hypertension in men. Am J Hypertens 2007;20:370–7.
- Sullivan PWP, Morrato EHM, Ghushchyan VM, Wyatt HRM, Hill JOP. Obesity, inactivity and the prevalence of diabetes and diabetes-related cardiovascular comorbidities in the U.S. 2000–2002. Diabetes Care 2005;28:1599–603.
- HDS Study Group. Hypertension in Diabetes Study (HDS): II.Increased risk of cardiovascular complications in hypertensive type 2 diabetic patients. J Hypertens 1993;11:319–25.
- Choe HM, Townsend KA, Blount G, Lo CH, Sadowski L, Standiford CJ. Treatment and control of blood pressure in patients with diabetes mellitus. Am J Health Syst Pharm 2007;64:97.
- The Diabetes Prevention Program Research Group. Lipid, lipoproteins, C-reactive protein, and hemostatic factors at baseline in the diabetes prevention program. Diabetes Care 2005;28:2472–9.
- U.S. Department of Health and Human Services. Physical activity and health: a report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, 1996.
- U.S. Department of Health and Human Services. Healthy People 2010. (cited 2004 Feb 4) 2001. Available from: http://
- www.healthypeople.gov/Document/HTML/Volume2/22Physical.htm.
- Nayak NN, Shankar K. Yoga: a therapeutic approach. Phys Med Rehab Clin N Am 2004;15:783–98.
- McCaffrey R, Ruknui P, Hatthakit U, Kasetsomboon P. The effects of yoga on hypertensive persons in Thailand. Holistic Nurs Prac 2005;19:173–80.
- Bijlani RL, Vempati RP, Yadav RK, Ray RB, Gupta V, Sharma R, et al. A brief but comprehensive lifestyle education program based on yoga reduces risk factors for cardiovascular disease and diabetes mellitus. J Altern Complem Med 2005;11:267–74.
- Kreitzer MJ, Gross CR, Ye X, Russas V, Treesak C. Longitudinal impact of mindfulness meditation on illness burden in solid-organ transplant recipients. Prog Transplant 2005;15:166–72.
- Madanmohan, Udupa K, Bhavanani AB, Shatapathy CC, Sahai A. Modulation of cardiovascular response to exercise by yoga training. Indian J Physiol Pharmacol 2004;48:461–5.
- Tindle HA, Davis RB, Phillips RS, Eisenberg DM. Trends in use of complementary and alternative medicine by US adults: 1997–2002. Altern Ther Health Med 2005;11:42–9.
- Khalsa SB. Treatment of chronic insomnia with yoga: a preliminary study with sleepwake diaries. Appl Psychophysiol Biofeedback 2004;29:269–78.
  Kunz R, Oxman AD. The unpredictability paradox: review of empirical comparisons
- Kunz R, Oxman AD. The unpredictability paradox: review of empirical comparisons of randomised and non-randomised clinical trials. Br Med J 1998;317:1185–90.

- Harinath K, Malhotra AS, Pal K, Prasad R, Kumar R, Kain TC, et al. Effects of Hatha yoga and Omkar meditation on cardiorespiratory performance, psychologic profile, and melatonin secretion. J Altern Complem Med 2004;10:261–8.
- Latha, Kaliappan KV. Yoga, pranayama, thermal biofeedback techniques in the management of stress and high blood pressure. J Indian Psychol 1991;9:36–46.
- Agte VV, Tarvadi K. Sudarshan Kriya yoga for treating type 2 diabetes: a preliminary study. Altern Complem Ther 2004;10:220–2.
- Malhotra V, Singh S, Singh KP, Gupta P, Sharma SB, Madhu SV, et al. Study of yoga asanas in assessment of pulmonary function in NIDDM patients. Indian J Physiol Pharmacol 2002;46:313–20.
- Malhotra V, Singh S, Tandon OP, Madhu SV, Prasad A, Sharma SB. Effect of Yoga asanas on nerve conduction in type 2 diabetes. Indian J Physiol Pharmacol 2002;46:298–306.
- Singh S, Malhotra V, Singh KP, Madhu SV, Tandon OP. Role of yoga in modifying certain cardiovascular functions in type 2 diabetic patients. J Assoc Physicians India 2004;52:203–6.
- Malhotra V, Singh S, Tandon OP, Sharma SB. The beneficial effect of yoga in diabetes. NMCJ 2005;7:145–7.
- Damodaran A, Malathi A, Patil N, Shah N, Suryavansihi, Marathe S. Therapeutic potential of yoga practices in modifying cardiovascular risk profile in middle aged men and women. J Assoc Physicians India 2002;50:633–40.
- Vijayalakshmi P, Madanmohan, Bhavanani AB, Patil A, Babu K. Modulation of stress induced by isometric handgrip test in hypertensive patients following yogic relaxation training. Indian J Physiol Pharmacol 2004;48:59–64.
- Murugesan R, Govindarajulu N, Bera TK. Effect of selected YOGIC PRACTICES on the management of hypertension. Indian J Physiol Pharmacol 2000;44:207–10.
- Selvamurthy W, Sridharan K, Ray US, Tiwary RS, Hegde KS, Radhakrishan U, et al. A new physiological approach to control essential hypertension. Indian J Physiol Pharmacol 1998;42:205–13.
- Mahajan AS, Reddy KS, Sachdeva U. Lipid profile of coronary risk subjects following yogic lifestyle intervention. Indian Heart J 1999;51:37–40.
- Manchanda SC, Narang R, Reddy KS, Sachdeva U, Prabhakaran D, Dharmanand S, et al. Retardation of coronary atherosclerosis with yoga lifestyle intervention. J Assoc Physicians India 2000;48:687–94.
- Sivasankaran S, Pollard-Quintner S, Sachdeva R, Pugeda J, Hoq SM, Zarich SW. The effect of a six-week program of yoga and meditation on brachial artery reactivity: do psychosocial interventions affect vascular tone? Clin Cardiol 2006;29:393–8.
- Yogendra J, Yogendra HJ, Ambardekar S, Lele RO, Shetty S, Dave M, et al. Beneficial effects of yoga lifestyle on reversibility of ischaemic heart disease: camp heart project of International Board of Yoga. J Assoc Physicians India 2004;52:283–9.
- Jemal A, Ward E, Hao Y, Thun M. Trends in the leading causes of death in the United States 1970–2002. JAMA 2005;294:1255–9.