



## A Randomised Study on Outcome of Dynamic Hip Screw Fixation Versus Helical Hip Screw System In Intertrochanteric Fracture

**DR.SURENDRA  
KUMAR PADARYA**

P.D. Hinduja National Hospital and Medical Research Centre,  
Mumbai

**DR.ASHWANI  
BILANDI**

P.D. Hinduja National Hospital and Medical Research Centre,  
Mumbai

### ABSTRACT

**Introduction :** Trochanteric fractures occur in the younger population due to high velocity trauma, whereas in the elderly population it is most often due to trivial trauma. In United States 49% of hip fractures are inter trochanteric. The most commonly used device for fixation is the Dynamic Hip Screw with Side Plate assemblies. We aimed to compare outcome of dynamic hip screw fixation versus helical hip screw system in intertrochanteric fracture

**Material methods :** this was a prospective study conducted at P. D. Hinduja National Hospital and M R C, Mumbai. 25 patients with comparable characteristics in each the DHS group and the spiral blade plate group were selected for the study randomizing them into the 2 groups for comparison. Radiological outcome on the basis of union, complications with respect to implant cut-out and malalignment and Functional outcome on the basis of Harris Hip Score were recorded.

**Results:** The mean time taken for surgery in the DHS group (in mins) mean was 57.44(SD+9.305), and in the DHHS group mean was 48.44(SD+ 6.740) (p 0.00014). There was no significant statistical difference (p=0.399) in The mean time taken for union(in weeks) There was no failure or complication in both groups.

**Conclusion:** Although the mean operative time was almost 10 mins more in the DHS group, there is no statistically significant difference in the out come between the DHS group and DHHS group.

### KEYWORDS

Intertrochanteric fracture , Dynamic Hip Screw(DHS) , Dynamic helical hip screw system(DHHS).

### INTRODUCTION:

Trochanteric fractures are common in the elderly people. The frequency of these fractures has increased primarily due to the increasing life span and more active life style brought on by urbanization. Trochanteric fractures occur in the younger population due to high velocity trauma, whereas in the elderly population it is most often due to trivial trauma.<sup>1</sup> The incidence of trochanteric fractures is more in the female population compared to the male due to osteoporosis. In 1990, the number of hip fractures in persons aged 50 years or older was 1.7 million worldwide<sup>1</sup> and data from the United States suggest that 49% of hip fractures are intertrochanteric<sup>2</sup>. The trochanteric fractures can be managed by conservative methods and there is usually union of the fracture. If suitable precautions are not taken the fracture undergoes malunion, leading to varus and external rotation deformity, shortening and limitation of hip movements<sup>7</sup>.

There are various forms of internal fixation devices used for trochanteric fractures; of them the most commonly used device is the Dynamic Hip Screw with Side Plate assemblies. This is a collapsible fixation device, which permits the proximal fragment to collapse or settle on the fixation device, seeking its own position of stability. In spite of the advances in anaesthesia, nursing care and the surgical techniques, hip fractures remain a significant cause of morbidity and mortality in the elderly population. Clawson introduced the dynamic hip screw (DHS) in 1964 and has been widely adopted as the implant of choice for these fractures. The failure rates associated with the DHS vary between 5-23%. However, more recent studies have reported improved results.<sup>9</sup>

It is well known that accurate positioning of the cephalic screw of a sliding hip screw (SHS) or intramedullary (IM) nail in the femoral head determines the outcome following open reduction internal fixation of trochanteric fractures<sup>5,6,7</sup>. The most common mode of failure of fixation is cut-out of the lag screw from the femoral head<sup>8</sup>, and it has been shown<sup>5,6,7,9,10</sup> that the tip-apex distance (TAD) is the most valuable factor in

determining the likelihood of lag screw cut-out, with a distance >25 mm an indicator of unsatisfactory screw placement and a statistically increased rate of cut-out. In addition to TAD the position of the cephalic screw in one of nine zones in the head has been described<sup>11</sup>. The ideal position is centre-centre, but a short screw which is centre-centre may still allow for a TAD >25 mm. Thus, it is both the correct length of the screw within the head, as well as the central position of the screw, that will help to prevent cut-out of the cephalic implant. Since the position of the implant in the head is dependent upon the initial guide wire, and since the position of the guide wire is solely under the control of the surgeon, proper placement of any of these implants is the best way to ensure a satisfactory outcome<sup>12,13</sup>. However, even with a very low rate of fixation failure and lag screw cut-out there has been a shift in the type of cephalic implant that is used with either a SHS or IM nail. A helical blade has been introduced for use with either of the two implants, with the idea that there will be an improved hold in the femoral head and a reduced rate of cut-out. Although there are biomechanical studies<sup>13,14</sup> which give support to this idea, it has not been confirmed by clinical studies. Additionally, there is no information in the literature concerning any difference between a helical blade and screw with respect to placement of the implant within the femoral head. In view of these considerations, the present study of Surgical Management of Peritrochanteric Fractures is taken up.

### MATERIAL AND METHODS

The study was a prospective one where the fractures were classified on the basis of AO system as 31-A1/ 31-A2. The surgeries were performed at P. D. Hinduja National Hospital and M R C, Mumbai with surgeon whose experience in the field of orthopaedic traumatology was factored in. Informed consent was obtained from all patients before the operation. The patients operated from Aug 2011- June 2013 with DHS and DHHS (spiral blade plate) were included. 25 patients in each the DHS group and the spiral blade plate group were selected for the study randomizing them into the 2 groups for comparison. The many characteristics of both the groups of the

patients were comparable. All fractures operated with a plate using either a sliding hip screw or a helical blade, which were less than 10 days old and in patient older than 18 years of age were included. Reverse intertrochanteric fracture and patients who had a concomitant ipsilateral /contralateral other lower limb injuries were excluded.

Management of patients: As soon as the patient with suspected intertrochanteric fracture was seen, necessary clinical and radiological evaluation was performed and necessary resuscitation and splintage using skin traction was. All routine pre-operative investigations were done along with X-rays of Pelvis with both hips-AP view, Involved side hip with femur full length-AP and Lateral view in all patients.

Associated injuries were evaluated and treated simultaneously. All the patients were operated on elective basis. The implant choice was independent of age, sex and fracture configuration. All implants used belonged to the Depuy Synthes. A standard operative technique (lateral approach) used for all cases. The duration of the surgery from skin incision to skin closure was recorded. The following outcomes were measured:

- (1) Radiological outcome on the basis of union, complications with respect to implant cut-out and malalignment and
- (2) Functional outcome on the basis of Harris Hip Score.

Radiological assessment was done intra-operatively in terms of guidewire placement and the screw size and location. This assessment was purely based on the surgeon's experience and the instrumentation used in the system. The second assessment was done immediately post-operatively in terms of reduction, zone of screw placement and TAD. It was accurate and was done on the basis of PACS (picture archiving and communications systems) which was available in the institute.

The post-operative protocol was similar for both sliding screw and the spiral blade. The patient was asked to bear full weight on both the limbs with the help of a walker. If the first 3-4 post-operative days were uneventful, they were discharged and asked to follow-up at 10 days post operation. The patient was asked to avoid squatting/sitting cross-legged. They were advised to stop using walker and start complete free weight bearing when tolerated. Serial post operative X-rays were done and the same parameters were followed in 1 month, 3 months, 6 months and at 1 year. At 6 months follow-up, the functional outcome was assessed on the basis of Harris Hip Score. A questionnaire was prepared on the basis of Harris Hip Score and given to the patient. Also, the surgeon evaluated the objective points of assessment in Harris Hip Score and noted them down. The overall outcomes of both the modalities of fixation were then assessed and the results were derived

## RESULTS:

In our study 50 patients with intertrochanteric fracture were randomised for treatment, 25 patients in each group. The DHS and DHHS (Helical Blade Plate) group were similar statistically in demographical (age, sex) and surgical. In this study age group in both groups is almost similar mean age in DHS group was 76.04 (SD +10.03) and in DHHS group was 78.36(SD +10.39). In this study we compared time taken for surgery, tip apex distance and time taken for union (Radiologically). The DHS group had mean Tip Apex Distance (TAD) of 17.05 mm (SD + 4.633 ) and in DHHS group mean TAD was 19.40 mm (SD + 5.430 ). There was a statistical difference between the TADs of the two groups ( $p=0.046$ ). The mean time taken for surgery in the DHS group (in mins) mean was 57.44(SD+9.305), and in the DHHS group mean was 48.44(SD+ 6.740). Statistically difference in these group was significant ( $p=0.00014$ ). The mean time taken for union (in weeks) in the DHS group was 16.21(SD+3.15) and in the DHHS group was 16.45 (SD+3.05). There was no significant statistical difference ( $p=0.399$ ). There was no failure in both groups (DHS and DHHS). Postoperatively there was no complications like superficial or deep infection, implant cut out, re-

vision or segmental osteonecrosis. At 6 months mean Harris Hip Score was in the DHS group was 84.5 (SD +8.153) and in the DHHS group 82.744 (SD +11.78), which statistically had no difference ( $p=0.288$ ).

**TABLE 1 : DHS GROUP**

	Mean	Range	SD
AGE (years)	76.04	66-88	+ 10.03
TIME TAKEN FOR SURGERY (mins)	57.44	40-75	± 9.305
TIME TAKEN FOR UNION(weeks)	16.21	12-22	+ 3.15
HHS	84.5	62.7-92.4	+ 8.15
TAD (mm)	17.05	10.2-29.6	+ 4.633

**TABLE : HELICAL BLADE PLATE**

	Mean	Range	SD
AGE (years)	78.36	61-97	+10.03
TIME TAKEN FOR SURGERY (mins)	48.44	36-60	9.305
TIME TAKEN FOR UNION (weeks)	16.45	12-22	+ 3.05
HHS	82.744	61.6-94.6	± 11.78
TAD (mm)	19.40	13-30.8	+ 5.430

## DISCUSSION

Intertrochanteric hip fractures are common and resulted in heavy burden to affected patients<sup>15</sup>. Sliding hip screws have been widely used but it is associated with problems<sup>16</sup>. There have been various biomechanical studies comparing the strength of helical blade plate and DHS screw.<sup>13,14</sup>. Ekmke et al<sup>17</sup> measured forces and energy required to insert DHS screw and DHHS blade and found that total energy required for DHS screw insertion is much higher than DHHS blade. Kyle et al evaluated sliding characteristics of different versions of DHS screws. They noted 4 factors in the initiation of sliding, including coefficient of friction between shaft and barrel, length of screw out of barrel, length of contact between barrel and shaft of screw, and force acting perpendicular to the screw<sup>18</sup>. Our study didn't have any exclusion criteria as far as age was concerned. Also the quality of bone was not considered a deciding factor for the choice of implant. A large prospective randomized series done by Stern et al<sup>13</sup> wasn't able to demonstrate any difference in complications of a blade vs a screw in terms of re-operation rates and cut outs in 1 year. The series included blade vs screw involving both the plate and nail. It did not address issues like the quality of reduction, time taken for the surgery, incidence of varus malunion and time taken to heal<sup>19</sup>. Also, considering it was a prospective study, it didn't take into account intra-operative factors like the time taken for the surgery and immediate post-operative morbidity for the two modalities of fixation. Also there was no difference in time taken for union between 2 groups. On an average time taken for union in our study was 16 weeks. In our study we included time taken for surgery, which has significant difference ( $p=0.046$ ), DHHS group took less time than DHS group though time taken for union was insignificant ( $p=0.399$ ). In our study TAD in DHS group was 17.05 mm and in DHHS group was 19.40 mm which was statistically insignificant but clinically irrelevant. Baumgaertner et al<sup>6,10</sup> demonstrated in their series that if the TAD was within 25 mm there wouldn't be any cut outs of the screw and this variable would be by far the most important predictor of outcomes of screw. In our study TAD was within 25 mm except in 5 patients where TAD was more than 25 mm (3 in DHHS group and 2 in DHS group). However there was no complication in any of these cases.

## CONCLUSION:

In conclusion our study shows that, there is no difference in the clinical outcomes between the DHS group and DHHS group. However the mean operative time was almost 10 mins more in the DHS group. There was no complication in both the groups. The supposed biomechanical stability of the spiral blade plate has not been converted to a definite clinical superiority in our study. Key to success is maintaining the Tip - apex distance to within 25 mm Consistent with the Bau-

emgarter series with the DHS. However it has to be kept in mind that cost of Helical blade plate is significant higher than DHS screw.

#### REFERENCES:

1. J.David, G.Lavelle. Fractures and dislocations chapter-52 in CAMPBELL'S OPERATIVE ORTHOPAEDICS, eleventh edition.VOL-3 pages; 3237-3308.
2. Kyle Richard F, Campbell Sara J. intertrochanteric fractures,chapter-40 in MICHAEL W.CHAPMAN OPERATIVE ORTHOPAEDICS.VOL-1,pages 600-603.
3. "The association of age, race and sex with the location of proximal femoral fractures in elderly". JBJS 1993;75(5),752-9.
4. Boyd H B, GRIFFIN "Classification and treatment of trochanteric fractures" Arch surgery, 1949;58;853-866.
5. Barton TM, Gleeson R, Topliss C, Greenwood R, Harries WJ, Chesser TJ (2010) A comparison of the long gamma nail with the sliding hip screw for the treatment of AO/OTA 31-A2 fractures of the proximal part of the femur: a prospective randomized trial. J Bone Joint Surg Am 92:792-798
6. Baumgaertner MR, Solberg BD (1997) Awareness of tip-apex distance reduces failure of fixation of trochanteric fractures of the hip. J Bone Joint Surg Br 79:969-971 3. Davis TR, Sher JL, Horsman A, Simpson M, Porter BB, Checketts RG (1990) Intertrochanteric femoral fractures. Mechanical failure after internal fixation. J Bone Joint Surg Br 72:26-31.
7. Geller JA, Saifi C, Morrison TA, Macaulay W (2010) Tip-apex distance of intramedullary devices as a predictor of cut-out failure in the treatment of peritrochanteric elderly hip fractures. Int Orthop 34:719-722.
8. Kyle RF (1994) Fractures of the proximal part of the femur. J Bone Joint Surg Am 76:924-950.
9. Walton M, Barnett A, Jackson M (2008) Tip-apex distance as a predictor of failure following cephalo-medullary fixation for unstable fractures of the proximal femur. Eur J Trauma Emerg S 34:273-276.
10. Baumgaertner MR, Curtin SL, Lindskog DM, Keggi JM (1995) The value of the tip-apex distance in predicting failure of fixation of peritrochanteric fractures of the hip. J Bone Joint Surg Am 77:1058-1064.
11. Cleveland M, Bosworth DM, Thompson FR, Wilson HJ Jr, Ishizuka T (1959) A ten-year analysis of intertrochanteric fractures of the femur. J Bone Joint Surg Am 41:1399-1408
12. Sahni V, Sureen S, Shetty V (2001) Who is responsible for failure of a dynamic hip screw by cutout, the patient or the surgeon? In: Proceedings of the 68th annual meeting of the American Academy of Orthopaedic Surgeons, vol 2, p 385
13. Stern R (2007) Are there advances in the treatment of extracapsular hip fractures in the elderly? Injury 38(Suppl 3):S77-S87
14. Al-Munajjed AA, Hammer J, Mayr E, Nerlich M, Lenich A (2008) Biomechanical characterisation of osteosyntheses for proximal femur fractures: helical blade versus screw. Stud Health Technol Inform 133:110.
15. Thakar C, Alsousou J, Hamilton TW, Willett K. The cost and consequences of proximal femoral fractures which require further surgery following initial fixation. J Bone Joint Surg Br 2010;92:1669-77
16. Crawford CH, Malkani AL, Cordray S, Roberts CS, Sligar W. The trochanteric nail versus the sliding hip screw for intertrochanteric hip fractures: a review of 93 cases. J Trauma 2006;60:325-29.
17. Ehmke L, Kam B, Sommers M, Bottlang M. Screw and Blade type implants produce distinct insertion forces. 4 world congress of biomechanics, 2004.
18. Kyle R, Wright T, Burstein A, Biomechanical analysis of sliding characteristics of compression hip screw. The journal of Bone and Joint Surgery 1980;62;1308-14.
19. Kancherla R, Sankineni S, Trikha V, Kumar R, Malhotra R (2012). Comment on Stern et al.: Prospective randomised study comparing screw versus helical blade in the treatment of low-energy trochanteric fractures. Int Orthop 36: 1109 - 1110.