



ORIGINAL RESEARCH PAPER

Radiodiagnosis

ROLE OF ULTRASONOGRAPHIC AND LAPAROSCOPIC EXAMINATION IN CHRONIC PELVIC PAIN

KEY WORDS: Chronic pelvic pain, clinical examination, ultrasonography, laparoscopy.

Dr. Mukhtar Ahmad

Radiologist, Department of Radiology, Rajiv Gandhi Cancer and Research Institute, New Delhi.

Dr. Nahida Hassan*

Consultant, Department of Obstetric and Gynecology, Rajiv Gandhi Cancer and Research Institute, New Delhi. *Corresponding Author

Dr. Dipesh Yadav

Radiologist, Department of Radiology, Rajiv Gandhi Cancer and Research Institute, New Delhi.

ABSTRACT

Background : Chronic pelvic pain is defined as pain localized to pelvis or lower abdomen below the line joining the two anterior superior iliac spines of at least 6months duration .Chronic pelvic pain is a common medical problem affecting men and women .
Aims and Objective: 1. To find out the correlation and compare the efficacy of ultrasonographic examination and diagnostic laparoscopy
Materials and Methods: This prospective study was conducted Rajiv Gandhi cancer and research institute,New Delhi. The study was conducted over a period of one year and total number of 100 patients were enrolled of age group 20-50yrs with pelvic pain of >6months duration.
Results: Maximum number of cases with chronic pelvic pain belonged to age group of 31-40yrs.Mean age of patients was 34.2+69 yrs. Mean duration of pain was 30.3+_25.67(7,120) months. The associate complaints were dysmenorrhea(50%), menorrhagia (24%) and infertility(10%). Among the 87 patients with abnormal findings on laparoscopic examinations 58 had abnormal clinical findings. Laparoscopy was more predictive in evaluation causes of CPP such as pelvic inflammatory disease (36%), pelvic adhesions(27%),endometriosis(29%),and pelvic congestions(13%).
Conclusion: Diagnostic laparoscopy is more sensitive method for evaluation of chronic pelvic pain.

INTRODUCTION

Chronic pelvic pain is defined as pain localized to pelvis or lower abdomen below the line joining the two anterior superior iliac spines of at least 6 months duration¹.Although no universally accepted definition of chronic pelvic pain exists more clinicians agree that chronic pelvic pain is characterized by pain that is menstrual or non-menstrual in nature, lasting 6 months or more, localized below the umbilicus or in pelvis, severe enough to cause functional disability or require medical or surgical treatment². Chronic pelvic pain is a more of constant or intermittent, cyclic or acyclic pelvic pain^{3,4}. Chronic pelvic pain is best classified as a symptom and not a disease.¹

Prevalence of chronic pelvic pain has been reported as 3.8% in women aged 15-73 years.^{5,6} Prevalence rate increased with age from 18.2/1000 in 15-20 years old to 27.6/1000 women >60 years of age.^{7,8,9} Symptoms persist longer in older age group.^{8,9}

The causes of chronic pelvic pain may be somatic or non-somatic (psychogenic).¹⁰ The causes of chronic pelvic pain can be broadly divided into gynecological and non-gynecological causes.¹¹

GYNAECOLOGICAL CAUSES

Gynecological causes account for 90% of chronic pelvic pain when there is specific diagnosis.⁹

(I) UTERINE CAUSES:

- (a) Adenomyosis.² The incidence of adenomyosis ranges from 5% to 70%.¹²
- (b) Leiomyomas^{2,13} Uterine fibroids have been detected in 5.2% to 8.9% in women who had undergone laparoscopy for chronic pelvic pain.¹³
- (c) Chronic endometritis.¹⁴
- (d) Chronic cervicitis¹⁴
- (e) Prolapse¹¹
- (f) Retroverted uterus^{2,11}
- (g) Intrauterine contraceptive device¹⁴

(ii) EXTRA-UTERINE CAUSES

OVARY

- Benign ovarian cyst.²
- Polycystic ovarian disease.²

Endometrioma: is most common morbidities associated with chronic pelvic pain approximately 70%.^{15,16,17}

Ovarian remnant syndrome Cyclic pain is associated with development of ovarian follicles within hormonally active ovarian tissue.^{18,19}

Ovarian retention syndrome (residual ovarian syndrome) occurs in a patient who has undergone hysterectomy with ovarian conservation.^{18,19}

PELVIS

Endometriosis^{2,17}

Pelvic congestion syndrome Pelvic varicosities causing chronic pelvic pain were first described by Richet in 1857 and elaborated by Beard in 1884²⁰.

Pelvic adhesive disease is the result of previous surgery or infection.²¹ Adhesions are diagnosed in approximately 25% of women with chronic pelvic pain.^{4,14}

Chronic pelvic inflammatory disease Caused by the gonorrhea, Chlamydia, mycobacterium tuberculosis and IUCD.¹⁴

OTHER CAUSES

- (a) Valvular vestibulitis^{22,23}
- (b) Vulvodynia²³
- (c) Malignancies of female genital tract.²

NON-GYNAECOLOGICAL

(I) UROLOGICAL CAUSES

- (a) Chronic urinary tract infection is one of the most common causes of irritative voiding syndrome in women.²⁴
 - (b) Interstitial Cystitis is a chronic inflammatory condition of the bladder presenting as lower abdominal pain.^{24,23}
 - (c) Subureteral diverticulitis the classic triad of symptoms is dysuria, dysparenia and post void dribbling.²⁴
 - (d) Urethral symptoms are a symptom complex that can include dysuria, urinary frequency, urgency, suprapubic discomfort and straguria.^{2,23}
 - (e) Malignancy.
- (ii) GASTROINTESTINAL CAUSES:**

- (a) Chronic Appendicitis: Presented with localized right lower quadrant pain unresponsive to conservative treatment.¹¹
- (b) Diverticular Disease: Commonly found in sigmoid colon. Infection of diverticuli and micro-abscess formation present as abdominal pain and tenderness.^{2,11}
- (c) Inflammatory bowel disease: Due to ulcerative colitis and Cohn's disease may be associated with chronic pelvic pain.²⁵
- (d) Irritable bowel syndrome: characterized by a complex of symptoms associated with abnormal gastrointestinal motility. Symptoms of IBS may be seen in 50% to 80% of patients with chronic pelvic pain.^{26,27}
- (e) Neoplasia: Carcinoma colon or rectum should be included in differential diagnosis of chronic pelvic pain.¹¹
- (iii) MUSCULOSKELETAL CAUSES:
 - (a) Coccydynia a fall or any trauma to coccyx can cause coccygeal pain.²²
 - (b) Levator ani syndrome can be diagnosed when a patient has chronic pelvic pain in absence of another explanation of pain caused by spastic pelvic floor muscle.^{11,22}
 - (c) Myofascial pain syndrome it is a disorder of muscle characterized by the presence of trigger points that when stimulated gives rise to localized pain.²²
 - (d) Prolapsed intravertebral disc¹¹
 - (e) Tumors of pelvic bones and muscles¹¹
 - (iv) Psychogenic^{11,28}
 - (v) Idiopathic^{2,11}

Material and method

This prospective study was conducted in Lalla Ded Hospital; an associated hospital of Government Medical College Srinagar Jammu & Kashmir. The duration of study was from March 2011-September 2012 with sample size of 100 patients. Aim and objectives: To evaluate various causes of chronic pelvic pain. To find out the correlation between clinical and laparoscopic examination. To compare the efficacy of clinical examination and diagnostic laparoscopy and to make a medical or surgical intervention as well if possible.

Inclusion Criteria

Women in age group of 20-50 years.
Women presenting with history of pelvic pain of 6 months or more.

Exclusion Criteria

Women with < 20 years or > 50 years of age.
Women with non-gynecological etiology.
The following data were recorded.
General characteristics of patients.
Demographic data

Detailed history was taken including onset, progression, location, characteristics of pain, duration of pain, radiation of pain, factors aggravating and relieving pain, association of pain with posture relation, menstrual periods and dyspareunia. Associated symptoms like nausea, vomiting, constipation, decreased appetite, fatigue and fever will be noted. History of previous surgery and infections was recorded. Obstetrical history and menstrual history was also recorded.

A thorough physical examination was done including: General physical examination, vitals, and systemic examination. Laboratory measures such as complete haemogram, KFT and LFT, urine examination, stool examination, ECG all leads, Chest X-ray and ultrasonography of pelvic and abdominal organ was done and was recorded.

After history taking, clinical examination and investigations, patients with obvious non-gynecological etiology was excluded. The remaining patients were investigated by performing diagnostic laparoscopy to determine type of pathology existing and if deemed necessary surgical or medical treatment were done. Diagnostic laparoscopic examination was performed under general anesthesia, pelvic and abdominal organ was visualized and pathology in any organ was noted.

The result was presented as descriptive statistics, Kappa statistics³¹

was used to estimate the extent of correlation between clinical examination and diagnostic laparoscopic findings.

We plan this study to detect the potentially treatable pathology by use of laparoscopy that was not detailed and detectable by other types of evaluation such as clinical examination. While subjecting patient to diagnostic laparoscopy various interventions were done where ever possible, i.e. adhesiolysis, drainage of chocolate cyst, cystectomy, drilling of polycystic ovaries, and aspiration of fluid in POD. In patients of infertility while subjected to diagnostic laparoscopy, chromotubation was done to assess the patency of tubes, and their further management was planned accordingly.

DISCUSSION

In the study, majority of the patients were in the age group of 31-40 years (63%), followed by 20-30 years (24%) and 13% of patients were in the age group of 41-50 years. The age wise distribution of patients in our study were in congruence with the study of Kamilya Gourisankar et al¹ where 54% patients belonged to the age group of 31-40 years, 32% patients belonged to the age group of 21-30 years and 14% to the age group of above 40 years. The mean duration of pain in the studied subjects was 30.3 months with a standard deviation of 25.6 months (range from 6 months to 10 years).

In our study, duration of pain significantly increased with the age of patients ($p < 0.002$) from the mean of 19 months with the standard deviation of 2.9 months (range 6 months – 5 years) in 20-30 years to 39.5 months with standard deviation of 8.67 months (range 6 months – 8 years) in 41-50 years. There was statistically significant association between duration of pain with age of patients. This is similar to the results of Zonderva KT et al³² study where duration of symptoms increases with age ($p < 0.001$) from the mean of 13.7 months in 13-20 years old to 20.2 months in age over 60 years, as symptoms persist longer in the older women. Similar results were found in Kamilya Gourisankar et al¹ study where duration of symptoms increased with age.

The commonest associated symptom was dysmenorrhea (50%) in the enrolled patients. Our results are in conformity with the study of Shripad Hebbar et al³², Kamilya Gourisankar et al¹, and Zonderva KT et al³¹, where 55.3%, 56% and 45% had dysmenorrhea respectively. In the enrolled patients, other complaints were menorrhagia 24% and infertility 11%. Our observations were in conformity with the observations of the study by Kamilya Gonasankar et al¹ On performing clinical examination, 42% of patients showed normal pelvic findings, our observations were in conformity with the observation in the studies of Shripad Hebbar et al³², Vercellini et al³⁴, HK Ho et al²² and Kamilya Gourisankar et al¹ where 42% , 40.4% , 44% and 47% of patients had normal clinical examination respectively. Similar observations were found in studies of HK Ho et al²² and Kamilya Gourisankar et al¹ where 44% and 47% of patients had normal clinical findings. In the study group, 58% of enrolled patients had abnormal clinical finding, which was in congruence with the study of Kamilya Gourisankar et al¹, where 53% of patients had abnormal clinical examination.

In the studied group, on performing clinical examination on enrolled patients, 22% had enlarged uterus (out of this 6 had fibroid uterus), 15% had retroverted uterus, 14% had restricted mobility of uterus, followed by 25% who had mass felt through fornices, 18% had thickened fornices, 6% mass in pouch of Douglas and 3% had thickened pouch of Douglas. Our observations were similar to the study conducted by Kamilya Gourisankar et al¹, where 22% had enlarged uterus, 25% had retroverted and restricted mobility of uterus, 33% had mass felt through fornices, 20% had thickened fornices and 9% had nodules pouch of Douglas.

The enrolled patients when subjected to diagnostic laparoscopy, 20% had no visible pathology. Our observations are in conformity with the observations of the studies conducted by Murphy and Flienger et al³⁵, Jacobson and Westron et al³⁶, Fear et al³⁷ and Goldstein et al³⁸ where 20%, 23%, 26% and 14% had normal diagnostic laparoscopy respectively.

On performing diagnostic laparoscopy on the study group, 80%

had abnormal laparoscopic findings. Our observations are in conformity with the observations reported in studies of Shalaya Chetteri et al¹¹, Mara et al²⁹, Goldstein et al⁴⁰ and Chatmen and Ward et al⁴¹, where 80%, 82%, 86% and 88% had visible pathology on laparoscopic examination respectively. The commonest laparoscopic diagnostic pathology in the enrolled patients was chronic PID 36% that is in conformity with the study of Kamilya Gourisanker et al¹ where 36% had chronic PID.. In the study group pelvic adhesions were second most common abnormality on diagnostic laparoscopy 32%, similar to the study of Drozyyik et al¹⁴ and Kamilya Gourisanker et al¹ where 32.5% and 30% had adhesions on laparoscopic examination respectively. In the study group, endometriosis was found in 47% of patients by laparoscopic examination (out of which 19 had chocolate cysts), that is in conformity with the study of Carter et al⁴², Newham et al and Vereline et al⁸ where endometriosis was diagnosed in 48%, 40% and 40.4% on diagnostic laparoscopy respectively. In the enrolled patients, pelvic congestion (diagnosed by diagnostic laparoscopy as a generalized hyperemia and large dilated pelvic vessels on broad ligament or infundibulopelvic ligament), was diagnosed in (17%) of patients on diagnostic laparoscopy. That was in conformity with the study of Kamilya Gourisanker et al¹ and Shrippad Hebbar et al³⁴ where 13% and 18.6% had pelvic congestion respectively.

The study showed that, on comparing the diagnostic laparoscopy with the clinical examination, diagnostic laparoscopy was more predictive in diagnosing PID 36%, pelvic adhesion 22% pelvic congestion, 17% and endometriosis in 47%. Similar observations were reported in the study of Kamilya Gourisanker et al¹.

Our study showed that on comparing the laparoscopic examination with the clinical examination. The sensitivity of clinical examination was 72.5%, this was in conformity with the study of Kamilya Gourisanker et al¹ were the sensitivity of clinical examination was 71.6%. The negative predictive value of clinical examination was 47.6%, that was in conformity with study of Kamilya Gourisanker et al¹ and Ozaksit et al²⁸ who had reported negative predictive value of 55.3% and 42.8% respectively. On comparing diagnostic laparoscopy with clinical examination, the accuracy was 71.6%.

Our study showed that by applying Kappa statistics³² there is fairly good correlation between number of normal and abnormal cases diagnosed by laparoscopy verses clinical examination Kappa= 0.51. This study showed that laparoscopic examination is an excellent tool in evaluation of patients with pelvic pain. There was a better correlation between an abnormal clinical examination and abnormal laparoscopic examination 72.5% that is in congruence with the study of Fear³⁸, Jacobsen, Kamilya Gourisanker et al¹, and Weston et al³⁷ and Lundberg et al⁴⁵ who had observed 74%, 77% and 59% correlation respectively.

CONCLUSION:

Laparoscopy is gold standard for diagnosis of chronic pelvic pain. Good clinical examination is informative but it is not conclusive in many patients. But there arise a need for imaging of pelvic organs by ultrasonography. Hence ultrasonography will be informative where clinical examination is inconclusive and laparoscopic facilities are not available.

RESULT AND OBSERVATION

The results and observations obtained are depicted in the following tables and graphs.

Table – 1: Depicting the age distribution of enrolled patients

Age (Years)	N	%
20 to 30	24.0	24.0
31 to 40	63.0	63.0
41 to 50	13.0	13.0

Majority of patients belonged to the age group of 31-40 years (63%) followed by 24% to the age of 24-30 years and 13% were in the age group of 31-50 years old.

Table – 2: Depicting duration of pain lower abdomen (month) in the study group.

Duration of Pain (months)	N	%
≤ 12	32	32.0
13 to 24	16	16.0
25 to 36	19	19.0
37 to 48	14	14.0
> 48	19	19.0
Mean±SD	30.3±25.6 months (6, 120)	

The mean duration of pain is 30.3 months with a standard deviation of 25.6 months (range 6 months to 10 years).

Table – 3 :Depicting relation of duration of pain with age of study group

Age (month)	Mean ± SE	p value
20 to 30	19.0 ±2.9 (7,60)	< 0.002 (Sig)
31 to 40	36.9 ± 3.3 (7, 120)	
41 to 50	46.4 ± 8.1 (7, 108)	

Pain significantly increase with the age of patients (p < 0.001) from the mean of 19 months with the SD of 2.9 months (range 6 months – 5 years) in 20-30 years to 39.5 months with SD of 8.67 months (range 6 months – 8 years) in 41-50 years.

Table – 4 :Depicting Associated Symptoms in enrolled patients

Associated Symptoms	N	%
Dysmenorrhea	50	50.0
Menorrhagia	24	24.0
Infertility	10	10.0
Surgical Intervention	8	8.0

Shows that dysmenorrhea was commonest associated symptom in 50% followed by menorrhagia in 24%, infertility in 10% and surgical intervention in 8%.

Table – 5: Depicting per speculum examination findings in the enrolled patients

Per Speculum examination	N	%
Unhealthy Discharge	8	8.0
Hypertrophy	6	6.0
Congested	4	4.0
Excessive Discharge	2	2.0

Majority of patients had unhealthy discharge (8%), followed by hypertrophied cervix (6%), congested cervix (4%) and excessive discharge (2%).

Table – 6: Depicting per vaginal examination findings of study group

Structure	Abnormality	n	%
Uterus	Enlarged	22	22.0
	Retroverted	15	15.0
	Restricted Mobility	14	14.0
Fornices	Thickened	18	18.0
	Mass	25	25.0
POD	Mass	6	6.0
	Thickened	3	3.0

As depicted by table, 25% of patients had mass felt in fornices, 22% had enlarged uterus, 18% had thickened fornices, 15% had retroverted uterus with restricted mobility of uterus in 14%, 6% had mass in POD and 3% had thickened POD.

Table – 7: Depicting ultrasonographic findings in the study group

Structure	Abnormality	N	%
Uterus	Enlarged	41	41.0
Ovarian Mass	Tubo-Ovarian Mass	17	17.0
	Cyst	16	16.0

Fallopian Tubes	Hydrosalpinx	7	7.0
POD	Mass	8	8.0
	Fluid	3	3.0

Majority of patients with enlarged uterus 41%, 17% had tubo-ovarian mass, 16% ovarian cyst, 7% had hydrosalpinx, 8% had mass seen in POD and 3% had significant fluid in POD.

Table – 8: Depicting laparoscopic abnormalities in the study group

Structure	Abnormality	n	%
Uterus	Enlarged	13	13.0
	Endometriosis	3	3.0
	Adhesions	5	5.0
	Fibroid	6	6.0
Ovaries	Functional Cyst	8	8.0
	Endrometroma	19	19.0
	Tubo-Ovarian Mass	22	22.0
	Polycystic Ovary	6	6.0
	Adhesions	10	10.0
Fallopian Tubes	Tortuous	17	17.0
	Adhesions	13	13.0
	Hydrosalpinx	8	8.0
POD	Endometriosis	21	21.0
	Adhesions	32	32.0
	Fluid	22	22.0
	Congestion	17	17.0

Majority of the patients had adhesions 32%, 22% had fluid in POD and tubo-ovarian mass, 21% had endometriosis followed by 17% with pelvic congestion, 13% had hydrosalpinx and 13% had enlarged uterus out of which 6% had fibroid uterus.

Table 9: Comparing Laparoscopic examination in relation with clinical examination in the study group

	Laparoscopy	Clinical Findings	
		Abnormal	Normal
Abnormal	80	58 (72.5)	22 (27.5)
Normal	20	0	20

The table shows that 80% of women had abnormal findings on laparoscopic examination, 58% of women had abnormal pelvic examination.

Table 10: Depicting laparoscopic examination in relation to clinical findings in the study group.

	Sensitivity	Specificity	PPV	NPV	Accuracy
Clinical Findings	72.5	100	100	47.6	78

Table shows that clinical examination had sensitivity of 72%, negative predictive value of 47.6% with accuracy of 78%.

Table11: Showing Comparison of Laparoscopic and Clinical Examination in the Enrolled Patients

	Clinical	Laparoscopic
PID	22	36
Myoma	5	16
Cyst	7	15
Endometriosis	6	29
Congestion	0	15
Adhesions	0	27


As shown in table that diagnostic laparoscopy is more predictive in diagnosing PID36%, endometriosis 29%, adhesions 27% and pelvic congestion in 15%.

Table 12: Depicting correlation between laparoscopic and clinical examination findings in the study group.

Laparoscopic Findings				
Clinical Findings	Abnormal		Normal	
	n	%	n	%
Abnormal	58	72.5	0	0.0
Normal	42	27.5	13	100.0

As shown in table there is better correlation between clinical and laparoscopic examination of 72.5%. Thus there is better correlation between laparoscopic examination and clinical examination

Receiver Operating Characteristic Curve Showing Comparison of Laparoscopic Examination with Ultrasonographic and Clinical Findings in the study group.



REFERENCES

- Kamela Gaurisankar, Mukherji Joydev, Gayen, Abhimanyu. Different methods of evaluation of chronic pelvic pain. J Obstet Gynecol 2005; 55(3): 251-253.
- Alexandra Simopoulou, Sohel Siddique. Chronic pelvic pain. In: The John Hopkins Manual of Gynaecology and Obstetrics Ed. 3rd Editors Kimberly B. Fartner, Linda M. Szy Maniskie, Harold E Fox, Edward E. Wallach. Page No. 312.
- Kontoravdis A, Hassan E, Hassiakos D et al. Laparoscopic evaluation and management of chronic pelvic pain during adolescence. Clin Exp Obstet Gynecol 1999; 26: 76-7.
- Rapkin AJ. Adhesion and pelvic pain: a retrospective study. Obstet Gynecol 1986; 68:13.
- Howard FM. The role of laparoscopy as a diagnostic tool in chronic pelvic pain. Bailcers Best Pract Res Clin Obstet Gynecol 2000; 14: 467-92.
- Howard FM. Chronic pelvic pain. Obstet Gynecol 2003; 101(3): 594-611.
- Kresch AL, Seifes DB, Sachs LB. Laparoscopy in 100 women with chronic pelvic pain. Obstet Gynecol 1984; 64: 72.
- Kontoravdis A, Chrysiopoulos A, Hassiakos D et al. The diagnostic value of laparoscopy in 2365 patients with acute and chronic pelvic pain. Int J Gynaecol Obstet 1996; 52: 243-8.
- Lawrence MD. Effect on negative laparoscopy rate in chronic pelvic pain using patients assisted laparoscopy. JSJS 1997; 1(4): 319-321.
- Cunanan RG, Courey WG, Lippes J. Laparoscopic findings in patients with chronic pelvic pain. Am J Obstet Gynecol 1983; 136: 589-591.
- Shallaya Chhetre, S Khanna, A Pooman, B Sen. Laparoscopy as a diagnostic tool in evaluation of chronic pelvic pain in women. J of Nepal Health Research 2009; 7: 45-48.
- L S F LO. Management of chronic pelvic pain, a gynecologist's perspective. The Hong Practitioner 1999; 21: 564-571.
- Nikkanen V, Punnonen R. Clinical significance of adenomyosis. Am Chir Gynecol 1980; 69: 278-80.
- Drozgyik I, Vizer M, Szabo I. Significance of laparoscopy in the management of chronic pelvic pain, Eur J Obstet Gynecol Reprod Biol 2007; 133(2): 223-26.
- Ness RB, Soper DE, Holley RL et al. Effectiveness of inpatients and outpatients treatment strategies for women with PID: results from PID evaluation and clinical health (PEACH) randomised trial. Am J Obstet Gynecol 2002; 186: 929-37.
- David D. Ortiz, MD, Christus. Chronic pelvic pain in women. Am Fam Physician 2008; 77: 1535-1542.
- Rawson RL. Prevalence of endometriosis in asymptomatic women. J Reprod Med 1991; 36: 513-55.
- Sernn K. Laparoscopy in gynecology. Geburtshilfe Frauenmheilkd. 1967; 27(11): 1029-42.
- Abu Rafeh B, Vilos GA, Mishra M. Frequency and laparoscopic management of ovarian remnant syndrome. Am Assoc Laparosc 2003; 10: 37-47.
- Dekel A, Efrat Z, Orieto R et al. The residual ovarian syndrome: a 20 years experience. Eur L Obstet Gynecol 1996; 68: 159-64.
- American Fertility Society. The American Fertility Society classification of adnexal adhesions, distal tubal occlusion, tubal occlusion secondary to ligation, tubal pregnancies, Mullerian anomalies and intrauterine adhesion. Fertil Steril 1988; 49: 944-55.
- H K HO, L N Sim, L K HO. The diagnostic value of laparoscopy in women with pelvic pain. Sing Med 1989; 30: 453-456.
- Steege S, Dodson JF, Hughes WC, Claude L. Relationship of laparoscopic findings to self report of pelvic pain. Am J Obstet Gynecol 1991; 2: 93-78.
- Vincent K. Chronic pelvic pain in women. Post Grad Med J 2009; 85: 24-29.
- Whitmore K, Seegel JK, Killog SP. Incidence of painfull bladder syndrome as a cause of sexual pain in women. J Sex Med 2007; 4: 720-7.
- Standford EJ, Koziol J, Fing A. The prevalence of interstitial cystitis endometriosis, adhesion and vulval pain in women with chronic pelvic pain. 2005; 12:43-9.
- Nikkanen V, Punnonen R. Clinical significance of adenomyosis. Am Chir Gynecol 1980; 69: 278-80.
- Richter HE, Holley RL, Chandraiah S, Varner RE. Laparoscopic and psychological evaluation of women with chronic pelvic pain. Int J Psychiatr Med 1998; 28: 243-53.
- Riggs, Rivan M, M Carthy, Joen M et al. The utility of the Beck Depression Momentary in the evaluation of chronic pelvic pain. 2008; 63: 148-149.
- Nezhat FR, Crystal RA, Nezhat CH et al. Laparoscopic adhesiolysis and relief of chronic pelvic pain. JSLS 2000; 4:281-5.
- Pereera Maxell F. A to Z of medical statistics. A comparison for critical appraisal London Arnold 1998; 99-100.

32. Zonderva KT, Yudin PL, Vessey MP et al. Patterns of diagnosis and referral in women consulting for chronic pelvic pain in United Kingdom primary care. *Br J Obstet Gynecol* 1999; 99:99-100.
33. Shripad H, Chander C. Role of laparoscopy in evaluation of chronic pelvic pain. *Min Access Surg* 2008; 1: 116-20.
34. Vercellini P, Fedele L, Arcaini L et al. Laparoscopy in the diagnosis of chronic pelvic pain in adolescent women. *J Reprod Med* 1989; 34: 827-30.
35. Murphy A, Flinger J. Diagnostic laparoscopy: Role in management of pelvic pain. *Med J Aust* 1981; 1: 571.
36. Jacobsen L, Westrom L. Objectivised diagnosis of acute pelvic inflammatory disease. *Am J Obstet Gynecol* 1969; 105: 1088-98.
37. Fear RE. Laparoscopy. A valuable aid in gynecologic diagnosis. *Obstet Gynecol* 1968; 31: 297-309.
38. Goldstein DP. New insights into the old problem of chronic pelvic pain. *Pediatr Surg* 1979; 14: 675.
39. Manish K. Singh MD, Elizabeth E Puscheck MD, Jaswant Patil. Chronic pelvic pain. *eMedicine Obstet Gynecol* 2010.
40. Goldstein DP, Emans SJ, Levanthal JM. Laparoscopy in the diagnosis and management of pelvic pain in adolescents. *J Reprod Med* 1980; 24: 251.
41. Chatman DL, Ward AB. Endometriosis in adolescents. *Reprod Med* 1982; 27: 156
42. Carter JE. Combined hysteroscopic and laparoscopic finding in chronic pelvic pain. *J Am Assoc Gyneacol Laparosc* 1994; 2: 43-7.
43. Lunberg WI, Wall JE, Malairs JR. Laparoscopy in evaluation of pelvic pain. *Obstet Gynecol* 1973; 42: 872-6.