



## ORIGINAL RESEARCH PAPER

Pathology

## EFFICACY OF ENTECAVIR IN HBeAg POSITIVE TREATMENT NAIVE CHRONIC HEPATITIS –B PATIENTS

**KEY WORDS:** Chronic hepatitis B, Entecavir, (ETV) HBV DNA.**Manjushree Nayak**

Associate Professor, Pathology, SCB Medical College, Cuttack

**Rajendra Kumar Panda**

Assistant Professor, Pharmacology, SCB Medical College, Cuttack

**Umesh Chandra Patra**

Professor &amp; Head, Department of Hepatology, SCB Medical College &amp; Hospital Cuttack, Odisha, India -7530007.

## ABSTRACT

**Background:** Entecavir is a nucleoside analogue drug that has selective anti-hepatitis B virus (HBV) activity. It is indicated for the treatment of chronic hepatitis B virus infection in adults with evidence of active viral replication, persistent elevations in serum aminotransferase and histologically active disease. The present study was conducted to evaluate the efficacy of entecavir (ETV) in hepatitis B e antigen (HBeAg) positive- chronic hepatitis B (CHB) treatment naive Indian patients in real life scenario.

**Materials & Methods:** This is a single center open label prospective observational study to evaluate the efficacy of Entecavir in chronic hepatitis B patients. A total of 140 patients were enrolled in the study from January 2015 to December 2016. Parameters were evaluated at baseline and at 12, 24 and 40 weeks of the treatment, Efficacy of Entecavir was assessed by evaluating hepatitis B virus DNA (HBV DNA), serum alanine aminotransferase (ALT) and sero-conversion.

**Result:** The mean HBV DNA at baseline was 5.99 log [on a base-10 scale] IU/mL which decreased to 2.12 log IU/mL at the end of 40 weeks. Thus there was a mean change of 3.87 log IU/ml which was statistically significant ( $P < 0.0001$ ). Out of 140 HBeAg positive subjects, 50 (35.71 %) had become negative at the end of 40 weeks. Only 10.71% of the patients had normal ALT ( $< 40$  IU/l) values in the beginning which increased to 100% at the end of 40 weeks.

**Conclusion:** Entecavir significantly improves virological, biochemical and serological markers in HBeAg positive treatment naïve chronic hepatitis B patients.

## INTRODUCTION :

Chronic hepatitis B affects an estimated 400 million people worldwide and causes more than 5 million deaths due to complications of chronic infection<sup>1</sup>. Nearly 40 million people out of the global HBV infection pool are from India and every year over 100,000 Indians die due to complications associated with chronic infection<sup>2</sup>.

Chronically infected patients with prolonged elevated HBV DNA level, elevated ALT level, and presence of HBeAg are at increased risk of developing progressive liver disease, cirrhosis, hepatic decompensation, hepatocellular carcinoma (HCC) and death<sup>1</sup>. HBV DNA level of more than 2000 IU/ml ( $10^4$  copies/ml) is a strong precursor for development of complications like cirrhosis and HCC<sup>3</sup>. Patients with chronic HBV infection who have ALT levels that are near the upper limit of the normal range are at a significantly higher risk for complications of cirrhosis and HCC than patients with ALT levels that are less than half the upper limit of the normal range. The highest risk of complication of cirrhosis and HCC occurs in patients with ALT levels that are one to two times the upper limit of the normal range<sup>4</sup>.

Seropositivity for HBeAg and/or HBV-DNA  $> 2,000$  IU/ml are significant risk factors for cirrhosis and HCC development, even in asymptomatic subjects with chronic HBV infection<sup>5</sup>. The goal of treatment is to suppress HBV replication and ensure the loss of HBeAg with ALT normalization at the end of the treatment, thus decreasing progression of the liver disease to cirrhosis and HCC. Drugs approved for HBV treatment include interferon, nucleoside or nucleotide analogs (lamivudine, adefovir, entecavir, tenofovir, and telbivudine)<sup>6</sup>.

Interferon requires parental administration, and causes many side effects especially in cases of cirrhosis<sup>7</sup>. A substantial percentage of patients particularly those with high levels of viral replication did not respond to treatment with interferon alfa alone or after a short course of corticosteroids<sup>8</sup>. Nucleoside and nucleotide analogs are administered orally, they cause more profound HBV DNA suppression. However these drugs are associated with rebound increase of HBV DNA levels or reactivation of hepatitis if discontinued prematurely. In addition, long-term use of these

drugs is compromised by the development of resistance<sup>6</sup>.

ETV is a potent and selective inhibitor of HBV DNA polymerase<sup>9</sup>. In preclinical studies performed in chronic woodchuck hepatitis virus infections, ETV showed potent and sustained suppression of viral DNA and an absence of both viral rebounds and emerging ETV resistance (ETVr<sup>10</sup>). ETV reduced the covalently closed circular DNA viral reservoir to undetectable levels and extended the lives of treated woodchucks by preventing HCC<sup>10</sup>. Therapy with ETV in HBeAg-positive CHB patients demonstrated superior histologic, virologic, and biochemical responses compared with lamivudine at 48 weeks<sup>1</sup>.

In a study by Leung et al, comprehensive monitoring of genotypic and phenotypic antiviral resistance was performed on 673 ETV treated nucleoside naive HBV patients; only 3% of ETV treated patients exhibited virologic rebound by the end of 96<sup>th</sup> week<sup>11</sup>. Therefore we conducted a prospective observational study to evaluate the efficacy of entecavir in seropositive chronic hepatitis B treatment naive Indian population in real life scenario.

## MATERIALS &amp; METHOD

**Study design:** This is an open label prospective observational study to evaluate the antiviral efficacy of ETV in treatment naive chronic hepatitis B patient. The study was conducted from January 2015 to December 2016 in the department of Hepatology, Pathology and Pharmacology at SCB Medical College & Hospital, Cuttack, Odisha after approval from Institutional Ethical Committee. Data were collected from patients with HBeAg positive chronic hepatitis B. Written informed consent was taken after explaining the details of the study protocol.

## Patient selection:

Patients were enrolled as per the following inclusion & exclusion criteria. HBeAg positive chronic hepatitis B men and women older than 16 years of age, patients with compensated liver disease document by elevated serum ALT levels, patients having HBV DNA levels greater than 2000 IU/ml, nucleoside naive patients, compliant patients were included in the study. Women of childbearing potential willing to use an acceptable method of

contraception to avoid pregnancy during treatment and for 8 weeks after completion of the study were also included in the study.

HBeAg positive CHB patients diagnosed or suspected hepato cellular carcinoma (HCC), associated diseases including heart disease, immunologic disease, malignant tumors, hypersensitive to nucleoside or nucleoside (acid) analogues or with a history of nucleoside antiviral drug treatment, evidence of decompensated liver disease and history of drug abuse or alcohol abuse were excluded from the study. Pregnant or lactating women not using adequate contraceptive measures were excluded from the study

**Intervention**

Patients received medication as a part of their usual treatment. They were not exposed to any experimental investigations. After enrollment in the study, all patients received 0.5mg of ETV orally on empty stomach once daily for a period of 40 weeks. The data for 140 patients were analyzed to evaluate the efficacy of ETV. The study comprised of 4 visits, viz. baseline visit and visits after 12, 24, and 40 weeks of treatment. Of the 140 patients enrolled in the study all completed the study and were considered for evaluating efficacy data.

**Data Collection**

Data collection was done at baseline (visit 1) and at visits 2, 3 and 4 at initiation i.e. at 12, 24, 40 weeks of treatment. The data collected at baseline were demographics, laboratory investigation, comorbid conditions, concomitant medications and drug allergies. At subsequent visits data useful to evaluate the efficacy of ETV i.e. HBV DNA load, serum ALT and HBeAg were collected.

**Assay Methodology**

Serum HBV DNA was quantified using the taqMan Real-Time polymerase chain reaction. The lower limit of detection was (<20 IU/ml). Serum ALT level was quantified by local laboratory, normal value range 0-40IU/L.

**Efficacy Analysis**

The efficacy of antiviral therapy of chronic hepatitis B is measured using surrogate markers. These include undetectable levels of HBV DNA, normalization of ALT, and HBeAg seroconversion at the end of 40 weeks.

**Statistical methods & analysis**

The software package used for statistical analysis was SAS software package version 9.1.3. Paired t-test was used for finding changes from any two visits when the data followed normality assumption and Wilcoxon signed rank test was used when data did not follow normality assumption. McNemar's test was used for paired categorical data for finding change between visits.

**OBSERVATION**


**Demographic summary:** Demographics for all the subjects are shown in **Table 1**. The mean age of subjects was 37.6 years, with 22 years as minimum and 65 years as maximum. There were more male subjects (N=100, 71.43%) than female subjects (N=40, 28.57%). The mean (SD) weight was 60.1 kgs, with a minimum of 46 kgs and maximum of 75 kgs. All patients were HBeAg positive.

**Table 1: Demographic and disease characteristics at the base line**


Demography	
Number of Patients	N=140
Male/female (n %)	(71.43%/28.57%)
Age (years)	37.6±10.86
Weight (Kgs)	60.5±8.43
DISEASE CHARACTERISTICS AT BASELINE	
Mean HBVDNA	Log 5.99± 1.116 IU/ml
Mean Serum ALT	69.7 ±25.88 IU/L
HBeAg positive	N=140 (100%)

**Efficacy analysis :  
HBV DNA (Viral Load):**

**Fig 1** depicts the transition in HBV DNA Levels in our patients across the study period. There are no cases with undetectable Levels of HBV DNA at visit one which improved to 3.57% in visit 2, 32.14 % in visit 3 and 64.29% in visit 4. After considering the logarithmic transformation of the viral load, the mean HBV DNA (log) was gradually decreased from visit 1 (5.986 log IU/ml), to visit 4 (2.120 log IU/ml) and this is statistically significant. P<0.05. It is graphically represented in **Fig-2**



**Figure 1**



**Figure 2  
Serum ALT levels:**

The mean of ALT in visit 1 was 69.7 IU/L ± 25.88 IU/L. It decreased to 41.1 IU/L ± 17.98 IU/L in visit2, 27.1 IU/L ±5.44 IU/L in visit3 and 24.8 IU/L ± 2.39 IU/L in visit 4.

Thus, there was a mean decrease of 44.90 IU/L ± 25.70 IU/L from the baseline to the end of the treatment at 40 weeks which was statistically significant (P<0.0001). It is observed that the normal ALT level (40 IU/ml) was 10.71% in first visit, 60.71% in second visit, 96.43% in third visit and 100% in fourth visit.

**Seroconversion of HBeAg -** Out of 140 HBeAg positive subjects in visit 1, 50 (35.71%) had become negative in fourth visit. **[Table 2.]**

**Table -2 HBeAg seroconversion**

	Baseline	40 weeks
HBeAg positive	(100.0%)	90 (64.29%)
HBeAg negative	00 (00.0%)	50 (35.71%)

**DISCUSSION:** Although the efficacy of ETV to suppress the HBV DNA load have been demonstrated in various clinical trials there is limited data on its use in real clinical practice in India. This prospective, observational study assessed the efficacy of ETV in the routine clinical management of treatment naive CHB patients. Our study demonstrates that ETV is associated with significantly higher rates of serologic, viral, and biochemical improvement. Undetectable levels of HBV DNA by PCR assay were noted in 64.29% of patients by 40 weeks of treatment. Starting from the beginning of the study to its completion at 40 weeks ALT normalization was noted in 100% and HBeAg Seroconversion in 35.71% patients. Our results are in agreement with findings of Chang et al<sup>1</sup> who showed in their study that a significantly greater number of patients on ETV achieved undetectable levels of HBV DNA than patients on lamivudine when monitored with PCR assay (67% vs.36%, p<0.001) and ALT levels (68% vs.60%).

**LIMITATIONS OF THE STUDY :**

The limitation of our study were that safety parameters were not

taken into consideration ,data of the patients after 40 weeks of treatment was not available therefore further studies are recommended to evaluate rebound increase in viral DNA load, development of ETV resistance and long term safety of the drug.

#### CONCLUSION:

Our study cohort was small and patients had no other co-morbid conditions or any concurrent drug therapy. Thus further studies evaluating the efficacy in such complex settings. We conclude that entecavir significantly improves viral biochemical and serological markers of HBeAg positive chronic hepatitis treatment naive patients who have not received nucleoside analogs for their condition.

#### CONFLICT OF INTEREST :NIL

#### REFERENCES

1. Gish RG, Lok AS, Chang TT, de Man RA, Gadano A, Sollano J, et al Entecavir therapy for up to 96 weeks in patients with HBeAg-positive chronic hepatitis B. *Gastroenterology*. 2007 Nov;133(5):1437-44.
2. Choudary A, Santra A, Chakravorty A, Banarji A, Pal S, Dhali G K, et al Community-based epidemiology of hepatitis B virus infection in West Bengal, India: Prevalence of hepatitis B e antigen-negative infection and associated viral variants *Journal of Gastroenterology and Hepatology* November 2005.
3. Lai CL, Yuen MF Chronic hepatitis B- New goals, new treatment. *N Engl J Med*. 2008 Dec 4;359(23):2488-91.
4. Yang MF, Yuan HJ, Wong DK, Yuen JC, WM, Chan AO, et al. Prognostic determinants for chronic hepatitis B in Asian: therapeutic implications. *Gut*. 2005;54:1610-4.
5. Yang HI, Lu SN, Liaw YF, You SL, Sun CA, Wang LY, et al Taiwan community –based cancer screening project group. Hepatitis B e antigen and the risk of hepatocellular carcinoma, *N Engl J Med*. 2002;18;347(3):168-74.
6. Perrillo RP, Marcellin P Effect of newer oral antiviral agents on future therapy of chronic hepatitis B, *Antivir Ther*. 2010;15(1):13-22.
7. De Man RA, Wolters LM, Nevens F, Chua D, Sherman M, Lai CL, et al Safety and efficacy of oral entecavir given for 28 days in patients with chronic hepatitis B virus infection *Hepatology*. 2001 Sep;34(3):578-82.
8. Perrillo RP, Schiff ER, Davis GL, Bodenheimer HC Jr, Lindsay K, Payne J, Dienstag JL, O'Brien C, Tamburro C, Jacobson IM, et al A randomized, controlled trial of interferon alfa-2b alone and after prednisone withdrawal for the treatment of chronic hepatitis B. The Hepatitis Interventional Therapy Group *N Engl J Med*. 1990 Aug 2;323 (5): 295 -301 .
9. Yao G, Chen C, Lu W, Ren H, Tan D, Wang Y, Xu D, Jiang Z, Liu J, Xu D, macdonald L; for the A1463023 Study Group Efficacy and safety of entecavir compared to lamivudine in nucleoside –naïve patients with chronic hepatitis B: a randomized double-blind trial in China *Hepato Int*. 2008Mar;2(1):136.
10. Colonna RJ, Rose R, Baldick CJ, Levine S, Pokornowski k, Yu CF, Walsh Entecavir resistance is rare in nucleoside naïve patients with hepatitis b *Hepatology*. 2006 Dec;44(6):1656-65.
11. Leung N, Peng CY, Hann HW, Sollano J, Lao-Tan J, Hsu CW, et al Early Hepatitis B virus DNA Reduction in Hepatitis B e Antigen-positive Patients with Chronic Hepatitis B: A Randomized International Study of Entecavir Versus Adefovir *Hepatology*. 2009 Jan;49(1):72-9.