



ORIGINAL RESEARCH PAPER

Radiology

ROLE OF MRI (MAGNETIC RESONANCE IMAGING) BRAIN IN EPILEPSY

KEY WORDS: AUB secretary endometrium POC

Dr. Sahil N. Shah 2nd year Resident, Department of Radiology, VSGH

Dr. Dharita S. Shah Prof. and HOD, Department of Radiology, VSGH

Dr. Darshini R. Patel 2nd year Resident, Department of Radiology, VSGH

Dr. Hardik G. Rabari 2nd year Resident, Department of Radiology, SCL.

ABSTRACT

The main purposes of neuroimaging in epilepsy patients are to identify underlying structural or metabolic abnormalities that require specific treatment and to aid in formulating a syndromic or etiological diagnosis. Many imaging modalities for investigating seizures are available like neurosonogram, computerized tomography (CT scan) of brain, magnetic resonance imaging (MRI) of brain, functional MRI, positron emission tomography (PET), single photon emission computed tomography (SPECT). Of these various modalities MRI is most important modality for evaluation of structural disorders causing epilepsy. So, role of MRI to be evaluated in patients with epilepsy.

AIMS AND OBJECTIVES:

- To evaluate role of MRI in differentiating primary and secondary epilepsy.
- Investigate the incidence of structural abnormalities in epilepsy.
- To identify different structural (except neoplasm) and metabolic causes in patient with secondary epilepsy.
- Classify causes of epilepsy according to age of patients & specific seizure type.
- Systematically evaluate role of MRI epilepsy protocol in epileptic patients.
- To evaluate sensitivity and specificity of hippocampal pathology with epilepsy protocol.
- Establish role of MRI in management and follow-up of epilepsy.

INTRODUCTION

Epilepsy is one of the **common neurological disorder** world wide. At least fifty million people in the world suffer from recurrent non provoked seizures. Incidence of epilepsy is clearly higher in developing countries than in developed countries.

A seizure is defined as a paroxysmal alteration in neurologic function due to excessive electrical discharge from the central nervous system. Epilepsy is defined as a condition of recurrent seizures, and medical intractability as recurrent seizures despite optimal treatment under the direction of an experienced neurologist over a 2-3-year period.

The **main purposes** of neuroimaging in epilepsy patients are to identify underlying **structural or metabolic abnormalities** that require specific treatment and to aid in formulating a **syndromic or etiological diagnosis**. Neuroimaging is even more important for those patients who have medically intractable seizures. Advances in technology to localize epileptogenic focus substantially improved the success of surgical treatment.

Structural disorders and metabolic disorders can be associated with seizure and detected on imaging. Structural disorders can be hippocampal or mesial temporal sclerosis, cortical developmental malformations or neuronal migration disorders, phakomatoses, vascular abnormalities, infections, neoplasms and scar epilepsy. However neoplastic causes of epilepsy are excluded in this study. Many imaging modalities for investigating seizures are available like neurosonogram, computerized tomography (CT scan) of brain, magnetic resonance imaging (MRI) of brain, functional MRI, positron emission tomography (PET), single photon emission computed tomography (SPECT). Of these various modalities **MRI is most important modality** for evaluation of structural disorders causing epilepsy. So, role of MRI to be evaluated in patients with epilepsy.

MATERIAL AND METHODS

This is a **prospective, single-center, observational study**. The Department of Imaging & Radiology provides a full range of services of diagnostic imaging to all the patients of hospital.

We included **60 patients** with epilepsy between the period of JUNE 2015 to JUNE 2016

Each patient was studied in detail with relevant to clinical history, examination and laboratory investigation.

They all are examined using magnetic resonance imaging of brain as prime diagnostic modality at our institution.

INCLUSION CRITERIA

- Patients with any of the following are included in present study.
- All patients with known case of partial seizures between age ranging since birth to 35 years.
- Onset of generalized or unclassified seizures in the first year of life, or in adulthood
- Evidence of a fixed deficit on neurological or neuropsychological examination
- Difficulty obtaining seizure control with first-line antiepileptic drugs (AEDs).

EXCLUSION CRITERIA


- All patients with seizures due to neoplasm
- All patients with electrolyte disturbances, alcohol and drug abuse
- All patients with neonatal asphyxia, perinatal trauma and congenital infections

RESULTS

A total 60 patients with up to 35 years of age were studied. The detailed information regarding Age, sex, clinical presentation, different lesion causing epilepsy with their radiological findings


and further management & follow up are given in master chart. Following data are observed in present study.

Table No.1 AGEWISE DISTRIBUTION OF LESION



Maximum number of patients were seen in age group 16 to 20 years. Youngest patient was 4 months old having lissencephaly & oldest patient was 34 yrs old having scar epilepsy.

Table No.2: SEXWISE DISTRIBUTION OF LESION



In study there were 36(60%) male patient & 24(40%) female patients

Table No.3: AGEWISE DISTRIBUTION OF PATIENTS WITH RESPECT TO SEX

Age Group(in years)	Male	Female	Total	Percentage
Up to 5	4	2	6	10%
6-10	5	4	9	15%
11-15	8	6	14	23.3%
16-20	14	5	19	31.7%
21-25	2	2	4	6.7%
26-30	2	2	4	6.7%
31-35	1	3	4	6.7%
TOTAL	36	24	60	100%

In present study maximum no. of male and female patients was in age group of 16 to 20 years.

Table No.4 CLINICAL PRESENTATION

Clinical presentation	No. of cases	Percentage
Partial Seizure	26	43.3%
(1) Simple	9	15%
(2) Complex	17	28.3%
Partial seizure with secondary generalization	14	23.3%
Generalized seizure	20	33.3%
(1) Absence Seizure	3	5%
(2) Generalized tonic clonic seizure	16	26.7%
(3) Myoclonic	1	1.7%
Focal neurological Sign	25	41.7%

In this study, the most common seizure presentation in patients was complex partial seizure (28.3 %) followed by generalized tonic clonic seizures (26.7 %).


Table No.5: DISTRIBUTION AND FREQUENCY OF VARIOUS LESIONS

Lesion	Total cases	% incidence
Cortical malformation	7	11.7%
Schizencephaly	1	1.7%

Lissencephaly	1	1.7%
Cortical heterotopia	2	3.3%
Focal cortical dysplasia	3	5%
Hippocampal pathology	14	23.3%
Unilateral hippocampal sclerosis	9	15%
Bilateral hippocampal sclerosis	5	8.3%
Infection	9	15%
Tuberculoma	6	10%
Neurocysticercosis	2	3.3%
Abscess	1	1.7%
Vascular Malformation	3	5%
Pail AVM	3	5%
Abnormal white matter	5	8.3%
Leukodystrophy(LD)	2	3.3%
Nonspecific hyperintensity	3	5%
Phakomatosis	2	3.3%
Tuberous sclerosis	1	1.7%
Sturge weber syndrome	1	1.7%
Gliosis(Scar Epilepsy)	4	6.7%
H/O previous trauma	3	5%
H/O cerebro vascular accident	1	1.7%
Normal MRI study	16	26.7%
Total	60	100%

In this study most common lesion was hippocampal pathology (23.3%) followed by tuberculoma (10%).

Table No.6: DISTRIBUTION OF LESION ACCORDING TO ETIOLOGY



Most of epilepsy are idiopathic (26.7%), however hippocampal pathology (23%) and infections (15%) are common causes of epilepsy.

Table No. 7: AGE WISE DISTRIBUTION OF VARIOUS LESIONS IN EPILEPSY

Age (in years)	Idiopathic	Cortical malformation	Hippocampal Pathology	Phakomatosis	Vascular malformation	Infection	Gliosis	Abn. White matter Hyperintensity	Total
<5	-	4	-	-	-	-	-	1	5
5-10	3	2	-	1	1	2	-	1	10
11-20	9	1	13	1	1	3	2	3	33
21-30	2	-	1	-	1	3	1	-	8
31-35	2	-	-	-	-	1	1	-	4
Total	16	7	14	2	3	9	4	5	60

Most common cause of epilepsy according to age group

First decade - cortical malformation.
 Second decade-hippocampal pathology
 Third decade-infection

Table No. 8: SEIZURE TYPEWISE DISTRIBUTION OF VARIOUS LESIONS IN EPILEPSY

	Partial seizure	Partial seizure with secondary generalisation	Generalized seizure	Total
Hippocampal pathology	11	3	-	14
Cortical malformation	4	2	1	7
Schizencephaly	1	-	-	
Lissencephaly	-	-	1	
Cortical heterotopia	1	1	-	
Focal cortical dysplasia	2	1	-	
Phakomatosis	-	1	1	2
Tuberous sclerosis	-	-	1	
Sturge weber Syndrome	-	1	-	
Vascular malformation	2	1	-	3
Infection	1	4	4	9
Abscess	-	1	-	
Tuberculoma	-	2	4	
Neurocysticercosis	1	1	-	
Gliosis	2	1	1	4
Abn. White matter hyperintensity	1	-	4	5
Leukodystrophy	-	-	2	
Nonspecific	1	-	2	
Normal MRI findings	5	2	9	16
Total	26	14	20	60

Most common lesion causing partial seizure is hippocampal sclerosis. While partial seizure with secondary generalization is commonly seen in infections, cortical malformation and idiopathic epilepsy. Generalized seizures is most common feature of idiopathic epilepsy followed by abnormal white matter hyperintensity and infection.

Table 9 - MRIFINDINGS OF DIFFERENTBRAIN LESIONS IN EPILEPTIC PATIENTS

MRI Finding	Number of cases	Percentage
Abnormal hyperintensity on T2W & FLAIR	37	61.6%
Cortical hyperintensity	3	5%
Hippocampal hyperintensity	12	20%
Subcortical hyperintensity	1	1.7%
White matter hyperintensity	19	31.7%
Unifocal	10	16.7%
Multifocal	6	10%
Diffuse	3	5%
Subependymal nodular grey matter	1	1.7%
Transmantle grey-matter lined cleft	1	1.7%
Abnormality on T1W-IR	21	35%
Hippocampal atrophy	12	20%
Cortical atrophy	1	1.7%
Cortical thickening with blurring of grey-white interface	2	3.3%
Thick grey matter with smooth interface with hour-glass configuration	1	1.7%
Subependymal/ subcortical nodular grey matter	2	3.3%
Transmantle grey-matter lined cleft	1	1.7%
Flow void on T2W and T1W	3	5%
Blooming on T2W-GRE	6	10%
Rim Post contrast enhancement on T1W	9	15%

Abnormal T1W hypointensity	19	31.7%
-----------------------------------	-----------	--------------

The most common findings on MRI were ,abnormalhyperintensity on T2W & FLAIR (61.6%).

Table No. 11: ROLE OF EPILEPSY PROTOCOL IN VARIOUS LESIONS IN EPILEPSY

LESION	Routine protocol	Epilepsy protocol+/- volumetry
Cortical malformation	5	7
Schizencephaly	1	1
Lissencephaly	1	1
Cortical heterotopia	1	2
Focal cortical dysplasia	2	3
Hippocampal pathology	8	14
Unilateralhippocampal sclerosis	6	9
Bilateralhippocampal sclerosis	2	5
Phakomatosis	2	2
Tuberous sclerosis	1	1
Sturge weber syndrome	1	1
Vascular Malformation	3	3
Pail AVM	3	3
Infection	9	9
Tuberculoma	6	6
Neurocysticercosis	2	2
Abscess	1	1
Gliosis(Scar Epilepsy)	4	4
H/O previous trauma	3	3
H/O cerebro vascular accident	1	1
Abnormal white matter	5	5
Leukodystrophy(LD)	2	2
Nonspecific hyperintensity	3	3
Normal MRI study	24	16

Out of 14 patients with hippocampal pathology diagnosed on epilepsy protocol with or without volumetry, only 8 patients were diagnosed with routine protocol. While 7 patients with cortical malformation were diagnosed on epilepsy protocol with or without volumetry, only 5 patients were picked up on routine protocol.

Table 12: FOLLOW UP AND FURTHER MANAGEMENT

Out of 60 patients presented with epilepsy 45% of patients responded to antiepileptic drugs, 13.3% of patients required specific medical management and 26.7 % of patients treated with surgical management. 5% of patients were expired. Rest of the patients could not be traced for follow-up.

Type of Lesions	Anti epileptic drugs	Specific Medical Mx	Surgical Mx	Expired	Follow-up not available	Total
Cortical malformation	2	-	2	1	2	7
Schizencephaly	1	-	-	-	-	1
Lissencephaly	-	-	-	1	-	1
Cortical heterotopia	1	-	-	-	1	2
Focal cortical dysplasia	-	-	2	-	1	3
Hippocampal pathology	3	-	11	-	-	14
Unilateral hippocampal sclerosis	2	-	7	-	-	9
Bilateralhippocampal sclerosis	1	-	4	-	-	5
Infection	-	8	1	-	-	9

Tuberculoma	0	6	-	-	-	6
Neurocysticercosis	0	2	-	-	-	2
Abscess			1			1
Vascular Malformation	1	-	2	-	-	3
Pail AVM	1	-	2	-	-	3
Abnormal white matter	3	-	-	2	-	5
Leukodystrophy (LD)	-	-	-	2	-	2
Nonspecific hyperintensity	3	-	-	-	-	3
Phakomatosis	2	-	-	-	-	2
Tuberous sclerosis	1	-	-	-	-	1
Sturge weber syndrome	1	-	-	-	-	1
Glios(Scar Epilepsy)	4	-	-	-	-	4
H/O previous trauma	3	-	-	-	-	3
H/O cerebro vascular accident	1	-	-	-	-	1
Normal MRI study	12	-	-	-	4	16
TOTAL	27	8	16	3	6	60

DISCUSSION

- Among all the patients with epilepsy causes in descending order are idiopathic (26.7%) followed by hippocampal pathology (23.3%), Infections(15%), Cortical malformation(11.7%), abnormal white matter hyperintensity (8.3%), gliosis (6.7%), AVM(5%) and phakomatoses (3.3%).
- Among epileptic patients with abnormal MRI findings, unilateral hippocampal sclerosis was most common structural lesion comprising 15% followed by tuberculoma comprising 10%.
- Among patients with epilepsy, cortical malformations was most common cause of epilepsy in <10 years of age. While hippocampal pathology and infections were commonest structural abnormality in 11-20years and 20-35 years of age respectively.
- Most common lesion causing partial seizure is hippocampal sclerosis (42.3%). While partial seizure with secondary generalization is commonly seen in infections (28.6%). Generalized seizures is most commonly associated with idiopathic epilepsy (45%) followed by abnormal white matter hyperintensity and infections (20%).
- Most of structural brain abnormalities lead to epilepsy were located in temporal lobe (30%) followed by frontal lobe(13.3%). Thus temporal lobe lesions were more epileptogenic.
- Abnormal white matter hypointensity on T1W images and hyperintensity on T2W images were most common finding on MRI study comprising 31.7%.

CONCLUSION

MRI reveals significant number of structural abnormalities in patients with focal seizures, symptomatic generalized seizures, refractory epilepsy, patients with focal neurological deficit and patients with early(infants) or late(adult) onset of epilepsy. Thus incidence of structural abnormality is more in this group of patients.

Structural abnormalities are more common with partial seizure than generalized seizures. Underlying cause of epilepsy varies significantly with age.

Epilepsy protocol has better yield, sensitivity and specificity than routine protocol in patients with hippocampal pathology and

cortical malformation. In order to maximize the potential of the technique, the protocol should be crafted to the needs of the individual epilepsy patient.

MRI has a major impact on management of epileptic patients. MRI identify causative lesion and localize it, thus helps to identify further line of management (medical or surgical). MRI is useful to plan surgical management and for post-operative followup.

Thus, we conclude that MRI is most important & reliable structural neuroimaging modality to evaluate the patients with epilepsy.

REFERENCES

1. Blume WT, LAders HO, .ET al Glosaery of descriptive terminology for ictal semiology; report of the ILAE task force on classification and terminology epilepsya 2001m 42.
2. Commission on classification and terminology of the international league against epilepsy proposal for revised clinical and electroencephalographic classification of epileptic seizures epilepsya 1981.22
3. Engel J Jr. report of the ILAE classification core grupu epilepsya 2006.47
4. Engel J Jr. Pedley TA (eds), epilepsy a comprehensive textbook vols 1, 2 and 3 Philadelphia Lippincott-raven publishers 1997.
5. Jack C, Bentley MD Twomey CK Zinsmeister AR MRI based volume measurements of the hippocampal formation and anterior temporal lobe validation studies Radiology 1990, 176, 205-209.
6. Elster A Gradient - echo MRI imaging techniques and acronyms Radiology 1993. 186(1) 1-B.
7. Jackson G new techniques in magnetic resonance and epilepsy radiology 1994-35 (suppl 6)S2-S13.
8. Barkovich A, Kjos B Gray matter heterotopias MR characteristics and correlation with development and neurologic manifestations Radiology 1992. 182(2) 493-499.
9. Smith A Weinstein MA, Quencer RM, Muroff LR, Stonesifer KJ, Li FC, Wener L, Solomon MA, Cruse RP, Rosenberg LH. Association of heterotopic gray matter with seizures. MR imaging work in progress. Radiology 1988, 168(1) 195-198.
10. Eksioglu Y, Scheffer IE, Cardenas P Knoll J, DiMario F, RamsbyG., BergM, Kamurok, Berkovic SF, DuyK GM, Parisi J, Huttenlocher PR, Walsch CA, periventricular heterotopia an X-linked dominant epilepsy locus causing aberrant cerebral cortical development neuron 1996. 16(1) 77-87.
11. Taylor DC, Falconer MA, Bruton CJ, et al. Focal dysplasia of the cerebral cortex in epilepsy. J Neurol Neurosurg Psychiatry 1971. 34 369-87.
12. Barkovich AJ subcortical heterotopia a distinct clinicoradiology entity Am J Neuroradiol 1996. 17(7) 1315-1322.
13. Palmimi A Andermann F, Aicardi J, Dulac O, Chaves F, Ponsot G, Pinard JM, Goutieres F, Livingston J, Tampieri D, Diffuse cortical dysplasia or the double cortex syndrome the clinical and epileptic spectrum in 10 patients Neurology 1991, 41(10) 1656-1662.
14. Shepherd C, Beard CM,Gomez MR, Kurland LT, Whisnant JP, Tuberous sclerosis complex in Olmsted County , Minnesota, 1950-1998. Arch Neurol 1991, 48(4) 400-401.
15. Schmauzer I, Bittner R, MR-imaging findings in children with Sturge-Weber syndrome neuropediatrics 1990, 21, 146-152.
16. Jackson GD, Briellmann RS, Kuzniecky RL, temporal lobe epilepsy in kuzniecky RL, Jackson GD .ed.
17. Jennett W, Lewin W,Traumatic epilepsy after closed head injuries J. neurol neurosurg Psychiatry 1960, 23 295-301.
18. Luhdorf K, Jensen L, Plesner A, etiology of seizures in the elderly epilepsya 1986 27(4) 458-463.
19. Kotila M, Waltimo O epilepsy after stroke Epilepsia 1992, 33(3) 495-498.
20. Aebi C, Kaufmann F, Schaad U Brain abscess in childhood-long-term experiences Eur J Pediatr 1991, 150(4) 282-286.
21. Choi K, A clinical study of adult onset seizure disorder J Korean Med Assoc 1986 29189-197.
22. Carbalal JR, Palacios E, Azar Kia B, Churchill R Radiology of cysticercosis of the central nervous system including computed tomography radiology 1977. 125(1) 127-131.
23. Creasy J, Alarcon J magnetic resonance imaging of neurocysticercosis top magnetic resonance imaging 1994 6 (1) 59-68.
24. Oguni H, Andermann F, Rasmussen T the syndrome of chronic encephalitis and epilepsy A study based on the MNI senes of 48 cases. Adv neurol 1992 57 419-433.
25. De vivo DC, Hirano M, DiMauro S, Mitochondrial disorders in Moser HW ed handbook of clinical Neurology Amsterdam Elsevier 1997 389-446.
26. Briellmann RS, Pell GS, Wellard RM, et al MR imaging of epilepsy state of the art at 1.5 T and potential of 3.T Epileptic Disorder 2003, 5. 3a-332.
27. Connelly A, MR diffusion and perfusion in epilepsy in Kuzniecky RL, Jackson GD, eds. Magnetic Resonance in Epilepsy . Neuroimaging Techniques 2nd ed. London Elsevier 2005, 315-332.
28. Jackson GD, Berkovic SF, Duncan JS, et al, Optimizing the diagnosis of hippocampal sclerosis using MR-imaging Antero-medial J neuroradiol 1993, 14, 753-762.
29. Magnetic Resonance in Epilepsy Neuroimaging Techiques 2nd ed. London Elsevier, 2005 99a-176.
30. Jackson GD, Kuzniecky RL, Pell GS, principles of magnetic resonance imaging in kuzniecky RL, Jack GD eds. Magnetic Resonance in epilepsy neuroimaging Techniques 2nd ed. London Elseveir 2005.