



**ORIGINAL RESEARCH PAPER**

**Medical Science**

**A PROSPECTIVE COMPARATIVE STUDY BETWEEN TRANSVAGINAL SONOGRAM AND HISTOPATHOLOGICAL EXAMINATION IN WOMEN WITH PERIMENOPAUSAL AND POSTMENOPAUSAL BLEEDING.**

**KEY WORDS:** Transvaginal, Histopathology, postmenopausal bleeding, endometrial thickness.

**DR.K. SHANTHI ELANGO**

MD, DGO, GOVT. RSRM LYING IN HOSPITAL, ATTACHED TO STANLEY MEDICAL COLLEGE, ROYAPURAM, CHENNAI-13, TAMILNADU.

**Dr. ANITHA VIRGIN KUMARI**

MD (O.G), DGO, GOVT. RSRM LYING IN HOSPITAL, ATTACHED TO STANLEY MEDICAL COLLEGE, ROYAPURAM, CHENNAI-13, TAMILNADU.

**ABSTRACT**

**Aim:**To determine the value of transvaginal sonography in detecting endometrial pathology, and its correlation with histopathology in cases of perimenopausal and postmenopausal beeding.  
**Methods:**It is a prospective study done on 100 women who were enrolled between August 2016 to July 2017. Out of 100 cases, 50 women were in perimenopausal age group and 50 were in postmenopausal age group.  
**Result:**From this study 50% of women with atrophic endometrium had a mean endometrial thickness <5mm and 50% of women with endometrial pathology had a thickness>5mm.  
**Conclusion:**TVS is useful as a first step diagnostic procedure in the evaluation of peri and postmenopausal bleeding.This study proves that this diagnostic tool correlates well with the histopathology findings.TVS with HPE correlation helps us to identify atypical hyperplasia at the earliest which is a precursor of adenocarcinoma in 28%.

**INTRODUCTION:**

Abnormal Uterine bleeding (AUB) is one of the most frequently encountered conditions in gynaecologic practice and forms about 10% of all gynaecological admissions. A menstrual abnormality is the cause of discomfort, inconvenience and disruption of healthy, social and physical life styles among women worldwide. It is responsible for more than 23.6% of all visits to our outpatients clinic and may accounts for more than 24% of all hysterectomies.

The main concern in postmenopausal perimenopausal bleeding is, that the bleeding could be only the external manifestation of a hidden serious pathology such as endometrial carcinoma. There are many diagnostic modalities for evaluating endometrial pathologies in women who have AUB. These include dilation & Curettage, and hysteroscopy. Recently transvaginal sonogram (TVS) has permitted the use of higher frequency ultrasound waves at greater proximity of the uterus. It is relatively cheap, simple outpatient method of studying the endometrium for detecting malignant lesions or their precursors at an earlier stage. It is a non invasive diagnostic modality of studying the endometrial pattern and its thickness and at the same time to exclude organic pathology in case of AUB. The thickness and internal echo texture of the endometrium in the various phases of the menstrual cycle as seen in transvaginal sonogram correlates well with endometrial pathology.

**AIM OF THE STUDY:**

To correlate the findings transvaginal sonogram and histopathological examination in women with peri ad postmenopausal bleeding namely transvaginal sonogram and histopathological examination.

**METHODS:**

It is a prospective study of 100 women were enrolled who were coming to OPD of Stanley Medical College, Obstetrics and Gynaecology Department. 50 women with perimenopausal bleeding (age >40yrs not attained menopause) and 50 women with postmenopausal bleeding (age >40 yrs who attained menopause). Patients included were with H/o of excessive bleeding per vaginum, who were >40yrs and both nulliparous and parous women. Patients with medical disorders, ca cervix, fibroid polyp which is visualized by speculum examination and adnexal swelling were excluded. These women examined with detailed history, pelvic examination with basic lab investigations & TVS done for endometrial thickness along with any pathology, later they were subjected to fractional curettage for histopathological study and results were correlated.

**RESULTS:**

In our study, out of 100 women who were in the age group of >40 yrs were included. These were divided into 2 group like, perimenopausal age as group A, and post menopausal age as group B. 50%of cases in our study were in perimenopausal age and 50% were in post menopausal age.

**COMPARISON OF HPE WITH TVS MEASUREMENT OF ENDOMETRIAL THICKNESS IN GROUP A (POST MENOPAUSAL BLEEDING)**

NO	HPE REPORT	NO.OF CASES	MEAN ENDOMETRIAL THICKNESS IN MM	PERCENTAGE
1.	Atrophic Endometrium	18	3.87mm(2-5)	36%
2.	Tissue Insufficient for Diagnosis (TIFD)	8	3.58mm(2.6-4)	16%
3.	Cystoglandular hyperplasia	7	13.57mm(11-17)	14%
4.	Nonsecretory endometrium	3	7mm(6-8)	6%
5.	Secretory endometrium	6	9.16mm(8-10)	12%
6.	Pyometra/Endometritis	4	9mm(8-11)	8%
7.	Endometrial carcinoma	4	17.2mm(16-18)	8%

Thus it can be seen that 50 % of women with atrophic endometrium had a mean endometrial thickness <5mm and 50% of women with endometrial pathology namely proliferative phase, hyperplasia, carcinoma of endometrium amd pyometra had a mean thickness >5mm.

**COMPARISON OF HPE WITH TVS MEASUREMENT OF ENDOMETRIAL THICKNESS IN GROUP B (PERI MENOPAUSAL BLEEDING)**

NO	HPE REPORT	NO.OF CASES	MEAN ENDOMETRIAL THICKNESS IN MM	PERCENTAGE
1.	Adenomatous hyperplasia	1	22mm	2%
2.	Proliferative Endometrium	4	8.75mm(8-10)	8%
3.	Cystoglandular hyperplasia	11	15.36mm(13-18)	22%

4.	Nonsecretory endometrium	26	8.73mm(6-11)	52%
5.	Secretory endometrium	4	10.25mm(9-12)	8%
6.	Endometritis	1	18mm	2%
7.	Endometrial carcinoma	2	25.5mm(25-28)	4%
8.	Atypical hyperplasia	1	24mm	2%
<b>TOTAL</b>		<b>50</b>		

Therefore it can be seen that endometrial thickness <13mm the HPE report was normal endometrium either secretory or nonsecretory. An endometrial thickness >13mm has been found to be associated with hyperplasia including one adenomatous and one atypical hyperplasia.

**COMPARISON OF HPE WITH TVS MEASUREMENT OF ENDOMETRIAL VOLUME IN GROUP A (POST MENOPAUSAL BLEEDING)**

NO	HPE REPORT	NO.OF CASES	MEAN ENDOMETRIAL THICKNESS IN CM <sup>3</sup>	PERCENTAGE
1.	Atrophic Endometrium	18	12.72(8.46-14.2)	36%
2.	Tissue Insufficient for Diagnosis (TIFD)	8	11.5(10.45-12.46)	16%
3.	Cystoglandular hyperplasia	7	59.3(28.8-95.7)	14%
4.	Nonsecretory endometrium	3	43.87(43.16-45.3)	6%
5.	Secretory endometrium	6	48.78(34.6-63.4)	12%
6.	Pyometra/Endo metritis	4	49.33(26.8-68.5)	8%
7.	Endometrial carcinoma	4	136(102.3-175.8)	8%
<b>TOTAL</b>		<b>50</b>		

This table shows mean endometrial volume was <13 cm<sup>3</sup> HPE report is atrophic endometrium. An endometrial volume >13cm<sup>3</sup> is associated with endometrial pathology.

**DISCUSSION:**

Abnormal uterine bleeding is common in peri menopausal & postmenopausal women. The etiology varies from simple dysfunctional uterine bleeding to benign lesions like polyp, fibroid & malignancies. Apart from the clinical examination, various diagnostic modalities are available to confirm our diagnosis. In our study the efficacy of trans vaginal sonogram for diagnosing abnormal uterine bleeding & its correlation with histopathology of endometrium by fractional curettage were studied. In our study 100 cases of AUB were enrolled, out of which, 50 cases (50%) in peri menopausal and 50 cases (50%) in post menopausal age group respectively.

Regarding endometrial thickness by TVS and results were correlated with HPE reports. In postmenopausal group 18(36%) had atrophic endometrium, 8(16%) had Tissue insufficient for diagnosis, 7(14%) had endometrial hyperplasia with ET Thickness of 13.57mm, 4(8%) had endometrial carcinoma (ET Thickness-17.2mm). In peri menopausal group 11(22%) had endometrial hyperplasia with ET Thickness of 15.36mm, 1(2%) had adenomatous hyperplasia, another 2(4%) had endometrial carcinoma of ET Thickness (25.5mm).

The endometrial volume in women with postmenopausal bleeding ranged from 8.46cm<sup>3</sup> in atrophic endometrium to 175.84cm<sup>3</sup> in a patient with endometrial carcinoma

In the present study of 50 women with postmenopausal bleeding using 5 mm and perimenopausal bleeding using 13mm as a cut off limit, vaginal sonography successfully diagnosed endometrial

abnormality. The specificity and sensitivity were both 100%. There were no false negative or false positive with a positive predictive value of 100% and a negative predictive value of 100%.

**CONCLUSION:**

Transvaginal sonogram is a simple, non invasive convenient way to indirectly visualise the endometrial cavity. It is useful as a first step diagnostic procedure in evaluation of peri and postmenopausal bleeding. This study proves that the diagnostic tool correlates well with its histopathological findings. Intrauterine pathology of the endometrium and myometrium are well delineated and endometrial carcinoma detected with precision. It can find out associated ovarian pathology. It appears that . ultrasonogram will continue to take the role of stethoscope for gynaecologist.

**REFERENCES:**

1. Ettinger B: Optimal use of post menopausal Hormone Replacement therapy. *Obstet. Gynaecol.* 1988, 72:315-365.
2. Najeeb R, Awan AS, Bakhtiar U, Akhter S. Role of TVS in assessment of abnormal uterine bleeding in perimenopausal age group. *J Ayub Med Coll Abbottabad* 2010; 22:87-90.
3. Vilos GA, Lefebvre G, Graves GR. Guidelines for the management of abnormal uterine.
4. Speroff L, Fritz MA. Menopause and the perimenopausal transition, clinical endocrinology. In; Speroff L, Fritz MA, editor. *Clinical gynecologic endocrinology and infertility.* 7th ed. Philadelphia, London: Lippincott Williams & Wilkins, 2005:628.
5. Goldstein SR. The endometrial echo revisited: have we created a monster? *Am J Obstet Gynecol.* 2004; 191:1092-6.
6. Vakian M, Vavilis D, Agorastos T; Histopathology findings of endometrium in patients with DUB; *clinics Exp. Obstetrics Gynecology*; 1996; 23(4):236-9.
7. Mark H Emanuel, MD, Marion J, Verdel, MD, Kees Wamstekjer, MD, PhD, and Frits B. Lammes, MD, PhD, A prospective comparison of TVS and diagnostic hysteroscopy on the evaluation of patients with AUB. *Clinical implications. AmJ Obstet Gyne* 1995; 172:547-52.
8. Anjali singh, Saroj singh, Veena mathur, Kalpana singh; TVS in DUB and its correlation with histopathology; *Journal of obste and Gyne of India*, 2001; 51(6): 116-121.