



ORIGINAL RESEARCH PAPER

Medicine

HIGH LEVEL RESISTANCE TO GENTAMICIN AND VANCOMYCIN IN ENTEROCOCCI ISOLATED FROM POULTRY IN ARGENTINA

KEY WORDS: antimicrobial, resistance, enterococci, poultry

Gastón Delpech

Microbiología Clínica, Medicina, Universidad Nacional del Centro de la Provincia de Buenos Aires, Av. Pringles 4375 (7400), Olavarría, Argentina.

Mónica Sparo*

Microbiología Clínica, Medicina, Universidad Nacional del Centro de la Provincia de Buenos Aires, Av. Pringles 4375 (7400), Olavarría, Argentina. *Corresponding Author

Beatriz Baldaccini

Microbiología Clínica, Medicina, Universidad Nacional del Centro de la Provincia de Buenos Aires, Av. Pringles 4375 (7400), Olavarría, Argentina.

Mónica Ceci

Centro de Estudios Bioquímicos, Av. Colón 1388 (7000), Tandil, Argentina.

Leonardo García Allende

Centro de Estudios Bioquímicos, Av. Colón 1388 (7000), Tandil, Argentina.

ABSTRACT

In chicken husbandry, antimicrobial resistance among enterococci has been observed. Glycopeptides and gentamicin Minimum Inhibitory Concentrations in fecal poultry enterococci were determined. Glycopeptide and high-level gentamicin resistance (HLGR) genes were investigated. *E. faecalis* (n = 105) and *E. faecium* (n = 70) were the predominant species. In *E. faecalis*, HLGR (19%) was observed. In high-level gentamicin resistant *E. faecalis*, presence of *aac* (6')-*le-aph* (2'')-*la* gene was confirmed. *VanA* *E. faecium* (7.1%) were found, with HLGR (20%) and high-level streptomycin resistance (40%). *vanA* genotype was detected in glycopeptide resistant *E. faecium* (100%). These results highlight the presence of horizontally transferable antimicrobial resistance determinants in *E. faecalis* and *E. faecium* recovered from poultry. Existence of enterococcal reservoirs in poultry should be taken into account as a potential source of antimicrobial resistance genes, strengthening their potential spread to humans through the food chain of horizontal transferable resistance determinants.

INTRODUCTION

The emergence of antimicrobial resistance is a priority for human health. In this sense, detection of reservoirs of potentially pathogenic bacteria with transferable resistance mechanisms that may spread from the environment to man is an important challenge¹⁻³.

In animal husbandry, antimicrobials are heavily used for prevention and treatment of bacterial diseases; but also as growth promoters in animal feed^{4,6}. Among Gram-positive bacteria, the genus *Enterococcus* shows a wide distribution in the environment. Enterococci are recognized as part of the indigenous microbiota of animal and human gastrointestinal tract. In fecal samples from farm animals, such as poultry, resistance to antimicrobials used in humans (e.g. aminoglycosides, glycopeptides) has been observed^{7,8}.

Enterococci can acquire high-level gentamicin resistance (HLGR) determinants by horizontal gene transfer^{2,9}. Their expression result in the loss of synergistic bactericidal effect achieved with cell wall-active agents, posing a serious therapeutic problem for invasive infections such as enterococcal endocarditis^{10,11}.

Among enterococci with HLGR, *aac* (6')-*le-aph* (2'')-*la* is the most prevalent gene, which encodes a bifunctional enzyme, AAC(6')-APH(2''), that confers resistance to aminoglycosides, except for streptomycin. Other monofunctional chromosomal [e.g. *aph* (2'')-*lb*, *aph* (2'')-*ld*] and plasmidic [e.g. *aph* (2'')-*lc*] genes also encode gentamicin resistance¹²⁻¹⁴.

Glycopeptide resistance in enterococci is mediated by *van* genotypes. Expression of *vanA* gene leads to inducible high-level vancomycin (Minimum Inhibitory Concentration, MIC \geq 64 μ g/ml) and teicoplanin (MIC \geq 16 μ g/ml) resistance¹⁵.

In farm animals, such as poultry, *E. faecalis* are antimicrobial resistance determinants' reservoirs (HLGR, glycopeptide)¹⁶⁻¹⁷. In addition, there is evidence supporting the feasibility of genetic determinants transfer between poultry and human *E. faecalis*¹⁸. In Argentina, however, there is no available information in this regard.

The aim of this study was to investigate HLGR and vancomycin resistance determinants in fecal poultry enterococci isolated from farms located at a region in Argentina.

Material and methods

Fecal samples (N = 120) were randomly collected from two poultry farms (GPT-A; GPT-B) in Tandil district, Buenos Aires Province (Argentina), from March to September 2015. GPT-A: a 9,000 broilers/year average production. GPT-B: a 7,000 broilers/year average production.

Samples (0.5 g) were added to 5 mL of Azide-Glucose broth. Cultures at 35 \pm C for 18 h in Bile-Esculin Azide agar (BEA) were carried out. Black pigmented colonies were selected for characterization at species level¹⁹. Genotypic confirmation by PCR for *tuf* and *ddl* genes (Table 1) was performed^{1,20}.

Resistance to vancomycin, teicoplanin, gentamicin and streptomycin was investigated. MIC was determined by the agar dilution method. Quality control strains: *E. faecalis* ATCC 29212 and *E. faecalis* ATCC 51299²¹.

Detection of glycopeptide resistance (*vanA*, *vanB*) and HLGR [*aac* (6')-*le-aph* (2'')-*la*, *aph* (2'')-*lb*, *aph* (2'')-*lc*, *aph* (2'')-*ld*] genes was carried out by PCR (Table 1)^{20,22}.

Table 1. Primers employed in PCR.

Gene	Primers (sequence 5' to 3')	Reference
<i>Tuf</i>	TACTGACAAACCATTTCATGATG AACTTCGTCCACCAACGCGAAC	²⁰
<i>ddl</i> _{<i>E. faecalis</i>}	ATCAAGTACAGTTAGTCT ACGATTCAAAGCTAACTG	²⁰
<i>ddl</i> _{<i>E. faecalis</i>}	TAGAGACATTGAATATGCC TCGAATGTGCTACAATC	²⁰
<i>vanA</i>	GGGAAAACGACAATTGC GTACAATGCGGCCGTTA	²⁰
<i>vanB</i>	ATGGGAAGCCGATAGTC GATTTCGTCTCGACC	²⁰

<i>aac</i> (6') - <i>le-aph</i>	GAGCAATAAGGGCATACCAAAAATC CCGTGCATT TGTCTTAAAAAAGTGG	22
<i>aph</i> (2'')- <i>lb</i>	TATGGATTGATGGTTAACTTGACGCTGAG ATTAAGCTTCCTGCTAAAATATAACATCTC	22
<i>aph</i> (2'')- <i>lc</i>	GAAGTGATGGAAATCCCTTCGTG GCTCTAACCCCTCAGAAATCCAGTC	22
<i>aph</i> (2'')- <i>ld</i>	GGTG GTTTTACAGGAATGCCATC CCCTCTCATACCAATCCATATAACC	22

RESULTS

Enterococci were isolated from 80% (96/120) of the poultry fecal samples. Phenotypic and genotypic characterization showed that *E. faecalis* (n = 105) was the predominant enterococcal species found (52.5%), followed by *E. faecium* (n = 70).

In 19% (20/105) of poultry *E. faecalis* (FLS-29, FLS-38, FLS-59, FLS-77, FLS-81, FLS-84, FLS-90, FLS-91, FLS-100, FLS-103, FLS-110, FLS-115, FLS-122, FLS-128, FLS-138, FLS-146, FLS-150, FLS-153, FLS-164, FLS-171), HLGR was observed (MIC_{gentamicin} = 512-1,024 µg/mL). In high-level gentamicin resistant *E. faecalis* (100%), presence of *aac* (6') -*le-aph* (2'')-*la* gene was confirmed by PCR (Fig. 1a). High-level streptomycin resistance (HLSR) was not detected in *E. faecalis* isolates (MIC_{streptomycin} < 2,000 µg/mL). Glycopeptide resistant *E. faecalis* were not recovered from any of the fecal samples analyzed in this study (MIC_{vancomycin} < 4 µg/mL; MIC_{teicoplanin} < 8µg/mL).

Vancomycin and teicoplanin resistant *E. faecium* (7.1%, 5/70) were found (Table 2). In all cases, VanA phenotype was observed (MIC_{vancomycin} = 64-1,024 µg/mL; MIC_{teicoplanin} = 64-512 µg/mL). Molecular analysis confirmed *vanA* genotype in all glycopeptide resistant *E. faecium* (Fig. 1b). Glycopeptide resistant *E. faecium* (FCM-70) showed HLGR (20%) and HLSR (40%; FCM-08, FCM-62). In all HLGR *E. faecium* isolates, *aac* (6') -*le-aph* (2'')-*la* gene was detected.

Table 2. Glycopeptide and high-level aminoglycoside resistance in poultry *E. faecium*.

Isolate	Genotype	MIC _{van}	MIC _{tei}	HLGR genotype	MIC _{gen}	MIC _{str}
FCM-14	<i>vanA</i>	64	64	ND ^a	128	128
FCM-22	<i>vanA</i>	128	128	ND	256	2,048
FCM-70	<i>vanA</i>	64	128	<i>aac</i> (6') - <i>le-aph</i> (2'')- <i>la</i>	2,048	256
FCM-93	<i>vanA</i>	512	128	ND	256	256
FCM-97	<i>vanA</i>	1,024	512	ND	256	2,048

Genotype: glycopeptide resistance gene. MICs in µg/mL. van: vancomycin; tei: teicoplanin. HLGR: High-level gentamicin resistance. gen: gentamicin; streptomycin. ND: not detected.




Fig. 1. a) PCR for *aac* (6') -*le-aph* (2'')-*la* gene (369 bp) detection in cattle enterococcal isolates. Ladder 1: *Enterococcus faecalis* FLS-29. Lane 2: *E. faecalis* FLS-38. Lane 3: *E. faecalis* FLS-59. Lane 4: *E. faecalis* FLS-77. Lane 5: *E. faecalis* FLS-81. Lane 6: *E. faecium* FCM-70. Lane 7: MW marker. b) PCR for *vanA* gene (732 bp) detection in cattle enterococci. Lane 1: MW marker. Lane 2: *Enterococcus*

faecium FCM-14. Lane 3: *E. faecium* FCM-22. Lane 4: *E. faecium* FCM-70. Lane 5: *E. faecium* FCM-93. Lane 6: *E. faecium* FCM-97.

DISCUSSION

Over the last years, *E. faecalis* showed a dualistic behavior. Biotechnological and immunomodulatory properties have been proven in several strains²³⁻²⁵. However, nowadays, is considered as one of the most prevalent health-care associated infections pathogens²⁶.

In our investigation, *E. faecalis* was the most frequent isolated enterococci. In previous reports, it was the most (46.1%) or the least (11.5%) prevalent recovered species from poultry^{16,27}.

In this study, high-level gentamicin resistant *E. faecalis* (19%) were recovered and *aac* (6') -*le-aph* (2'')-*la* gene was detected. A low prevalence of enterococci harboring *aac* (6') -*le-aph* (2'')-*la* gene (2.7%) was observed for Australian poultry²⁷. Likewise, *aac* (6') -*le-aph* (2'')-*la* gene was detected as the only HLGR determinant in *E. faecalis* isolated from poultry in Korea (10.9%). In addition, all high-level gentamicin resistant enterococci showed co-expression of HLGR¹⁷. Interestingly, the most frequent HLGR gene detected in human enterococci is *aac* (6') -*le-aph* (2'')-*la*²⁸.

Recently, our group reported the presence of HLGR in *E. faecalis* from food of animal origin, as well as *in vivo* horizontal transfer of this resistance between food and human enterococci^{1,2}. Therefore, horizontal transfer of resistance determinants between enterococci from different origin could not be ruled out as source of antimicrobial resistance.

In Argentina, in human enterococci, antimicrobial resistance has been studied. *E. faecalis* has been reported as the most prevalent enterococcal species expressing HLGR in isolates recovered from hospitalized patients²⁹.

Vancomycin-resistant *E. faecalis* were not recovered during this investigation. In 2007, vancomycin resistance was found in 20% of poultry enterococci from Slovakia, mainly in *E. faecalis*³⁰. Chan *et al.*¹⁶ also found *aac* (6') -*le-aph* (2'')-*la* and *vanA* genes in *E. faecalis* (4%) recovered from Korean poultry. In Europe, previous studies have shown the relevance of these genetic determinants of HLGR and glycopeptide resistance in human enterococci^{28,31}.

Nowadays, *E. faecium* is also considered another enterococcal species with clinical relevance²⁶. In addition, previous reports also found that *E. faecium* was the second commonest enterococcal isolates^{16,17}.

HLGR was observed in vancomycin-resistant *E. faecium* (20%). In poultry enterococcal isolates from China, *aac* (6') -*aph* (2'') gene was detected in *E. faecium* although HLGR was not confirmed. Furthermore, in Australian range-meat chickens and indoor-meat chickens, it was reported the recovery of gentamicin-resistant *E. faecium*^{16,17}.

Along this investigation, glycopeptide-resistant *E. faecium* were recovered (7.1%). VanA phenotype is encoded by gene *vanA*, typically carried in plasmid-harbored transposons. The most frequent glycopeptide resistance genotype in clinical enterococci is *vanA*³². Nowadays, there is an increasing concern about *vanA* plasmid-mediated transfer to methicillin-resistant *Staphylococcus aureus*^{15,33}. In Chinese poultry, lower vancomycin resistance prevalence (1.3%) was detected for *E. faecium*, found in poultry drinking water. Recently, In Argentina, on the contrary, VanA *E. faecium* have been recovered from food of animal origin^{1,16}.

In a local nation-wide survey about resistance in clinical enterococci most of the *E. faecium* strains carried *vanA* gene (98%), expressing HLGR (77.2%) and high-level streptomycin resistance (95.8%) as well³².

In countries like the United States and in the European Union the use of antimicrobials in husbandry is documented. Aminoglycosides (gentamicin) are employed in poultry^{5,6}.

However, glycopeptides, such as avoparcin, were banned due to its linkage with spread of vancomycin-resistant enterococci. After the ban it was observed a significant decrease of glycopeptide resistance in Danish poultry isolates⁹. However, in Argentina, the lack of consistent information about antimicrobial use in poultry husbandry does not allow to rule out this practice as a source of glycopeptide or aminoglycoside resistance in poultry enterococci. It is important to note that in the two studied farms, antimicrobials were frequently administered.

This study reports the presence of horizontally transferable antimicrobial resistance determinants (HLGR, *vanA*) in *E. faecalis* and *E. faecium* recovered from poultry in the Province of Buenos Aires, Argentina. The existence of these enterococcal reservoirs in farm animals such as poultry should be taken into account as a potential source of antimicrobial resistance genes, strengthening their potential spread to humans through the food chain.

REFERENCES

1. Delpech G, Pourcel G, Schell C, De Luca M, Basualdo J, Bernstein J, Grenovero S, Sparo M. Antimicrobial resistance profiles of *Enterococcus faecalis* and *Enterococcus faecium* from artisanal food of animal origin in Argentina. *Foodborne Pathog Dis*. 2012; 9: 939-44.
2. Sparo M, Urbizu L, Solana MV, Pourcel G, Delpech G, Confalonieri A, Ceci M, Sánchez Bruni S. High-level resistance to gentamicin: Genetic transfer between *Enterococcus faecalis* isolated from food of animal origin and human microbiota. *Lett Appl Microbiol*. 2012; 54: 119-25.
3. Chioro A, Coll-Seck AM, Hrie B, Moeloeck N, Motsolaedi A, Rajatanavin R, Touraine M. Antimicrobial resistance: a priority for global health action. *Bull World Health Organ*. 2015; 93: 439.
4. van den Bogaard AE, Stobberingh EE. Antibiotic usage in animals-Impact on bacterial resistance and public health. *Drugs*. 1999; 58: 589-607.
5. Schwarz S, Kehrenberg C, Walsh TR. Use of antimicrobial agents in veterinary medicine and food animal production. *Int J Antimicrob Agents*. 2001; 17: 431-7.
6. Mathew AG, Cissell R, Liamthong S. Antibiotic resistance in bacteria associated with food animals: a United States perspective of livestock production. *Foodborne Pathog Dis*. 2007; 4: 115-33.
7. Hayes JR, English LL, Carr LE, Wagner DD, Joseph SW. Multiple-antibiotic resistance of *Enterococcus* spp. isolated from commercial poultry production environments. *Appl Environ Microbiol*. 2004; 70: 6005-11.
8. Ruzauskas M, Virgailis M, Šlugždinienė R, Sužiedėlienė E, Šeputienė V, Daugėlavicius R, Zienius D, Šengaut J, Pavilionis A. Antimicrobial resistance of *Enterococcus* spp. isolated from livestock in Lithuania. *Veterinariški Arhiv*. 2009; 79: 439-49.
9. Hollenbeck BL, Rice LB. Intrinsic and acquired resistance mechanisms in enterococcus. *Virulence*. 2012; 3: 421-33.
10. Leblanc DJ. 2006. *Enterococcus*. In: *The Prokaryotes: a handbook on the biology of bacteria*. New York: Springer.
11. Munita JM, Arias CA, Murray BE. Enterococcal endocarditis: can we win the war? *Curr Infect Dis Rep*. 2012; 14: 339-49.
12. Kobayashi N, Alam M, Nishimoto Y, Urasawa S, Uehara N, Watanabe N. Distribution of aminoglycoside resistance genes in recent clinical isolates of *Enterococcus faecalis*, *Enterococcus faecium* and *Enterococcus avium*. *Epidemiol Infect*. 2001; 126: 197-204.
13. Ramirez MS, Tolmasky ME. Aminoglycoside modifying enzymes. *Drug Resist Updat*. 2010; 13: 151-71.
14. Sparo M, Delpech G. 2014. High-level gentamicin resistance in *Enterococcus faecalis*: molecular characteristics and relevance in severe infections. In: *Enterococcus faecalis: molecular characteristics, role in nosocomial infections and antibacterial effects*. Nova Publishers: Hauppauge.
15. Cercenado E. *Enterococcus*: phenotype and genotype resistance and epidemiology in Spain. *Enferm Infect Microbiol Clin*. 2011; 29: 59-65.
16. Chan YY, Abd Nasir MH, Yahaya MA, Salleh NM, Md Dan AD, Musa AM, Ravichandran M. Low prevalence of vancomycin- and bifunctional aminoglycoside-resistant enterococci isolated from poultry farms in Malaysia. *Int J Food Microbiol*. 2008; 122: 221-6.
17. Choi JM, Woo GJ. Molecular characterization of high-level gentamicin-resistant *Enterococcus faecalis* from chicken meat in Korea. *Int J Food Microbiol*. 2013; 165: 1-6.
18. Donabedian SM, Thal LA, Hershberger E, Perri MB, Chow JW, Bartlett P, et al. Molecular characterization of gentamicin-resistant enterococci in the United States: Evidence of spread from animals to humans through food. *J Clin Microbiol*. 2003; 41: 1109-13.
19. Sparo MD, Mallo RA. Evaluation of bacterial flora from a natural maize silage (in Spanish). *Rev Argent Microbiol*. 2001; 33: 75-80.
20. Dutka-Malen S, Evers S, Courvalin P. Detection of glycopeptide resistance genotypes and identification to the species level of clinically relevant enterococci by PCR. *J Clin Microbiol*. 1995; 33: 24-7.
21. Clinical and Laboratory Standards Institute. 2015. Performance standards for antimicrobial susceptibility testing: 25th informational supplement. M100-S25 Wayne: CLSI.
22. Sparo M, Delpech G, Pourcel G, Schell C, de Luca MM, Bernstein J, Grenovero S, Basualdo JA. Cytolysin and high-level gentamicin resistance in *Enterococcus faecalis* from different origin. *RAZYElE*. 2013; 8: 5-10.
23. Sparo MD, Corso A, Galletti P, Delpech G, Ceci M, Confalonieri A, Urbizu L, Sánchez Bruni S. *Enterococcus faecalis* CECT7121: biopreservation of crafted goat cheese. *Int J Prob Preb*. 2012; 7: 145-52.
24. Sparo M, Delpech G, Batistelli S, Basualdo JA. Immunomodulatory properties of cell wall extract from *Enterococcus faecalis* CECT7121. *Braz J Infect Dis*. 2014; 18: 551-5.
25. Delpech G, Bistoletti M, Ceci M, Lissarrague S, Sánchez Bruni S, Sparo M. Bactericidal activity and synergy studies of peptide AP-CECT7121 against multi resistant bacteria isolated from human and animal soft-tissue infections. *Probiotics Antimicrob Proteins*. 2017; 9: 355-62.
26. Gilmore MS, Clewell DB, Ike Y, Shankar N (2014) In: *Enterococci: from commensals to leading causes of drug resistant infection*. Boston: Massachusetts Eye and Ear Infirmary.
27. Obeng AS, Rickard H, Ndi O, Sexton M, Barton M. Comparison of antimicrobial resistance patterns in enterococci from intensive and free range chickens in Australia. *Avian Pathol*. 2013; 42: 45-54.
28. del Campo R, Tenorio C, Rubio C, Castillo J, Torres C, Gómez-Lus R. Aminoglycoside-modifying enzymes in high-level streptomycin and gentamicin resistant *Enterococcus* spp. in Spain. *Int J Antimicrob Agents*. 2000; 15: 221-6.
29. Toledo C, Pérez ME, Rocchi M, Gribaudo G, Mangiaterra S, Monterisi A. Isolation of enterococci species causative of infections and sensitivity to antimicrobial drugs. *Rev Argent Microbiol*. 2004; 36: 31-5.
30. Krocko M, Čanigová M, Ducková V. Occurrence, isolation and antibiotic resistance of *Enterococcus* species isolated from raw pork, beef and poultry. *J Food Nutr Res*. 2007; 46: 91-5.
31. Werner G, Coque TM, Hammerum AM, Hope R, Hryniewicz W, Johnson A, et al. Emergence and spread of vancomycin resistance among enterococci in Europe. *Euro Surveill*. 2008; 13: 19046.
32. Corso AC, Galletti PS, Rodríguez MM, Melano RG, Ceriana PG, Faccone DF, Galas MF, VRE Argentinean Collaborative Group. Molecular epidemiology of vancomycin-resistant *Enterococcus faecium* in Argentina. *Int J Infect Dis*. 2007; 11: 69-75.
33. Cattoir V, Leclercq R. Glycopeptide resistant enterococci. *Med Sci (Paris)*. 2010; 26: 936-42.