



**ORIGINAL RESEARCH PAPER**

**Biochemistry**

**“STUDY OF PLASMODIUM LACTATE DEHYDROGENASE (pLDH) LEVEL AS AN INDIRECT BIOMARKER OF ACUTE PLASMODIUM FALCIPARUM MALARIA INFECTION”**

**KEY WORDS:** P.falciparum, pLDH,

**Dr(Smt.) Gulab Kanwar**

Senior Professor and HOD, Biochemistry Department, Government Medical College, Kota

**Dr Suman Meena\***

3rd year Resident in Biochemistry, Department of biochemistry, Government Medical College, Kota, Rajasthan \*Corresponding Author

**Dr Mohini**

Department of biochemistry, Government Medical College, Kota, Rajasthan

**ABSTRACT**

**Background :** Plasmodium lactate dehydrogenase (pLDH) level was assayed in the sera of 40 adult male and 40 adult female patients within the age group of 12-50 years presenting with acute, Plasmodium falciparum malaria infection and a control group of 40 healthy adults within the same age group.

**Study objective:** Study was conduct to determine diagnostic value of serum LDH level as a biomarker in patient with acute P.falciparum Infection.

**Methods:** Patient selection and pre-qualification were done by simple random sampling of individuals presenting at the New medical college hospital Outpatient Department (OPD) with a history of fever and malaise within a period of one to eight days, and who were confirmed to be infected with the P. falciparum malaria parasite by microscopically examination of Giemsa-stained thin blood slides.

**Results:** The mean serum LDH level in male patients was found to be 562 ±236.0 IU. This level is significantly higher than the control LDH activity of 236.10±19.0 IU (p-value is less than 0.05). The mean serum LDH activity among female patients was 465 ±177.0IU, which is a relatively higher activity compared to the control LDH activity of 236.10 ±19.0 IU (p-value is less than 0.05).

**Conclusion:** The combination of acute hepatocellular injury and red cell haemolysis induced by the invading merozoites may account for the increase in serum LDH activity during this infection. Therefore serum LDH activity is a potentially valuable enzymatic marker of acute uncomplicated P. falciparum malaria infection, especially in the absence of other complicating diseases known to be associated with the above normal serum LDH level.

**I. Introduction**

Malaria has been responsible for the highest mortality in most endemic countries. Even after decades of malaria control campaigns, it persists as a disease of high mortality due to improper diagnosis and rapid evolving drug resistant malaria parasites.

Lactate dehydrogenase (LDH) is an intracellular enzyme, which catalyses the readily reversible reaction involving the oxidation of lactate to pyruvate with nicotinamide adenine dinucleotide (NAD) serving as coenzyme<sup>1</sup>. LDH is an enzyme, which is classified as a true intracellular Enzyme<sup>2</sup> because of its high degree of tissue specificity where overall tissue concentrations are some 500-fold greater than serum levels under normal circumstances<sup>3</sup>. LDH have five theoretically possible forms , which are found in human tissues e.g. liver, heart, erythrocytes, skeletal muscles and kidneys<sup>4</sup>. So disease affecting these organs such as renal infarction myocardial infarction and haemolysis have been reported to be associated with significant elevations in total serum LDH activity. Such elevations have been widely applied as diagnostic indices for kidney, liver, heart and red blood cell dysfunction <sup>5-7</sup>. Additionally, high serum LDH activity has also been reported in small cell carcinoma of the lung , nephroblastoma, neuroblastoma and metabolic neuroendocrine tumour<sup>8</sup> measles and cervical lymphadenitis<sup>9</sup>, Hodgkin's disease and non-Hodgkin's lymphoma, and in the follow-up of ovarian Dysgerminoma<sup>10</sup>.

Plasmodium falciparum malaria infection is a febrile illness accounting for 300-500 million clinical cases annually worldwide. The life cycle of this parasite in the human host includes the developmental cycle in red blood cells, and the cycle taking place in the liver cell parenchyma, includes a series of transformations in the host hepatocytes. Pathophysiological processes usually associated with acute P. falciparum malaria infections, i.e., the hepatic activity of the invading sporozoites leading to centrilobular liver damage and the destruction of the host red blood cells consequent to erythrocytic merogony<sup>11</sup>. Being rich sources of LDH, the acute liver injury and red blood cell destruction will be followed by the release of LDH into the circulation. This finding has important implications because it highlights the potential of using serum LDH activity as an index in the monitoring of acute P.

falciparum malaria infection, particularly when all other possible causes of increased serum LDH levels have been eliminated.

**2. Aims and objective:-**

1. To determine potential of using serum LDH level as a biomarker in the monitoring of acute P.falciparum infection.
2. To determine diagnostic value of serum LDH level in patient with acute P.falciparum infection.

**3. Material and method:-**

Study was conducted at New medical college Hospital Kota during month of July to oct. 2016 when malaria endemicity is at its highest peak because of highest average rainfall.

Patients selection and prequalification were done by simple random sampling of individual admitted in medicine wards at New medical college Hospital Kota. Total 80 patients included in our study among them 40 were males and 40 were females, 40 persons also included as control group.

**4. Exclusion criteria:-**

Patients, whose case history showed a concomitant presentation with the following condition :-

- Acquired immune deficiency syndrome,
- Anemia (chronic anemia due to hypo proliferative, hemolytic and other known hemolytic disorder.)
- Liver cirrhosis,
- Alcoholism,
- Kidney disorder Patients on self-medication with any anti-malarial drugs prior to presentation.
- Malignancy.
- Any bone, muscle and joints diseases.

**5. Methods :-**

Venous blood (5ml) was obtained from each of the patients by venepuncture of the antecubital vein using a sterile needle and syringe between eight and ten o'clock in the morning. The blood samples were then transferred into clean, sterile centrifuge tubes and allowed to clot. Enzyme assay was carried out within 24 hours of collection. Statistically method used was unpaired Student t test

and variable by using Graph in stat version 3.10. was used to calculate p values. P value of <0.05 was considered to be statistically significant and value <0.001 considered as highly significant.(Normal LDH levels range -140 U/L to 280 U/L).

**6. Observation:-**

Total 80 hospitalized patient (male 40 and female 40) within the age group of 12-50 years presenting with acute, uncomplicated *Plasmodium falciparum* malaria infection and a control group of 40 healthy adults within the same age group were included in our study.

**Table I. Serum LDH level in adult male and female *P. falciparum* malaria patients and controls.**

S.N.	SUBJECT	MEAN SERUM LDH ACTIVITY (IU)
1.	Male patients (n-40)	562 ±236.0 IU.
2.	Female patients (n-40)	465 ±177.0 IU,
3.	Controls (n-40)	236.10 ±19.0

**Discussion:-**

In Our study the mean serum LDH level in male patients was found to be 562 ±236.0 IU. This was more than two times above the control LDH level of 236.10 ± 19.0 IU. Similarly, in female patients, the serum LDH level of 465 ± 177.0 IU is over twice of the control serum LDH level. Mean serum LDH level was significantly higher in patient infected with *p. falciparum* infection than control group (p<0.05). Among the patients, the males were found to have a significantly higher serum LDH. Our results were similar to the study by **Garba et al 12** in which mean serum LDH activity in male and female patient were 789.0 and 634.0, respectively. **Maegraith 11** postulated that the factors involved in hepatic dysfunction in acute *P. falciparum* malaria infection involve a synergy between local circulatory failure and centrilobular cellular damage. Since LDH is found in clinically-significant amounts both the liver and red blood cells, the observed increase in serum LDH activity during acute *P. falciparum* malaria infection in this study can be accounted for by a synergy between the two pathophysiological processes usually associated with acute *P. falciparum* malaria infections, i.e., the hepatic activity of the invading sporozoites leading to centrilobular liver damage and the destruction of the host red blood cells consequent to erythrocytic merogony**11**.

Being rich sources of LDH, the acute liver injury and red blood cell destruction will be followed by the release of LDH into the circulation. This finding has important implications because it highlights the potential of using serum LDH level as a biomarker in the monitoring of acute *P. falciparum* malaria infection, particularly when all other possible causes of increased serum LDH levels have been eliminated.

Although, diagnosis of malaria rest on the demonstration of asexual forms of the parasite in stained peripheral blood smear. Sometimes no parasites can be found in peripheral blood smears from patients with malaria, even in severe infections. This may be explained by partial antimalarial treatment or by sequestration of parasitized cells in deep vascular beds**13**. Interpretation of blood smear films require some experience because artifacts are common. Before a thick smear is judged to be negative, 100-200 fields should be examined under oil immersion. So indirect evidences for diagnosis of malaria becomes only the reasons to start or to justify treatment against malaria**14**.

LDH present abundantly in tissues (liver, red blood cells) which get infected by malarial parasite during completion of asexual cycle**15**. So raised serum LDH level may be considered as an evidence for *P. falciparum* infection.

**References:**

[1] Stryer L. Biochemistry. 2nd ed. New York: WH Freeman, 1982.  
 [2] Sullivan JM, Alpers JP. In vitro regulation of rat heart 5'-nucleotidase by adenine nucleotides and magnesium. J Bio Chem 1971; 246:3057-63.  
 [3] Podlasek SJ, McPherson RA. Streptokinase binds lactatedehydrogenase subunit-M, which shares an epitope with plasminogen. Clin Chem 1989; 35:69-73.  
 [4] Calbreath DF. Clinical Chemistry. Philadelphia: WB Saunders, 1992.  
 [5] Wills MR. The Biochemical Consequences of Chronic Renal Failure. New York: Harvey, Miller and Medcalf, 1971.

[6] Timmis AD, Nathan AW. Essentials of Cardiology. Oxford: Blackwell Scientific, 1993.  
 [7] Castaldo G, Oriani G, Cimino L, et al. Total discrimination of peritoneal malignant ascites from cirrhosis- and hepatocarcinoma-associated ascites by assays of ascitic cholesterol and lactatedehydrogenase. Clin Chem 1994; 40:478-83.  
 [8] Kanowski D, Clague A. Increased lactate dehydrogenase isoenzyme-1 in a case of glucagonoma. Clin Chem 1994; 40:158-9.  
 [9] Sugaya N, Kanno J, Nirasawa M, et al. Increased activities of cytosol aminopeptidase and lactate dehydrogenase in serum originate from lymphocytes in necrotizing lymphadenitis. Clin Chem 1990; 36:304-6.  
 [10] Pressley RH, Muntz HG, Falkenberry S, et al. Serum lactic dehydrogenase as a tumour marker in dysgerminoma. Gynecol Oncol 1992; 4:281-3.  
 [11] Maegraith B. Aspects of the pathogenesis of malaria. Southeast Asian Trop Med Pub Health 1981; 12:251-67.  
 [12] I H Garba, G A Ubom. Total serum lactate dehydrogenase activity in acute Plasmodium falciparum malaria infection. Singapore Med J 2005; 46(11): 632  
 [13] Noppadon Tangpukdee, Chatnapa Duangdee, Polrat Wilairatana, Srivicha Krudsood. Malaria Diagnosis: A Brief Review Korean J Parasitol. 2009 Jun;47(2):93-102.  
 [14] available at <http://www.parasitol.or.kr/kjp/search.php?where=aview&id=28586&code=0066KJP&mod>.  
 [15] Nicholus J.White, Joel G.bremen. Malaria. Fauci AS, Braunwald E, kasper DL, longo DL, hauser SL, Jameson JL et al editors—Harrison's Principles of Internal Medicine. Vol 1. 17th ed. New York: Mc Graw Hill Inc, 2008: 1280-94.