



ORIGINAL RESEARCH PAPER

Physiology

EFFECTS OF PRE AND POST EXERCISE EFFECT ON ECG IN FIRST YEAR MEDICAL STUDENTS OF NMC, PATNA

KEY WORDS:

Electrocardiogram, QRS duration, QRS axis, QRS voltage, QTc interval

Manish Ranjan

Tutor, Department of Physiology, Nalanda Medical College, Patna

Anil Kumar Singh*

Professor, Department Of Physiology, Nalanda Medical College, Patna *Corresponding Author

ABSTRACT

Background and Aim: A study was conducted to determine the effect of exercise on Electrocardiogram (ECG) variables in First year students of NMC, Patna, and gender variation of these variables was compared.

Materials and Methods: 100 medical students were considered (55 males and 45 females), ages 18-24 years. The volunteers were subjected to exercise stress test by using bicycle ergometer. They exercised till exhaustion or attainment of 85% Heart Rate Maximum (HRmax) (Modified McArdle Protocol). ECG was recorded pre and post-exercise.

Results: The weight ranges from 45.37 ± 6.53 kg. QRS duration ranged from 0.04 to 0.12 seconds mean of 0.064 seconds. QRS voltage in V6 varied from 0.30 to 2.1 mv with a mean of 1.14 mv. The QTc interval was 0.31 to 0.47 seconds mean was 0.39 seconds.

Conclusion: The study revealed an unexpected finding where PR, QRS and QT durations were prolonged post-exercise in apparently healthy young medical students.

Introduction

Sound health and physical fitness are positively associated with good mental health and well-being. People who have regular physical exercise tend to have less anxiety and depression and lower level of stress than do people with a sedentary lifestyle.¹ It is not only athletes that require physical exercise for better performance but also non-athletes for maintenance of physical and mental health. Buffalo health study concluded that cardiopulmonary function is the long term predictor for overall survival rates in both genders and could be used as a tool in general health assessment.^{2,3}

In our environment, preschool physical examinations do not include ECG for new students and baseline values are scarce. This study was conducted to assess the effect of exercise on ECG parameters in young adults at NMC, Patna. This study aim is to find early signs of cardiovascular disorders among students. Electrocardiograph (ECG) is an important non-invasive and acceptable tool for assessment of myocardial contractility and electrical conduction.⁴ Patterns of ECG vary

from person to person and even within a person, it varies when exposed to different physiological conditions. To the best of our knowledge, no such study has been done among the youths at NMC, Patna. Previous studies of prevalence of ECG findings usually uses subjects 25-75 years or above.⁵ The baseline data can also help in predicting those that may likely develop myocardial dysfunction later in life.⁶

To adequately guide this work, one major hypothesis was formulated, i.e. The null hypothesis is that exercise has no significant effect on cardiac parameters in young healthy subjects.

MATERIALS AND METHODS

The inclusion criterion for participating in this study was being a medical student of University of Abuja. There were 45 preclinical students in the new medical school. All the students volunteered to participate. Hundred(100) students aged 18-24 years qualified to participate after the exclusion criteria were applied.

Exclusion criteria :

Previous history of heart disease; Ingestion of digitalis in preceding six months; Hypertension and associated pulmonary disease (e.g. asthma, pneumonia etc.); Previous history of any surgery; On any medication.

Subjects were advised for relaxation, wear light clothing and take their normal meals 2-3hrs before the study.

Anthropometric measurements were taken. Pre-exercise blood

pressure, pulse rate and respiratory rates were recorded.

A qualified medical personnel supervised the period of exercise test to handle any emergency that may arise. Three technical staffs we also involved in the study. The procedure was demonstrated to the subjects before the exercise.

Recordings were carried out according to the specifications of the American Heart Association, i.e. subjects lying supine, arms by their side, chest electrodes in their correct positions, limb electrodes on the wrists and ankles, recording at 25 mm/sec, calibrated at 10 mm/mV.

Exercise stress test was done using bicycle ergometer. The age-predicted maximal heart rate (HRMax) was first calculated using the formula, HRMax = 220 - age. 75% (sub maximal level) of this value is the heart rate the subject aims to achieve during the stress test.⁸ Subjects sat comfortably on a bicycle with the seat adjusted in such a way that there was knee flexion when the contralateral knee was fully extended. The digital monitor was reset and resistance was fixed at stage 2. The subject started walking slowly and gradually increased the speed, until the attainment of 85% age-predicted HRMax or when exhausted and could not continue due to fatigue (modified McArdle protocol). The subject is asked to walk/run slowly for about 30 seconds. The subject was then asked to sit and relax and the ECG recorded again. The duration

of exercise, distance covered, calories burnt and the maximal heart rate were recorded by the digital monitor. The room temperature at the time of study was 34°C. The study was approved by the ethical committee of Nalanda Medical College, patna. The subjects gave written informed consent for participating in the study. It was according to Helsinki Declaration of June 1964 and amended in Seoul in October 2008.

Data obtained was interpreted using SPSS 13.0. The differences between pre and post-exercise values were evaluated using the pairwise t-test. All descriptive data was expressed as means ± standard deviations. Results were considered significant when p was <0.05.

The descriptive statistics: The ages of the subject ranged from 18-30 years, with average of 21 years among males and 23 years among the females. Their heights were between 1.5-2.0m with average in males being 1.77m and 1.48m in females. The subjects weighed between 44 - 92kg with the average weight in males being 55kg and in females 45kg. BMI of the subjects ranged between 17.72 and 28.57, with males having an average of 21.46 and females 22.52. Duration of QTC interval in males was significantly raised post-exercise but marginal in females (Table-1).

Pre-exercise PR interval was found to be higher in females as compared to male subjects. Post-exercise, the values were higher in males (Table-1). QRS duration was found to be higher in males both pre and post-exercise (Table-1). The ECG variables were

correlated with descriptive variables pre and post exercise. This was to ascertain the strength of relationship between the variables studied. The results are shown in ECG variables of pre- and post-exercise and correlated with descriptive variables in (Table-2).

Table-1: PR Interval, QRS Duration andQTIntervalof subjects pre and post-exercise (ms).

Sex	PR Interval				QRS Duration				QTInterval			
	Pre-Exercise		Post-Exercise		Pre-Exercise		Post-Exercise		Pre-Exercise		Post-Exercise	
Male	143.00	23.9	144.75	14.7	90.86	4.57	74.09	9.81	366.55	20.63	405.82	22.94*
Female	150.44	27.54	123.33	19.21	77.44	5.77	78.89	8.76	389.39	20.41	409.11	18.84
Paired Diff. of Means	-12.94	37.45	-0.13	25.01	3.94	8.32	6.33	13.46	-12.56	33.46	5.39	29.26

Values are mean SD

*Significant (p<0.05).

Table-2: Correlations between PR Interval, QRS Duration and QTInterval and measured variables Pre and Post-exercise.

Variables	PR Interval		QRS Duration		QTInterval	
	Pre	Post	Pre	Post	Pre	Post
Height	- 0.18	0.11	0.39*	0.29	- 0.11	0.08
Weight	- 0.19	0.08	0.19	0.10	- 0.05	0.18
BMI	- 0.04	0.02	- 0.19	-0.20	0.06	0.17

*Significant (p<0.05).

Discussion

In this study, QTC durations fell within the normal range, higher in the females both pre and post-exercise. An increase was recorded post-exercise in both sexes the increase in males was statistically significant (p-value <0.05).

Jonathan and Adams (2002)⁸ found the normal responses during exercise to be shortening of PR, QRS and QT internals. This was corroborated by Monero-Reviriego and Meniro.¹⁰ The obvious response to the increase in HR is shortening of the PR interval, QRS duration and QT interval.

These changes occur in normal subjects and are related to a normal HR response.⁸ But in our study, reverse was the case. It was only the PR intervals in females that showed a post-exercise decrease. Though the changes were not statistically significant except in QTC interval in males where there was significant increase post-exercise. There is thus need for further studies in this region to verify if there is post-exercise increase and look for possible confounding factors that may give these effects.

Conclusion

The paired difference between males and females were not significant in any of the measured variable. Conclusions drawn from the study revealed an unexpected finding where PR, QRS and QT durations were prolonged post-exercise in apparently healthy young medical students. Though they all fell within normal range, ideally these parameters decrease with exercise due to increase in heart rate.

REFERENCES

1. .Pratt M. Physical activity. In: Goldman L, Bennett J .C, editors. Cecil Text book of Medicine. 21st ed. Philadelphia: WB Sanders 2000; 1: 31-3.
2. Schunemann HJ, Dorn J, Grant BJ, Winkelstein WJ, Trevisan M. Pulmonary function is the Long term predictor of mortality in the general population: 29 years follow-up of the buffalo health study. Chest 2000; 118: 656-64.
3. Charles Fisch. Abnormal E.C.G. in normal individuals. J Am Med Assoc. 1984;9:250.
4. Blakman MD, Kuskin MD. Inverted T wave in the precordial E.C.G. of normal adolescents. Am Heart J. 1964;67:304.
6. Prakash S, Meshran S, Rantekkar U. Athletes, yogis and individuals with sedentary life styles; do their lungs function differs? Indian J PhysiolPharmacol 2007; 51: 76-80.
7. Litchfield RL. Non-invasive tests for cardiac risk stratification. Which ones are most prognostic? Postgrad Med 2004; 115: 30-6.
8. Daviglius ML, Liao Y, Greenland P, Dyer AR, Liu K, Xie X, et al. Association of nonspecific minor ST-T abnormalities with cardiovascular mortality: the Chicago Western Electric Study. JAMA 1999; 281: 530-6.
9. Extramiana F, Maison-Blanche P, Badilini F, Pinoteau J, Desco T, Coumel P. Circadian modulation of QT rate dependence in healthy volunteers: gender and age differences. J Electrocardiol 1999; 32: 33-43.

10. Tanaka H, Monahan KD, Seals DR. Age-predicted maximal heart rate revisited. J Am CollCardiol 2001; 37: 153-6.
11. Hill J, Timmis A. ABC of Clinical Electrocardiography- Exercise Tolerance Testing. BMJ 2002; 324: 1084-7.
12. McArdle WD, Katch FI, Katch VL Eds. In:Essentials of exercise Physiology. Philadelphia: Lea and Febger, 1994; pp271.
13. Moreno-Revirego S, Merino JL. Short QT Syndrome. E-Journal of the ECS Council for Cardiology Practice 2010; 9: 2.