



**ORIGINAL RESEARCH PAPER**

**Medical Science**

**A STUDY OF HEMODYNAMICS WHEN DIFFERENT DOSES OF DEXMEDETOMIDINE IS USED WITH PROPOFOL AS AN INDUCING AGENT**

**KEY WORDS:** Hemdynamics, dexmedetomidine, propofol.

**Dr. Tamanna Ahemad**

Post-Graduate, Department of Anesthesiology, A.J.Institute of Medical Sciences, Mangalore.

**ABSTRACT**

Intravenous dexmedetomidine is associated with a biphasic BP response. At low doses the dominant action of  $\alpha_2$  adrenoreceptor agonist activation is a reduction in sympathetic tone, mediated by a reduction of noradrenaline release at the neuroeffector junction, and a inhibition of neurotransmission in sympathetic nerves. The net effect of dexmedetomidine action is a significant reduction in circulating catecholamine's with a slight decrease in BP and a modest reduction in heart rate. When administered as a continuous infusion, it is associated with a stable hemodynamic response. Significant hypotension is usually only observed in patients with pre-existing hypovolemia or vasoconstriction. The bradycardia frequently seen after the administration of dexmedetomidine may be due to the central sympatholytic action, enhanced vagal activity and partly by baroreceptor reflex. Hemodynamic stability provided by  $\alpha_2$  adrenoreceptor agonists in the perioperative period leads to a reduction in perioperative myocardial ischemia.

**Introduction:**

Macewen, a Scottish surgeon, is credited with the first elective use of endotracheal intubation to administer anaesthesia in 1878. He performed an awake, digital, blind intubation under chloroform anaesthesia. Direct laryngoscopy was first performed in 1895 by Alfred Kirstein with a modified esophagoscope called autoscope. In 1913, Chevalier Jackson designed a laryngoscope and was the first to describe tracheal intubation under direct laryngoscopy. Sir Robert Macintosh (1897–1989) introduced the Macintosh curved laryngoscopic blade that has been in use since 1943. In 1940, Reid and Brace first described hemodynamic response to laryngoscopy and intubation. Since then various methods of attenuation of response to laryngoscopy and intubation are still in search. No single method has been established as the most appropriate for this purpose.  $\alpha_2$  adrenoreceptor agonists such as clonidine provide preoperative sedation and anxiolysis. Intraoperative use of  $\alpha_2$  agonists is associated with improved hemodynamic stability, reduced sympathoadrenal responses to laryngoscopy, reduced anaesthetic requirement. They also reduce incidence of postoperative pain and analgesic requisites. Dexmedetomidine, a recently introduced  $\alpha_2$  adrenoreceptor agonists, is being used as a premedication in doses ranging from 0.33 to 1  $\mu\text{g}/\text{kg}$ . It has certain side effects of which most common are bradycardia and hypotension especially in presence of other concomitant factors. Hence this study was conducted to compare and evaluate the effect of different dosages of dexmedetomidine on induction dose of propofol and hemodynamics.

**Aims and Objectives:**

To study the hemodynamics when different doses of dexmedetomidine is used with propofol as an inducing agent.

**Materials and Methods:**

This study was done in the Department of Anesthesia in A.J.Institute of Medical Sciences

This study was done using 60 patients. The study was done from July 2017 to June 2018.

They were divided into 4 groups

- Group A received 1  $\mu\text{g}/\text{kg}$  of dexmedetomidine.
- Group B received 0.6  $\mu\text{g}/\text{kg}$  of dexmedetomidine.
- Group C received 0.3  $\mu\text{g}/\text{kg}$  of dexmedetomidine.
- Group D received 20 ml of normal saline.

**Inclusion Criteria**

1. The patients were aged between 30-50 years
2. The patients had no co-morbidities

**Exclusion Criteria**

1. Aged below 30 and above 50 years

2. Patients with co-morbidities

All the statistics were done using the SPSS software 2015 (California)

**Results:**

**Graph 1: Heart Rate**






Figure -Heart rate

**Graph 2: Mean Blood Pressure**



**Graph 3: Saturation**



Note :Table 1

KEY FOR TIME INTERVALS	
1.	Baseline
2.	10 min post infusion
3.	20 min post infusion
4.	Post induction
5.	Before intubation
6.	1min post intubation
7.	2min post intubation
8.	5 min post intubation
9.	10 min post intubation

**Discussion:**

Significant intragroup ( $p < 0.001$ ) and intergroup ( $p = 0.001$ ) variations in heart rate were noted in the study population. Significant intergroup variations were noted at 10 minutes post infusion ( $p < 0.001$ ), 20 minutes post infusion ( $p < 0.001$ ), post induction ( $p = 0.002$ ), before intubation ( $p = .007$ ), 1 minute post intubation ( $p = 0.001$ ), 2 minutes post intubation ( $p = 0.002$ ) and 5 minutes post intubation ( $p = 0.004$ ).

The study population displayed significant intragroup ( $p < 0.001$ ) and intergroup ( $p = 0.001$ ) differences in mean BP during period of observation. Significant intergroup variations were noted 10 minutes post infusion ( $p = 0.001$ ), 20 minutes post infusion ( $p < 0.001$ ), 1 minute post intubation, 2 minutes post intubation, 5 minutes post intubation and 10 minutes post intubation ( $p < 0.001$ ).

There was no statistically significant difference in oxygen saturation between the four groups. Mean saturation remained above 98% in all four groups.

**Conclusion:**

Attenuation of hemodynamic response was best seen with 1 µg/kg followed by 0.6 µg/kg while hemodynamic profiles of 0.3 µg/kg of dexmedetomidine and placebo group were similar. Hence we conclude that 1 µg/kg and 0.6 µg/kg of dexmedetomidine offer desirable hemodynamics.

**REFERENCES:**

1. Szmuk Peter, Ezri Tiberiu, Evron Shmuel, Roth Jeffrey et al A brief history of tracheostomy and tracheal intubation, from the Bronze Age to the Space Age. *Intensive Care Med.* 2008; 34(2):222–228
2. Gurulingappa, Aleem MA, Awati MN, Adarsh S. Attenuation of Cardiovascular Responses to Direct Laryngoscopy and Intubation-A Comparative Study Between iv Bolus Fentanyl, Lignocaine and Placebo. *J Clin Diagn Res.* 2012 Dec;6(10):1749-52
3. Altan A, Turgut N, Yildiz F, Türkmen A, Ustun H Effects of magnesium sulphate and clonidine on propofol consumption, haemodynamics and postoperative recovery. *Br J Anaesth.* 2005 April; 94(4): 438–441
4. Siobal MS, Kallet RH, Kivett VA, Tang JF. Use of dexmedetomidine to facilitate extubation in surgical intensive-care-unit patients who failed previous weaning attempts following prolonged mechanical ventilation: A pilot study. *Respir Care* 2006;51:492-6.
5. Aantaa R, Kanto J, Scheinin M, Kallio A, Scheinin H. Dexmedetomidine, an  $\alpha_2$ -adrenoceptor agonist, reduces anesthetic requirements for patients undergoing minor gynecologic surgery. *Anesthesiology.* 1990 Aug;73(2):230-5
6. Cheung CW, Ying CL, Chiu WK, Wong GT, Ng KF, Irwin MG. Analgesic and sedative effects of intranasal dexmedetomidine in third molar surgery under local anaesthesia. *Anaesthesia.* 2007 Nov;62(11):1132-8
7. Candiotti KA, Bergese SD, Bokesch PM, Feldman MA, Wisemandle W, Bekker AY et al. Monitored anesthesia care with dexmedetomidine: a prospective, randomized, double-blind, multicenter trial. *Anesth Analg.* 2010 Jan 1;110(1):47-56