



**ORIGINAL RESEARCH PAPER**

**Orthopaedics**

**A CLINICAL STUDY OF THE SURGICAL MANAGEMENT OF SUPRACONDYLAR FEMUR FRACTURE TREATED BY RETROGRADE INTRAMEDULLARY(GSH) INTERLOCKING NAIL**

**KEY WORDS:** Supracondylar fracture,GSH,RETROGRADE NAILING

**Dr.Midhun Jose\***

Orthopaedics Resident , Dept. Of Orthopaedics,kurnool Medicalcollege, kurnool \*Corresponding Author

**Dr. Nagaraju**

Assistant Professor,Dept.of orthopaedics,Kurnool MedicalCollege,Kurnool

**ABSTRACT**

**BACKGROUND-** A prospective randomized study of 20 patients who underwent fixation of the supracondylar femur fractures with GSH nail was done to study the clinical and radiological outcome of GSH interlocking nail in supracondylar femur fractures. The study also evaluated the results of the retrograde nailing in supracondylar femoral fractures in relation to knee flexion, mobilization of patients and early weight bearing.

**MATERIALS AND METHODS**

A study was conducted on 20 patients with Supracondylar fracture femur admitted from OPD Clinic and Casualty clinic of GGH, Kurnool. The study was done over a period of 24 months from May 2018 to May 2020.

**RESULTS**

The duration of follow up ranged from 3 months to 23 months. Out of 20 patients, 1 patient developed knee sepsis, nail was removed immediately and patient was lost for follow-up. 65% good to excellent result were obtained using Neer's and Sander's evaluation scoring system.

**INTRODUCTION**

Supracondylar and intercondylar femoral fractures are often difficult to treat and they are notorious for many complications. The traditional management of displaced fracture supracondylar of femur was along the principle of Watson Jones<sup>1</sup> & John Charnley<sup>2</sup>.

Supracondylar fractures tend to collapse into varus. During application of AO blade plate or dynamic condylar screw, the shaft of femur is often pulled laterally displacing the line of weight bearing, lateral to the anatomic axis of condyle. This creates rotational movements at the fracture site that causes pulling off the blade plate or condylar screws leading to fatigue fracture of the plates. Also, the presence of osteoporotic bone leads to fixation failures with screws and plates cutting of the soft bone.

The obvious advantage of an intramedullary device is that it aligns the

femoral shaft with condyles reducing the tendency to place varus movement at the fracture site. And because the bending movement of an intramedullary device is substantially reduced failure of fixation in osteoporotic bone should be less.

In addition, a retrograde intramedullary supracondylar nail has got distinct advantages of preservation of fracture hematoma, decreased blood loss, minimal soft tissue dissection, less operative time and reduced rate of infection.

**OBJECTIVES OF THE STUDY**

This work has been undertaken to study the clinical and radiological outcome of GSH interlocking nail in supracondylar femur fractures and also to study and evaluate the results of the retrograde nailing in supracondylar femoral fractures.

**MATERIALS AND METHODS**

The study was conducted on 20 patients with Supracondylar fracture femur admitted from OPD Clinic and Casualty clinic of GGH, Kurnool. The study was done over a period of 24 months from May 2018 to May 2020.

**INCLUSION CRITERIA:-**

1. Age between 15-70 years
2. H/O Trauma (RTA, fall from height).

**EXCLUSION CRITERIA:-**

1. Mid shaft femur fractures
2. Non union.
3. Malunion.
4. Pathological fractures

**Implant Used:**

- The implant used was Orthocare supracondylar nail system with instrumentation set.
- The nails are available with outer diameter of 10, 11 and 12 mm
- There is 5 degree anterior bend and an anterior bow for anatomic fit. All sized nails have five interlocking holes in all lengths two proximal holes and three distal holes, which accept interlocking screws of 4.9mm thread diameter.

**RESULTS**

The method used for fracture fixation was closed or open reduction and internal fixation with retrograde intramedullary supracondylar GSH nail. The duration of follow up ranged from 3 months to 23 months. Out of 20 patients, 1 patient developed knee sepsis, nail was removed immediately and patient was lost for follow-up. 65% good to excellent result were obtained using Neer's and Sander's evaluation scoring system.

In this study, the youngest case was 25 years old male and the oldest was 54 years. Overall mean age was 36.15 years.

Seventy five (75%) percent fractures were sustained due to road traffic accidents and fall from height accounted for 25% of fractures. In this study right side affection was seen more than twice as common as left side.

**Table 1 Distribution of Fracture**

Nature of Fracture	Number of cases	Percentage
A1	8	40
A2	10	50
A3	2	10
Total	20	100

Average operative time was 96.50 minutes. 80% of case underwent closed reduction and 20% open reduction. In majority of patients at least 2 screws were used distally.

Average radiological union time was 16.21 weeks of 20 patients, one patient went into deep infection after 2 months.

Average full weight bearing was achieved by 11.68 weeks. Average flexion in this study was 105 degree with more than 50% patients having knee range of motion more than 110 degree. Average extensor lag in this study was 5.68 degrees. Out of 19 patients, 4 had shortening 2 shortening of 22 mm and 2 shortening of 25 mm.

**Table 2:Varus/Valgus Malalignment**

Malalignment of >5 degree	Number of cases	Percentage
Varus	5	25
Valgus	4	20


In this study, very few patients had significant varus/valgus malalignment.

Local symptoms at distal screws was found to be the commonest complications like pain and loosening of screws

**Neer's Rating:**

In 65% cases, there was good to excellent results.

**Graph -1: Neer's and Sander's Rating**



Average knee flexion for closed reduction- 105°

Average knee flexion for open reduction – 101.

The final knee flexion seems to become slightly better if nails with larger diameter were used.

8 patients (62%) out of 13 with 250mm long nail had ≥110 degree knee flexion.

5 out of 6 patients (83%) with 12 mm diameter had ≥ 110 degree flexion.

In one patient with 250 mm long and 11 mm diameter nail developed knee sepsis and the patient was lost for follow-up. There is no correlation between type of reduction and full weight bearing; 69% in closed reduction and 66.66% in open reduction had full weight bearing in 5-12 weeks.

Average weight bearing for closed reduction 11.5 weeks

Average weight bearing for open reduction 12 week

**DISCUSSION**

**Age Incidence:**

Mean age group reported in the previous series were: Lucas SE et al<sup>7</sup>(1993)reported 39 years as mean age group. Watanabe Y<sup>3</sup> (2002) reported 64 years as mean age group. In the present series, the mean age was 36.15 years.

**Sex Incidence:**

Watanabe y<sup>3</sup> (2002) study, whose mean age was 64 years, there were 4 male and 20 female patients.

In the present series, there were 18 male patients with age group 34.89 years, and 2 female with average age of 47.5 years.

**Mechanism of Injury:**

Studies conducted by Schatzker et al<sup>4</sup> (1974), Yang RS et al<sup>5</sup> (1990) and Leung KS et al<sup>6</sup> (1991), demonstrated road traffic accidents as the major causal factor. Lucas SE<sup>7</sup> (1993) reported 79% road traffic accident, 17% fall and 4% gunshot wound.

In the present series, road traffic accidents accounted for 75% of cases and 25% resulting from fall.

**Associated Injuries/ Illness:**

In the present series, 6 patients had associated injuries. Of the 6 injured patients, 2 with ipsilateral proximal tibial fracture had knee flexion of average 82.5° and extensor lag of > 10°. Patient with patellar fracture had knee flexion of 100° and extensor lag of 8°. Two patients with tibial spine fracture had average knee flexion of 102.5° and extensor lag of 9°.

Thus, it appears that though significantly less number of patients in the present series had associated trauma, it seems to affect the final outcome. This can be attributed to delayed mobilization and delayed weight bearing in these patients.

**Injury-Surgery Interval:**

Lucas SE<sup>7</sup>(1993) observed that average injury-surgery interval of 6 days

Watanabe Y<sup>3</sup> (2002) observed the average injury surgery interval of 3 days

In the present series, the injury-surgery interval was 10.3 days

**Operative Time:**

Gyning JB<sup>8</sup> (1999) reported average operative time as 112 minutes. Kumar A<sup>9</sup> (2000) reported average operative time as 58 minutes. Watanabe Y<sup>3</sup> (2002) reported average operative time as 108 minutes. Gellman GE<sup>10</sup> (1996) reported operative time for Type-A 113 minutes. Christodoulou A<sup>11</sup> (2005) reported average operative time of 92 minutes.

Average operative time in this series was 96.5 minutes (75-140 minutes).

**Radiological Union:**

Danziger MB<sup>12</sup> (1995) reported average radiological union time 12.4 weeks.

Gellmann GE<sup>10</sup> (1996) reported average radiological union time of 12 weeks.

Kumar A<sup>9</sup> (2000) reported average radiological union time of 14 weeks.

19 out of 20 cases in the present study united at an average of 16.2 weeks.

**Knee Flexion:**

Kumar A<sup>9</sup>(2000) obtained an average knee flexion of 100 degrees.

Watanabe Y<sup>3</sup> (2002) obtained an average knee flexion of 102 degrees. Ingman AM<sup>13</sup>(2002) obtained an average knee flexion of 101 degrees.

In the present study, the knee flexion was 105 degrees

**Knee Extension Lag:**

Watanabe Y<sup>3</sup>(2002) documented extensor lag of 5 degrees. The average extensor lag in the present series was 5.68 degrees.

**Shortening:**

In Lucas SE<sup>7</sup> (1993) study 1 patient had shortening of >3 cm while Gellmann GE (1996) reported 6 out of 24 patients having 2 cm shortening. In the present study, 4 patients had shortening with an average 2.35 cm and could be compensated by giving a shoe raise.

**Complications:**

**a) Infection:**

There has been no report of infection by any authors except for Lucas SE<sup>7</sup> (1993) who reported one case of septic knee.

In the present series, there was one case of septic knee, 1 month after nail insertion and nail was removed immediately and then patient was lost from followup.

**b) Non-Union:**

Four cases of non-union were reported by Iannacone WM<sup>14</sup> (1994), which were treated with bone grafting and revision fixation.

Kumar A9 (2000) reported one case of non-union but attributed it to technical error, rather than implant. All other studies reported good and solid union.

In the present series, also there was no case of non-union and all patients united radiologically at >3 months interval from surgery.

**c) Delayed Union:**

Danziger MB<sup>12</sup> (1995) also reported a case of delayed union.

In the present series, there was a case of A1-2 type of supracondylar fracture, associated with ipsilateral comminution fracture proximal tibia, immobilized the patient for 10 weeks and showed bridging callus and clinical signs of union at 24 weeks. No secondary procedure was done in that patient except for delayed full weight bearing.

**d) Distal Migration of Nail:**

Gynning JB<sup>16</sup> (1999) reported a case in which the distal locking screws broke at 3 months and the nail protruded in knee joint by 2 cm.

In the present series, there was a case of type-A3-2 fracture in which with weight bearing, the single distal screw cut through the femoral condyles and migrated into surrounding soft tissue. The patient had restricted knee flexion because of mechanical block and painful knee flexion. The implant was removed after the fracture had completely healed

**e) Distal Screw Breakage/ Local Symptoms at Distal Screw:**

Gynning JB<sup>16</sup> (1999) reported no screw breakage but in 5 of his patients, distal screws backed out and were removed under local anesthesia after fracture union. Kumar A9 (2000) reported a case of loosening of distal screw in 1 patient 4 weeks after surgery which was removed without affecting final outcome. In the present series, there were 3 patients who complained of pain at the site of distal screws, which required screw removal after fracture healing.

**f) Stress fracture:**

Kumar A<sup>9</sup> (2000) reported 2 cases of stress fracture at the proximal tip of nail who also had ipsilateral total hip stem. He attributed the fracture to high stress concentration between two intramedullary implants in an osteoporotic bone. In the present series, there was no case of stress fracture.

**g) Implant Failure:**

High implant failure rate was reported by Danziger MB<sup>15</sup>

(1995) and Iannacone<sup>14</sup> (1994) in their studies where nails with multiple holes were used along with 6.4 mm locking screws. With the modification of the nail to total 4 to 5 holes placed at both ends with screw size of 5.0mm, no implant failure has been reported in newer studies. In the present study also, there were no cases of implant failure.

**h) Impingement:**

In the present series 1 nail was kept slightly protruding in the intercondylar notch, due to error in technique. They were removed after solid bone union.

**I) Neer's and Sander's Rating:**

Neer et al<sup>17</sup> (1967) reported 52% satisfactory results with operative method. Janzing et al<sup>18</sup> (1998) used retrograde nailing to treat 26 distal femoral fractures and 72% had neer score of 85 points or more (excellent). In the present series, 65% had Neer score of good to excellent. Also 65% had Sander's score 37 of good to excellent

**CONCLUSION**

1. Retrograde intramedullary supracondylar nail is a good fixation system for distal third femoral fractures, particularly extra-articular type
2. The operative-time is lessened with decrease in blood loss.
3. Closed reduction can be achieved by not disturbing fracture hematoma and soft tissue.
4. Even with open reduction, there is less soft tissue trauma and less postoperative stiffness.
5. Distal screw related local symptoms is a common problem and is related to implant and technique.
6. Utmost great care is required to avoid infection.
7. There is no non-union, less delayed unions and rates of angular or rotational malunions.
8. Non-requirement of bone graft decreases the morbidity associated with donor site.
9. Early surgery, closed reduction, at least two screws in each fragment and early post-operative knee mobilization are essential for good union and good knee range of motion.
10. There is no much difference in individual fracture type healing and weight bearing.

Thus, supracondylar nail is the optimal tool for many supracondylar fractures of femur. It provides rigid fixation in a region of femur, where a widening canal, thin cortices and frequently poor bone stock make fixation difficult. Surgical exposure for nail placement requires significantly less periosteal stripping and soft tissue exposure than that of lateral fixation devices. Orthopaedic surgeons experienced with intramedullary nailing will find the supracondylar GSH nail a useful technique, but requires attention to prevent complications.

**REFERENCES**

1. Wilson JN. Watson Jones's: Fractures and joint injuries. 6th ed, pg. 1003- 070, 1982.
2. Charnley John. The closed treatment of common fractures. 3rd ed, Pg. 197-204
3. Watanabe Y, Takai S, Yamashita F, Kusakabe T. Second generation intramedullary supracondylar nail for distal femur fractures. International ortho (SICOT), 26:85-88, 2002.
4. Schatzker J, Home G, Waddell J. The Toronto experience with supracondylar fractures of femur. Injury, 6, pg. 113-28, 1975.
5. Yang-Rong-Sen, Hwa-Chang Liv et al. Supracondylar fractures of the femur. J Trauma, Vol. 30, pg. 315-19, Mar 1990.
6. Leung KS, Shen WY, Mui LT. Interlocking intramedullary nailing for supracondylar and intercondylar fractures of distal part of femur. JBJS, Vol. 73-A, pg. 332-40, Mar 1991.
7. Lucas. Quoted by Rockwood CA, Green DP. Fractures in adult, 4th ed, Vol. 11, pg. 1972-93, 1996.
9. Kumar A, Jasani VM, Butt MS. Management of distal femoral fractures in elderly patients using retrograde titanium supracondylar nails. Injury, 31(3): 169-73, Apr 2000.
10. Gellman RE, Guy D, Paiement, Hellary D, Green. Treatment of supracondylar femoral fractures with a retrograde intramedullary nail. CORR, No. 332:90-97, 1996.
11. Christodoulou A, Terzidis I, Ploymis A, Metsovitis S, Koukoulidis A, Toptsis C. Supracondylar femoral fractures in elderly patients treated with the dynamic

- condylar screw and the retrograde intramedullary nail: A comparative study of the two methods. *Arch Orthop Trauma Surg*, 125(2):73-9, March 2005. Epub 2004 Dec. 21.
12. Danziger MB, Louci D, Zecher SB. Treatment of intercondylar and supracondylar distal femur fractures using the GSH supracondylar nail. *Am J Orthop*, 24 (9):684-90, 1995.
  13. Ingman AM. Retrograde intramedullary nailing of supracondylar femoral fractures: Design & Development of a New Implant. *Injury*, 33(8);707- 12: 2002 Oct.
  14. Iannacone WM, Bennett FS, DeLong WG Initial experience with the treatment of supracondylar femoral fractures using the supracondylar nail: A Preliminary Report. *J OrthoTrauma*, 8(4):322-7, Aug 1994.
  16. Gynning JB, Hanson D. Treatment of distal femoral fractures with intramedullary supracondylar nails in elderly patients. *Injury*, 30(1):43- 46, Jan 1999.
  17. Neer CS, Gratham SA, Shelton ML et al Supracondylar fractures of adult femur. *JBJS*, Vol. 49-A, pg. 591-613, 1967.
  18. Janzing HM, Vaas F, Van-Damme G et al. Treatment of distal femoral fractures in the elderly: Results with the retrograde intramedullary supracondylar nail. *Unfallchir*, 1998;24: 55.