



ORIGINAL RESEARCH PAPER

Ayurveda

GLYCYRRHIZA GLABRA (LIQUORICE): ASTOUNDING MEDICAL BENEFITS

KEY WORDS: Glycyrrhiza, Anti-bacterial, tuberculosis

Danish K Chishti*

Research Associate Unani, RRIUM, Ministry of Ayush, Delhi. *Corresponding Author

ABSTRACT

At the point when the Greek civilisation was invaded by the Romans, an incredible number of books were lost. Luckily this treatise on *Materia Medica* (in an Arabic interpretation) persevered in the library of the House of Wisdom, an incredible assortment of logical and medical information in Baghdad. Specifically the popular doctor Ibn Baithar utilized the book widely in the thirteenth century in his compositions on therapeutics. This information was spread broadly through the Islamic caliphate, as far west as Spain and as far north to the zone presently known as the Balkans. Licorice (*Glycyrrhiza glabra*) has for quite some time been utilized for both culinary and clinical purposes. It is generally utilized in bronchial issues, for example, catarrh, bronchitis, cold, influenza and hacks. It diminishes aggravation of the throat but then has an expectorant activity. It delivers its demulcent and expectorant impacts. It is utilized in easing pressure. It is a powerful recuperating specialist for tuberculosis, where its belongings have been contrasted with hydrocortisone. *Glycyrrhiza* is likewise viable in assisting with lessening fevers (*glycyrrhetic* corrosive has an impact like anti-inflammatory medicine), and it might have an antibacterial activity too. Its utilization in the treatment of interminable irritations, for example, joint pain and rheumatic illnesses, Anti-provocative, constant skin conditions, and immune system maladies when all is said in done.

INTRODUCTION

Glycyrrhiza glabra, otherwise called licorice and sweetwood, is local to the Mediterranean and certain zones of Asia. Verifiably, the dried rhizome and foundation of this plant were utilized restoratively by the Egyptian, Chinese, Greek, Indian, and Roman civic establishments as an expectorant and carminative. In current medicine, licorice extricates are frequently utilized as a seasoning specialist to veil severe preference for arrangements, and as an expectorant in hack and cold arrangements. Licorice extricates have been utilized for over 60 years in Japan to treat incessant hepatitis, and furthermore have restorative advantage against different infections, including human immunodeficiency infection (HIV), cytomegalovirus (CMV), and Herpes simplex.¹ Deglycyrrhizinized licorice (DGL) arrangements are valuable in treating different sorts of ulcers, while topical licorice arrangements have been utilized to sooth and recuperate skin ejections, for example, psoriasis and herpetic sores.

Pharmacognosy

Glycyrrhiza glabra Linn is a tough lasting shrub, achieving a height up to 2.5 m. The leaves are compound, imparipinnate, substitute, having 4-7 sets of oval, circular or lanceolate leaflets. The blossoms are narrow, typically papilionaceous, borne in axillary spikes, lavender to violet in shading. The calyx is short, campanulate, with lanceolate tips and bearing glandular hairs. The organic product is a compacted vegetable or unit, up to 1.5 cm long, erect, glabrous, fairly reticulately hollowed, and as a rule contains 3-5 dark colored, reniform seeds. The taproot is around 1.5 cm long and subdivides into 3-5 backup roots, about 1.25 cm long, from which the level woody stolons emerge. These may arrive at 8 m and when dried and cut, together with the root, comprise business liquorice. It might be discovered stripped or unpeeled. The bits of root break with a stringy crack, uncovering the yellowish inside with a trademark scent and sweet taste.^{1,2}

Vernacular Names

Sanskrit: Yashti, madhuh, Madhuka
 Bengali: Jashtimadhu, Jaishbomodhu
 Gujarat: Jethimadhu
 Hindi: Jothi, madh, Mulhatti
 Kannada: Yastimadhuka, atimaddhura
 Oriya: Jatimadhu
 Tamil: Atimaduram
 English: Licorice, Liquorice, Sweet wood

Arab: Aslussiesa
 Persia: Ausareha mahaka
 France: Boisdoux

Description

The licorice shrub is an individual from the pea family and develops in subtropical atmospheres in rich soil to a stature of four or five feet. It has oval handouts, white to purplish blossom bunches, and level cases. Subterranean, the licorice plant has a broad root framework with a principle taproot and various sprinters. The fundamental taproot, which is collected for restorative use, is delicate, stringy, and has a splendid yellow interior.¹ *Glycyrrhiza* is gotten from the antiquated Greek term glykos, which means sweet, and rhiza, which means root.

Geographical Distribution

Native to central and south western Asia and the Mediterranean region. It is cultivated in the Mediterranean basin of Africa, in South Europe and in India.

Genetics

Glycyrrhiza glabra is a diploid with 2n=16.

Habitat

Glycyrrhiza glabra is a hard herb or under shrub accomplishing a tallness up to 6ft; leaves multifoliolate, imparipinnate, blossoms in axillary spikes, papilionaceous, lavender to violet in shading, cases compacted, and containing reniform seeds. The dried, stripped or unstripped underground stems and roots establish the medication, referred to in the exchange as Liquorice. Blossoms in March and natural products in August.

Active Constituents

Various parts have been detached from licorice, including a water-dissolvable, naturally dynamic complex that represents 40-50 percent of complete dry material weight. This complex is made out of triterpene saponins, flavonoids, polysaccharides, gelatins, basic sugars, amino acids, mineral salts, and different other substances.² *Glycyrrhizin*, a triterpenoid compound, represents the sweet taste of licorice root. This compound speaks to a blend of potassium-calcium-magnesium salts of *glycyrrhetic* corrosive that differs inside a 2-25 percent extend. Among the common saponins, *glycyrrhetic* corrosive is an atom made out of a hydrophilic section, two particles of *glucuronic* corrosive, and a hydrophobic piece, *glycyrrhetic* acid.² The yellow shade of

licorice is because of the flavonoid substance of the plant, which incorporates liquiritin, isoliquiritin (a chalcone), and other compounds.³ The isoflavones glabridin and hispaglabridins An and B have huge cell reinforcement activity,⁴ and both glabridin and glabrene have estrogen-like activity.⁵

Part Used

Root and Rhizomes

Phyto Chemistry

The major bio.active constituent of rhizomes is a triterpenoids saponin glycyrrhizin, glycyrrhizinic acid, glabrin A&B, glycyrrhetol, glabrolide, isoglabrolide, isoflavones, coumarins, triterpene sterols etc.

Physical Properties

- Total ash Not more than 7%
- Acid insoluble ash Not more than 2%
- Sulfated ash Not more than 10%
- Water soluble extractive Not less than 20%
- Diluted alcohol-soluble
- Extractive Not less than 25%
- Moisture 5.25%
- Ether extracts 16.85%
- Albuminoids 37.00 % (containing nitrogen 5.92%)
- Soluble carbohydrates 31.00%
- Woody fiber 5.05%
- Ash 4.80 % (containing sand 0.25%)

Substitutes Or Adulterants

Manchurian licorice is gotten from glycyrrhiza uralensis. Being a substitute it portion contain glycyrrhizin the dynamic standard yet almost no of free sugars. The normal adulterant is wild licorice likewise called Indian licorice, got from the underlying foundations of Abrus precatorious (leguminosae). Minutely the adulterant is described by stone cells.

Clinical Studies Done On Glycrrriza Glabra Chronic Hepatitis

In Japan, glycyrrhizin has been utilized for over 60 years as a treatment for interminable hepatitis C. Stronger Neo-Minophagen C (SNMC), a glycyrrhizin arrangement, has been widely utilized with significant achievement. In two clinical preliminaries, SNMC has been appeared to altogether bring down aspartate transaminase (AST), alanine transaminase (ALT), and gamma-glutamyltransferase (GGT) fixations, while all the while improving histologic proof of putrefaction and provocative injuries in the liver.^{6,7}

Oral Lichen Planus

Patients with chronic hepatitis C regularly experience oral lichen planus, a provocative infection described by lymphocytic hyperkeratosis of the oral mucosa. It is seldom relieved and compelling medicines are constrained. In an open clinical preliminary, 17 hepatitis C-positive patients with oral lichen planus were given either routine dental consideration or 40 mL IV glycyrrhizin day by day for one month. Among nine patients taking glycyrrhizin, six (66.7%) noted improved clinical side effects, for example, diminished redness, less white papules, and less disintegration of the mucosa. In the non-glycyrrhizin gathering of eight patients, just one (14.3%) announced any improvement.⁸

Peptic Ulcer Disease

Licorice has been utilized as a demulcent and emollient for a long time to advance the recuperating of ulcers by following up on the mucosal layer. Glycyrrhizin (as carbenoxolone sodium) speeds mending of gastric ulcers and ensures against ibuprofen prompted harm to the gastric mucosa. In a twofold visually impaired, fake treatment controlled examination, 70 patients with endoscopically-affirmed gastric or duodenal ulcers were given carbenoxolone sodium

300 mg or fake treatment day by day during the initial seven days, trailed by 150 mg day by day throughout the following 3-5 weeks. The creators finished up the carbenoxolone bunch had an expansion in pH at the stomach antrum from 1.1 to 6.0, and a decrease in basal and histamine-actuated gastric corrosive emission at pH 3 and 5. In general, 70 percent of ulcers in the glycyrrhizin bunch mended inside 3-5 weeks of starting treatment, contrasted with 36 percent utilizing placebo.⁹

Aphthous Ulcers

In a double-blind, placebo treatment controlled preliminary, 24 patients with repetitive aphthous ulcers were haphazardly dispensed to devour 2 g glycyrrhizin (carbenoxolone sodium) in 30 mL of warm water or a fake treatment multiple times day by day following suppers for about a month. Rather than the placebo treatment gathering, the utilization of the oral licorice mouthwash fundamentally diminished the normal number of ulcers every day, torment scores, and the advancement of new ulcers.¹⁰

Anti-bacterial Activity

Shirazi M.H. et al. had contemplated the in vitro inhibitory impacts of *G. glabra* extract against the development of *Salmonella typhi*, *S. paratyphi B*, *Shigella sonnei*, *S. flexneri* and enterotoxigenic *E. coli* (ETEC *E. coli*) was examined utilizing admirably and circle dissemination technique. *Salmonella paratyphi B* demonstrated no powerlessness to liquorice with focuses lower than 7.5%, anyway totally tried bacterial strains showed defenselessness to high grouping of liquorice.^{11,12}

Immunological Effect

Neutrophils treated with alcoholic extract of *Glycyrrhiza glabra* showed increase in phagocytic activity.¹³

Medicinal Uses Of Glycyrrhiza Glabra (licorice)

It has anti-inflammatory, antiviral, saponin glycyrrhizin and numerous significant synthetic substances that are required for acceptable wellbeing. It is principally utilized in treating cervical malignancy, kidney and bladder issue, HIV, hepatitis B, herpes, aggravation of mucous film, nourishment poison, stomachaches, hacks, horse-voice, bronchitis, asthma, ulcers, joint inflammation, shingles, sun-consumes, fevers in newborn children, and furthermore utilized for creepy crawly nibbles. Licorice glycyrrhizic corrosive is multiple times better than sucrose.⁹

It is said to empower the creation of hormones, for example, hydrocortisone, bolster the adrenal organs and animates the discharge of hormones from the adrenal cortex. Licorice has estrogen movement, used to settle menstrual cycle, and furthermore know generally as a decent solution for lungs and spleen protests. It is joined with different herbs in treating hacks, colds, sore throat, asthma, stomach and duodenal ulcers, hepatitis, madness, food contamination, hypoglycemia, bronchitis, colitis, diverticulitis, gastritis, some of stress related scatters, sickness, and irritation. Licorice purifies the colon, advances adrenal organ work, diminishes muscle fits, and builds the smoothness of bodily fluid from the lungs and bronchial tubes.^{8,9,10}

Licorice properties incorporate anodyne, cell reinforcement, antispasmodic, mitigating, demulcent, depurative, diuretic, cooling, expectorant, emollient, estrogenic, expectorant, hostile to ligament, tonic, energizer adrenal, against cholesterolemic, against gastritis and against allergenic. The Licorice glycoside is like the normal steroids of the body and has a decent impact in treatment of Addison's illness.¹⁰

Dosage

Because individual susceptibility to various licorice preparations is vast, it is difficult to predict a dose appropriate

for all individuals. Nevertheless, a daily oral intake of 1-10 mg of glycyrrhizin, which corresponds to 1-5 g licorice (2% glycyrrhizin), has been estimated to be a safe dose for most healthy adults.¹¹⁻¹²

CONCLUSION

Licorice (*Glycyrrhiza glabra* Linn) root, rhizome and its concentrate, for example, glycyrrhizin have a long history of utilization in traditional system of medicine, people cures, and as a sweetening and seasoning operator. Pharmacological investigations have assessed a few of the conventional wellbeing claims behind licorice use albeit a considerable lot of these reports have created conflicting outcomes. Deglycyrrhizinated licorice has likewise given some impact in the treatment of gastrointestinal ulcers, recommending the nearness of dynamic fixings other than glycyrrhizin, and many more studies have demonstrated it has many gainful impacts. The review highlighted the chemical constituent, pharmacological and therapeutic effects of *Glycyrrhiza glabra* as promising source of drugs because of its safety and effectiveness.

REFERENCES:

1. Olukoga A, Donaldson D. Historical perspectives on health. The history of liquorice: the plant, its extract, cultivation, and commercialisation and etymology. *JRSoc Health* 1998;118:300-304.
2. Obolentseva GV, Litvinenko VI, Ammosov AS, et al. Pharmacological and therapeutic properties of licorice preparations (a review). *Pharm Chem J* 1999;33:24-31.
3. Yamamura Y, Kawakami J, Santa T, et al. Pharmacokinetic profile of glycyrrhizin in healthy volunteers by a new high-performance liquid chromatographic method. *J Pharm Sci* 1992;81:1042-1046.
4. Vaya J, Belinky PA, Aviram M. Antioxidant constituents from licorice roots: isolation, structure elucidation and antioxidative capacity toward LDL oxidation. *Free Radic Biol Med* 1997;23:302-313.
5. Tamir S, Eizenberg M, Somjen D, et al. Estrogenlike activity of glabrene and other constituents isolated from licorice root. *J Steroid Biochem Mol Biol* 2001;78:291-298.
6. Van Rossum TG, Vulto AG, Hop WC, Schalm SW. Glycyrrhizin-induced reduction of ALT in European patients with chronic hepatitis C. *Am J Gastroenterol* 2001;96:2432-2437.
7. Tsubota A, Kumada H, Arase Y, et al. Combined ursodeoxycholic acid and glycyrrhizin therapy for chronic hepatitis C virus infection: a randomized controlled trial in 170 patients. *Eur J Gastroenterol Hepatol* 2008;11:1077-1083.
8. Da Nagao Y, Sata M, Suzuki H, et al. Effectiveness of glycyrrhizin for oral lichen planus in patients with chronic HCV infection. *J Gastroenterol* 1999;31:691-695.
9. Loginov AS, Speransky MD, Speranskaya IE, et al. The effectiveness of carbenoxolone in the treatment of gastro-duodenal ulcer patients. *Scand J Gastroenterol Suppl* 1980;65:85-91.
10. Poswillo D, Partridge M. Management of recurrent aphthous ulcers. *Br Dent J* 1984;157:55-57.
11. Walker BR, Edwards CR. Licorice-induced hypertension and syndromes of apparent mineralocorticoid excess. *Endocrinol Metab Clin North Am* 2004;23:359-377.
12. Shirazi MH, Ranjbar R, Eshraghi S, Sadeghi G, Jonaidi N, Bazzaz N, Izadi M, Sadeghifard N., An Evaluation of Antibacterial Activity of *Glycyrrhiza glabra* Linn Extract on the Growth of *Salmonella*, *Shigella* and *ETEC E. coli*, *J Biological Sciences*, 2007; 7(5):827-829
13. Al-Snafi AE. Medicinal plants possessed antioxidant and free radical scavenging effects (part 3)- A review. *IOSR Journal of Pharmacy* 2017;7(4):48-62