



ORIGINAL RESEARCH PAPER

Anesthesiology

A COMPARATIVE STUDY BETWEEN INTRAVENOUS DEXMEDETOMIDINE AND 50% MAGNESIUM SULFATE IN ATTENUATION OF CARDIOVASCULAR RESPONSE TO LARYNGOSCOPY AND ENDOTRACHEAL INTUBATION.

KEY WORDS: laryngoscopy, dexmedetomidine, magnesium sulphate, hemodynamic changes

Dr Deba Gopal Pathak

Prof(Head), Dept of Anaesthesiology And Critical Care, Silchar Medical College And Hospital, Silchar, Assam

Dr Puja Choubey*

Post Graduate Trainee, Dept of Anaesthesiology And Critical Care, Silchar Medical College And Hospital, Silchar, Assam *Corresponding Author

ABSTRACT

INTRODUCTION: Hypertension and tachycardia subsequent to tracheal intubation have been well documented. In susceptible patients it can cause increase in perioperative morbidity and mortality. Many agents have been used to attenuate these undesirable hemodynamic responses to laryngoscopy and intubation with varying success. Dexmedetomidine increases the hemodynamic stability by altering the stress induced sympatho-adrenal responses to laryngoscopy and intubation during surgery and emergence from anesthesia. On the other hand, magnesium sulfate can significantly attenuate the release of catecholamines at the time of laryngoscopy and tracheal intubation.

AIM: The present study was aimed to compare Inj, Dexmedetomidine 1µg/kg and Magnesium sulfate 50mg/kg in attenuation of cardiovascular response to laryngoscopy and intubation.

METHODS: A randomized prospective study was conducted in Silchar Medical College and Hospital, Silchar, Assam after obtaining Institutional Ethical Committee clearance and written informed consent from the patients.

INCLUSION CRITERIA: Patients between age 18-60 years, of both sexes, ASA grade I&II and scheduled for elective surgical procedure under general anesthesia

EXCLUSION CRITERIA: Refusal to informed consent, anticipated difficult airway, ASA grade III&IV, history of allergy to study drug, pregnant and lactating mother, any disorders of cardiovascular system, respiratory system, renal system, hepatic and neuromuscular conditions and duration of surgery less than 60mins. 100 patients were randomly allocated to two groups of 50 patients each. Group A received inj. dexmedetomidine (1µg/kg) and Group B received inj. Magnesium sulfate (50 mg/kg). Hemodynamic parameters of patients (HR, SBP, DBP, and MAP) were recorded immediately before anesthesia induction, just before intubation, just after intubation, at 2min, 5min and 10 minutes for the study purpose. All analyses were done by using two tailed test; p-value <0.05 was considered significant and <0.001 as highly significant.

RESULTS: There was no significant difference between the groups with regard to demographic variables. The mean fall in HR, SBP, DBP and MAP values at 2 and 5 minutes of intubation were statistically highly significant (p=0.000) in group A as compared to group B.

DISCUSSION: The results were compared with various other similar studies which had analyzed the effect of dexmedetomidine and magnesium sulfate for attenuation of cardiovascular response during laryngoscopy and intubation and our results correlated with those studies.

INTRODUCTION

It is well proven that direct laryngoscopy and intubation are very noxious stimuli and induce stress response in the form of laryngo-sympathetic stimulation and hemodynamic responses. These responses are well tolerated in an otherwise healthy individual, but in patients suffering from hypertension, coronary artery disease, cerebrovascular disease, intracranial aneurysms etc, these transient hemodynamic changes can result in potentially deleterious effects like left ventricular failure, pulmonary edema, myocardial ischemia, ventricular dysrhythmias and cerebral hemorrhage.¹

These changes are maximum in the first 1 minute and last for 10 minutes after the procedure. To attenuate these deleterious effects various methods and drugs have been tried⁽²⁻¹¹⁾.

Dexmedetomidine is a α2 receptor selective and specific adrenergic agonist.^{12,13} These drugs by virtue of their sympatholytic (i.e. antihypertensive and negative chronotropic) action, attenuate the hemodynamic response following laryngoscopy and endotracheal intubation⁸.

Magnesium is well known to block the release of catecholamines from both adrenergic nerve terminals and the adrenal gland, and intravenous magnesium sulfate inhibits catecholamine release associated with laryngoscopy. Moreover, magnesium produces vasodilator effect by acting directly on blood vessels, and high-dose magnesium attenuates vasopressin-stimulated vasoconstriction^{14,15}

The present study was undertaken to compare the efficacy of

magnesium sulphate and dexmedetomidine for attenuation of haemodynamic responses following laryngoscopy and endotracheal intubation.

METHODS

The study was undertaken after obtaining Institutional Ethical Committee clearance and written informed consent from the patients. A randomized prospective study involving 100 patients of both sexes between 18-60 years, requiring endotracheal intubation and general anaesthesia for various elective surgical procedures belonging to ASA grade I and II were included in the study.

The study population was divided into 2 groups with 50 patients in each group.

Patient with anticipated difficult airway, ASA grade III&IV, History of allergy to study drug, pregnant and lactating mother, patient with any disorders of cardiovascular system, respiratory system, renal system, hepatic and neuromuscular conditions, duration of surgery less than 60mins and refusing consent were excluded from the study.

On the day prior to surgery a thorough clinical examination of the patient was performed. All patients were explained about the anaesthetic technique and written informed consent was taken. Patients were kept NPO for 8 hours prior surgery. All patients were given tablet alprazolam 0.5 mg orally at bed time on the previous night of the surgery.

On arrival of the patient in the operating room, a 20-gauge intravenous cannula was secured and connected to IV fluid

ringer lactate. The baseline systolic, diastolic blood pressure, mean arterial pressure and heart rate was recorded after 5mins of settling in the operative room. All patients were being premedicated with intravenous (IV) glycopyrrolate (0.05mg/kg) midazolam 0.03mg/kg, IV fentanyl (2µg/kg). Patients in group A received inj. Dexmedetomidine 1µg/kg body weight diluted in 10 ml normal saline intravenously over 10 min and patients in group B received inj. Magnesium sulfate 50 mg/kg diluted to 10 ml with normal saline over 10 minutes.

The patients were then pre oxygenated for 3 minutes, anesthesia was induced with propofol (titrated till loss of verbal response). Endotracheal intubation was facilitated with IV Atracurium 0.5mg/kg three minutes prior to laryngoscopy and intubation. If time for laryngoscopy and intubation exceeded 15 seconds, such patients were excluded from the study. Anesthesia was maintained using 60% nitrous oxide and 40% of oxygen with isoflurane and inj. Atracurium.

Hemodynamic parameters of patients including systolic BP (SBP), diastolic BP (DBP), mean arterial pressure (MAP) and heart rate (HR) were recorded immediately before anesthesia induction, just before intubation, just after intubation, 2min, 5min, 10min, and every 15 mins after intubation till the operation ended as the hemodynamic changes and intubation mostly occurs in the first minute and last for approximately 10 mins after the procedure. At the end of the surgical procedure patients was reversed with neostigmine 0.05 mg/kg body weight and glycopyrrolate 0.02mg/kg.

STATISTICAL METHOD EMPLOYED

All data are presented as Mean ± SD (Standard Deviation). All Quantitative data are assessed using Student's t - test to analyse changes over a period of time. Qualitative data are assessed using Fisher exact Test or Chi- square test. P value < 0 .05 is considered significant. The statistical software **GRAPHPAD INSAT-3** was used for the analysis of data.

RESULTS:

The mean age, weight, height and duration of surgery of both the groups were comparable. There was no significant difference amongst the groups with regard to demographic variables (P value < 0.05).

CHANGES IN HEART RATE :

Table 1:- Comparison of mean heart rate between two groups

	Heart Rate (beats/min)				P- value
	Groups A (DEXMED) N=50		Groups B (MgSO4) N=50		
	Mean	SD	Mean	SD	
HR T0 (Baseline)	85.36	5.55	86.56	7.07	0.348
HR T (Before Induction)	84.82	5.72	85.46	6.55	0.604
HR T1 (Before Intubation)	74.68	6.45	77.52	6.95	0.037
HR T2 (Just After Intubation)	83.02	5.45	96.14	4.48	0.000
HR T3 (2mins After Intubation)	77.06	5.37	90.46	3.79	0.000
HR T4 (5mins After Intubation)	73.44	5.55	84.52	3.41	0.000
HR T5 (10mins After Intubation)	84.90	5.32	85.90	8.72	0.490

The mean HR decrease observed just after intubation, at 2 and 5 minutes after intubation in Group A was statistically highly significant compared to mean HR in group B (p=0.000).




Figure 1 – Line diagram showing comparison of mean heart rate between the two groups.

CHANGE IN SYSTOLIC BLOOD PRESURE

Table 2:- Comparison of mean systolic blood pressure in between two groups

	SBP				P- value
	Groups A (DEXMED) N=50		Groups B (MgSO4) N=50		
	Mean	SD	Mean	SD	
SBP(mmHg) T0 (Baseline)	127.68	7.05	129.96	6.76	0.102
SBP(mmHg) T (Before Induction)	126.62	7.58	128.64	7.72	0.190
SBP(mmHg) T1 (Before Intubation)	103.04	7.30	105.90	5.51	0.029
SBP(mmHg) T2 (Just After Intubation)	116.18	7.45	136.46	8.65	0.000
SBP(mmHg) T3 (2mins After Intubation)	108.28	5.73	128.38	7.08	0.000
SBP(mmHg) T4 (5mins After Intubation)	102.44	5.42	120.88	7.75	0.000
SBP(mmHg) T5 (10mins After Intubation)	126.68	7.04	127.36	8.77	0.670

The mean fall in SBP in group A just after intubation, at 2 and 5 minutes after intubation was statistically highly significant (p=0.000) compared to group B.




Figure 2– Line diagram showing comparison of mean systolic blood pressure between two groups.

DIASTOLIC BLOOD PRESSURE

Table 3:- Comparison of mean diastolic blood pressure between the two groups.

	DBP				P- value
	Groups A (DEXMED) N=50		Groups B (MgSO4) N=50		
	Mean	SD	Mean	SD	
DBP(mmHg) T0 (Baseline)	82.02	4.22	82.36	4.29	0.690
DBP(mmHg) T (Before Induction)	81.12	4.25	81.56	4.27	0.607
DBP(mmHg) T1 (Before Intubation)	72.12	5.36	73.94	3.80	0.053
DBP(mmHg) T2 (Just After Intubation)	68.62	6.06	69.96	5.23	0.240
DBP(mmHg) T3 (2mins After Intubation)	69.24	3.92	79.94	3.48	0.000

DBP(mmHg) T4 (5mins After Intubation)	66.34	3.85	75.94	3.38	0.000
DBP(mmHg) T5 (10mins After Intubation)	81.60	3.86	82.30	5.32	0.453

The fall in mean DBP values at 2 and, 5 minutes of intubation were statistically highly significant (p=0.000) in group A compared to group B.




Figure 3 – Line diagram showing comparison of mean diastolic blood pressure between the two groups.

MEAN ARTERIAL PRESSURE

Table 4:- Comparison of mean arterial pressure between the two groups.

	MAP				P-value
	Groups A (DEXMED) N=50		Groups B (MgSO4) N=50		
	Mean	SD	Mean	SD	
MAP(mmHg) T0 (Baseline)	96.98	4.31	97.86	3.99	0.292
MAP(mmHg) T (Before Induction)	95.98	4.47	97.02	3.81	0.214
MAP(mmHg) T1 (Before Intubation)	82.38	4.96	83.94	4.56	0.105
MAP(mmHg) T2 (Just After Intubation)	80.02	4.88	81.88	4.60	0.053
MAP(mmHg) T3 (2mins After Intubation)	81.98	3.62	95.50	3.62	0.000
MAP(mmHg) T4 (5mins After Intubation)	77.96	3.54	90.46	3.45	0.000
MAP(mmHg) T5 (10mins After Intubation)	96.04	4.25	96.84	5.48	0.417

The fall in mean MAP values in group A at 2 and 5 minutes of intubation were statistically highly significant (p=0.000) compared to group B.




Figure 4 – Line diagram showing comparison of MAP between two groups.

In our study, we noticed 6% incidence of PONV and bradycardia and 4% incidence of hypotension in group A. The incidence of hypotension and bradycardia was less with MgSO4 but the difference between the two groups was non-significant. Tachycardia was seen in Group B which were not significant when compared with group A. These side effects didn't require any intervention.

DISCUSSION.

Laryngoscopy and intubation is a routine practice in

anesthesiology. King et al., (1951)¹⁶ have described the circulatory responses to laryngeal and tracheal stimulation following laryngoscopy and tracheal intubation as reflex sympathoadrenal stimulation. Many agents have been used to attenuate undesirable haemodynamic responses to laryngoscopy and intubation with varying success. These include intravenous opioids, vasodilators, calcium channel blockers, intravenous and topical lignocaine and adrenoceptor blocking drugs alone or in combination with other drugs²⁻¹¹.

The present study aimed at comparing intravenous dexmedetomidine and 50% Magnesium sulfate for the hemodynamic attenuation during laryngoscopy and endotracheal intubation.

The study population consisted of 100 patients divided equally in two groups. Patients in group A received inj. Dexmedetomidine 1µg/kg body weight diluted in 10 ml normal saline intravenously over 10 min and patients in group B received inj. Magnesium sulfate 50 mg/kg diluted to 10 ml with normal saline over 10 minutes. Attempts have been made to suppress the circulatory response using various pharmacological agents aimed at afferent, efferent or both limbs of response.

Dexmedetomidine, a highly selective α2-adrenoreceptors agonist, increases the hemodynamic stability by altering the stress induced sympatho-adrenal responses to laryngoscopy and intubation during surgery and emergence from anesthesia. Dexmedetomidine by activating pre and post-synaptic α2-receptors of sympathetic system produces vasodilatation; also by acting on post-synaptic α2-receptors of vascular smooth muscle cells it produces vasoconstriction. It there by shows a biphasic, dose dependent response on blood pressure and heart rate, characterized by an initial short-term increase in BP followed by a longer lasting reduction in BP and HR^{17,18,19,20}.

Magnesium sulfate, on the other hand, has been described as the physiological calcium antagonist²¹ because it competes with calcium for membrane channels and can modify many calcium-mediated responses. The ability of magnesium ions to inhibit the release of catecholamine's from both the adrenal gland and peripheral adrenergic nerve terminals has been known for over 25 years²² and is now well established. So, magnesium sulfate can significantly attenuate the release of catecholamines at the time of laryngoscopy and tracheal intubation and thus reduce the severity of cardiovascular disturbances. Magnesium acts by slowing the atrial rate by inhibiting the calcium mediated depolarizing current in pacemaker tissue^{23,24} and, therefore, the overall effect is the mild increase in heart rate.

In our study we have compared dexmedetomidine with Magnesium sulfate to attenuate the stress response to laryngoscopy and endotracheal intubation. The mean fall in HR and SBP observed just after intubation, at 2 and 5 minutes after intubation in Group A was statistically highly significant compared to mean HR in B group (p=0.000).

The fall in mean DBP and mean MAP values at 2 and 5 minutes of intubation were statistically highly significant (p=0.000) in group A compared to group B.

In conclusion both dexmedetomidine and magnesium sulfate attenuated the cardiovascular response to laryngoscopy and intubation. Inj dexmedetomidine 1µg/kg was more efficient than inj. MgSO4 50mg/kg for attenuation of the cardiovascular response to laryngoscopy and intubation. Dexmedetomidine was associated with greater decrease in heart rate as compared to magnesium sulfate where a mild increase in HR could also be seen in some cases.

Similar findings were also found on several studies which corroborate the finding of the hemodynamic attenuation by the drugs. Chhaya Joshi et al²⁵, Krishna Chaithanya et al²⁶, Ismail M.A Ahmed et al²⁷, Vinit K. Srivastava et al²⁸, Lakshmi Mahajan et al²⁹, Kumkum Gupta et al³⁰, Nabin Pokhrel et al³¹.

CONCLUSION

From this study it was observed that Inj. Dexmedetomidine 1µg/kg caused better attenuation of cardiovascular response to laryngoscopy and intubation as compared to magnesium sulfate. It provided more stable HR, SBP, DBP and MAP during the stressful period following laryngoscopy and intubation. There were minimal side effects with no significant difference in both the groups.

On the basis of our present clinical comparative study, we can come to conclusion that

1. Both dexmedetomidine and magnesium sulfate attenuated the cardiovascular response to laryngoscopy and intubation.
2. Inj dexmedetomidine 1µg/kg was more efficient than inj. MgSO4 50mg/kg for attenuation of the cardiovascular response to laryngoscopy and intubation.
3. Dexmedetomidine was associated with greater decrease in heart rate as compared to magnesium sulfate where a mild increase in HR could also be seen in some cases.

1. Conflict of Interest- none
2. Source of Support - none

REFERENCES

1. Fox EJ, Sklar GS, Hill CH, Villanue Var, King BD. Complications related to the pressor response to endotracheal intubation. *Anesthesiology*. 1977; 47:524-5.
2. Kumar et al. Blocking Glossopharyngeal and superior laryngeal nerves to attenuate the cardiovascular response to laryngoscopy and endotracheal intubation. *IJA*, 1993; 41:20-25
3. Martin DE, Rosenberg H, Aukburg SJ. Low dose fentanyl blunts circulatory response to tracheal intubation. *Anesthesia Analgesia*, 1982; 6:680-84
4. Lev R, Rosen P. Prophylactic lidocaine use preintubation: a review. *Journal of Emergency medicine*, 1994; 12:499-506.
5. Miller RD, Eriksson LI, Fleisher LA, Wiener-Kronish JP, Young WL. *Miller's Anesthesia*. Philadelphia, Churchill Livingstone, 7th edition; 1585-1600
6. Mikawa K, Nishina K, Maekawa N, Obara H. Comparison of nicardipine, diltiazem, and verapamil for controlling the cardiovascular responses to tracheal intubation. *BJA*, 1996; 76:221-26
7. Stoelting RK. Attenuation of blood pressure response to laryngoscopy and tracheal intubation with sodium nitroprusside. *Anesthesia analgesia*, 1979; 58:116-19
8. Fassoulaki A, Kaniaris P. Intranasal administration of nitroglycerine attenuates the pressor response to laryngoscopy and intubation of the trachea. *BJA*, 1983; 5:49-52.
9. A. Altan, N. Turgut, F. Yildiz, A. Turkmen and H. Ustun: Effects of magnesium sulphate and clonidine on propofol consumption, haemodynamics and postoperative recovery. *BJA*, 2005; 94 (4): 438-41
10. Meiklejohn BH, Coley S. Pressor and catecholamine response to nasal intubation of the trachea. *BJA* 1989; 63 (3):283-286
11. Stoelting RK et al. Circulatory changes during direct laryngoscopy and tracheal intubation: Influence of duration of laryngoscopy with or without prior lidocaine. *Anesthesiology*, 1977; 47:381-84.
12. Stoelting RK, Hiller SC. *Textbook of pharmacology and physiology in anesthetic practice*. Philadelphia, Lippincott Williams and Wilkins, 4th edition; 2006:340-45.
13. Bloor BC, Ward DS, Belleville JP, Maze M. Effects of intravenous dexmedetomidine in humans. II Haemodynamic changes. *Anesthesiology* 1992; 77:1134-1142.
14. S. Herroeder, M. E. Schonherr, S. G. De Hert, and M. W. Hollmann, "Magnesium—essentials for anesthesiologists," *Anesthesiology*, vol. 114, no. 4, pp. 971–993, 2011.
15. S.-H. Do, "Magnesium: a versatile drug for anesthesiologists," *Korean Journal of Anesthesiology*, vol. 65, no. 1, pp. 4–8, 2013.
16. King BD, Harris LC, et al. Reflex circulatory responses to direct laryngoscopy and tracheal intubation performed during general anaesthesia. *Anesthesiology*, 1951; 12:556-566.
17. Link RE, Desai K, Hein L, Stevens ME, Chruscinski A, Bernstein D, et al. Cardiovascular regulation in mice lacking alpha2-adrenergic receptor subtypes b and c. *Science*. 1996; 273:803-805.
18. Kobilka BK, Matsui H, Kobilka TS, Yang-Feng TL, Francke U, Caron MG, et al. Cloning, sequencing, and expression of the gene coding for the human platelet alpha 2-adrenergic receptor. *Science*. 1987; 238:650-656.
19. Regan JW, Kobilka TS, Yang-Feng TL, Caron MG, Lefkowitz RJ, Kobilka BK. Cloning and expression of a human kidney cDNA for an alpha 2-adrenergic receptor subtype. *Proc Natl Acad Sci USA*. 1988; 85:6301-6305.
20. Snapir A, Posti J, Kentala E, Koskenvuo J, Sundell J, Tuunanen H, et al. Effects of low and high plasma concentrations of dexmedetomidine on myocardial perfusion and cardiac function in healthy male subjects. *Anesthesiology*. 2006; 105:902-910.

21. Iseri LT, French JH. Magnesium: Nature's physiologic calcium blocker. *Am Heart J* 1984; 108:18 & 93.
22. Douglas WW, Rubin RP. The mechanism of catecholamine release from the adrenal medulla and the role of calcium in stimulus-secretion coupling. *J Physiol* 1963; 167:28 & 310.
23. Turlapaty PDMV, Carrier O. Influence of magnesium on calcium induced responses of atrial and vascular muscle. *J Pharmacol Exy Ther* 1973; 187:8 & 98.
24. Somjen GG, Baskerville EN. Effect of excess magnesium and vagal inhibition and acetylcholine sensitivity of the mammalian heart in situ and in vitro. *Nature* 1968; 217:679-80.
25. Chhaya Joshi, Anilkumar Ganeshnavar, Shilpa Masur . A COMPARATIVE STUDY BETWEEN INTRA VENOUS DEXMEDETOMIDINE AND MAGNESIUM SULFATE IN ATTENUATION OF CARDIOVASCULAR RESPONSE TO LARYNGOSCOPY AND ENDOTRACHEAL INTUBATION – A RANDOMIZED CLINICAL TRIAL, *Intl. J Clin. Diag. Res.* 2016; 4(3):II
26. Krishna Chaithanya, Jagadish Vaddineni, Narasimha Reddy et al. "A Comparative Study between I.V 50% Magnesium Sulphate and Dexmedetomidine for Attenuation of Cardiovascular Stress Response during Laryngoscopy and Endotracheal Intubation". *Journal of Evolution of Medical and Dental Sciences* 2014; Vol. 3, Issue 32, August 04; Page: 8741-8749.
27. Ahmed IM, Abdelraouf HS. Magnesium sulfate, dexmedetomidine, and lignocaine in attenuating hypertension during