



ORIGINAL RESEARCH PAPER

Surgery

A COMPARATIVE STUDY BETWEEN OPEN INGUINAL HERNIOPLASTY VS LAPAROSCOPIC TRANSABDOMINAL PRE-PERITONEAL HERNIOPLASTY(TAPP)

KEY WORDS: Inguinal hernia repair, Laparoscopic hernioplasty, Open hernioplasty, Transabdominal Pre-Peritoneal Hernioplasty (TAPP)

Dr. A. A. Ghasura (HOU & HOD) Civil Hospital Ahmedabad

Dr. Ashini J Jadav* 1st year Surgery *Corresponding Author

Dr. Rishavkumar 1st year Surgery

ABSTRACT

Background: - The word "hernia" derived from Latin term meaning "a rupture"^[2] A hernia is defined as an area of weakness or complete disruption of fibromuscular tissue of body wall can pass through or herniate through defect.^[2] In all ventral abdominal hernias, Inguinal hernia is the most common. Inguinal hernia repair is common in surgical practice. In past, the repair was pure open tissue repair which was changed to open prosthetic mesh repair. Laparoscopic groin hernia repair was first performed by Ger in 1979.^[2] The laparoscopic approach to hernia repair has since evolved into common and effective procedure.

Methods: - The presented study is a prospective, open labelled, randomized, comparative study during a period from July-2020 to till date. After taking written consent from the patients were randomly allocated to two groups. **Group A**, Open hernioplasty were operated with open tension free Lichtenstein's hernioplasty. **Group B**, laparoscopic hernioplasty were operated with Transabdominal Preperitoneal (TAPP) repair. Those patients who fulfilled the inclusion criteria are included in study. The mode of presentation, surgical treatment and postoperative complications were all evaluated and compared.

Results: - During the study there were no major life-threatening complications were reported in either group. The postoperative pain and need of analgesics, presence of surgical site infections more in Open hernioplasty group compared to laparoscopy group. The time taken for return to routine daily activities and work was earlier in patients treated with laparoscopy compared to open surgery and good cosmetic results.

Conclusions: - Laparoscopic hernia repair has better, evolving & effective procedure in recent days compared to open hernia repair in terms of less postoperative complications, early recovery with good cosmetic results.

Aims & Objectives:-

- To compare the time taken for repair, patient's hospital duration of stay, surgical repair, complications that occur intraoperatively and postoperatively and outcome of open inguinal hernia mesh repair and laparoscopic hernia mesh repair
- To evaluate the limitations of laparoscopic TAPP Inguinal Hernioplasty

INTRODUCTION

Repair of inguinal hernia is one of the commonest operations performed by surgeons around the world. The treatment has been in evolution from the pure tissue repairs to the prosthetic repairs and in the recent laparoscopic repair. Preferred approach for open inguinal hernia repair is Lichtenstein's tension free open inguinal hernioplasty using a prosthetic mesh. The recurrence rate is 10.1% in laparoscopic group compared to open hernioplasty (4.9%)^[1,2]. Postoperative morbidity is low and recovery is quick in laparoscopy group.^[2]

Laparoscopy had gained widespread acceptance in today's era of surgery. Several studies have shown the benefit of the laparoscopic hernioplasty over open hernioplasty (OH) in terms of less postoperative pain and morbidity, wound complications, postoperative pain, early resumption of daily activities and work with better cosmetic results.^[2] But it had some limitations like longer operative time, longer learning curve, higher cost, a potential for serious life threatening accidents and a higher recurrence rate as compared with open surgery.^[2]

Laparoscopic hernioplasty can be accomplished in two ways i.e. Trans-Abdominal Preperitoneal repair (TAPP) and Totally Extraperitoneal repair (TEP). TEP repair and open hernioplasty does not need invasion of the peritoneal cavity. Technically it eliminates the hazards of intra operational injuries. The objective of the study was to evaluate the safety and efficacy of laparoscopic hernia repair & standard open anterior tension free Lichtenstein's hernioplasty

METHODOLOGY

The present study is a prospective study of 50 cases of an

inguinal hernia admitted in Civil Hospital Ahmedabad during the study period of July-2020 to till date. Written and informed consent was taken from the patients. All the laparoscopic hernia operation performed by a same operative surgeon. 50 cases for the purpose of the study were selected by the nonprobability (purposive) sampling method.

INCLUSION CRITERIA

Adults above 20 years and below 75 years of age of Male patients ASA (American Society of Anaesthesia) grade I & II with Primary unilateral or bilateral inguinal hernia are included in study.

EXCLUSION CRITERIA

Patients with complicated hernia (irreducible, obstructed, strangulated), those with recurrent hernia, female patients were excluded. Those unfit for general anaesthesia (ASA Grade ≥ III), laparoscopy or pneumoperitoneum i.e. those with cardiac diseases (MI, IHD), respiratory diseases (chronic asthma, COPD), renal or hepatic diseases, bleeding disorders, immunocompromised patients with malignancy etc are excluded from the study.

Choice of Procedure

The procedure was based on the general condition of patients and associated cost of the procedure.

All patients were clinically evaluated and underwent routine investigations for fitness. Elderly patients above 50 years of age are evaluated for prostate enlargement by digital rectal examination & ultrasonography. After taking written consent from the patients are randomly allocated with help of lottery technique into study group A & group B.

Surgical Procedure

1. Open Lichtenstein's Tension free inguinal hernioplasty

The patient was laid down in the supine position under spinal anaesthesia. After the asepsis using betadine solution, an oblique incision of approximately 4-6 cm long kept over inguinal canal. Dissection done upto separation of spermatic cord and hernia sac. Hernial sac opened & content reduced into abdominal cavity and closed with Polyglactin suture. With help of On-lay technique Polypropylene mesh placed & fixed. External oblique sheath, subcutaneous tissue & skin closed.

2. Transabdominal Preperitoneal Repair (TAPP)

For the TAPP procedure, the patient laid down in the Trendelenburg position under general anaesthesia. Pneumoperitoneum was created around the umbilical region inserting three trocars, a 10mm at umbilical level and two 5 mm in midclavicular plane on the left and right lumbar region. An incision was made on the peritoneal membrane from anterior superior iliac spine to pubic tubercle with help of electrocautery. The peritoneal membrane mobilized & preperitoneal space created. A Polypropylene mesh fixed using tacker. Peritoneum closed with vicryl suture. Port site closed with Polyglactin Port closure.




Fig 1: - Open Inguinal Hernioplasty




Fig 2: - Laparoscopic TAPP Hernioplasty

Patients were admitted one day prior to surgery. They were operated as per allocated group and relevant operative findings were noted. The antibiotic protocol was preoperative antibiotics consisting of intravenous dose of injectable amoxicillin+ Clavulanic acid 1.2 gm given in morning as prophylaxis. The analgesic used in postoperative period. The patients were encouraged to move in the early postoperative period and to take liquid diet on the evening of surgery. Foley's catheter, which was inserted in all patients in laparoscopic and bilateral inguinal hernia and was removed in the next morning and the patients were discharged in the 2nd & 3rd postoperative day after surgery.

Sutures were removed between 7-10 days. The wounds were checked and graded accordingly. Patients were postoperatively evaluated for presence of any cough impulse, swelling, and signs of recurrence. The follow up done during postoperative in 1st week, 2nd week, 1 month & 6 month. The scars were checked at each visit.

Descriptive statistics were used where ever is applicable. All the parameters are compared between the two groups. The mean age of the patients, time taken for surgery, duration of hospital stay & time taken to return normal activities are compared.

The distribution and type of hernias, ease of detection, amount of bleeding during surgery, operative time for surgery, postoperative surgical site infection and cosmetic appearance results are compared.

RESULTS

A total 50 Male patients are included in the study out of which 38 patients had open Hernioplasty allocated into group A and 12 patients had laparoscopic TAPP Hernioplasty into group B. There was no mortality or surgery related major complications in both groups.

The mean age of the patients in two groups are compared.

Table-1: - Comparison Of Mean Age In Study Groups

Type	Open Hernioplasty	Laparoscopic TAPP Hernioplasty	TOTAL
Mean age	39.63	45.20	42.41

Table-2: -site Of Hernia In Study Group

Site	Open Hernioplasty	Laparoscopic TAPP Hernioplasty	TOTAL
Right	18	6	24
Left	13	3	16
Bilateral	7	3	10
TOTAL	38	12	50




Table 3: -Types of Inguinal Hernia

Type	Open Hernioplasty	Laparoscopic TAPP Hernioplasty	TOTAL
Direct	11	10	21
Indirect	25	2	27
Pantaloon	2	0	2
TOTAL	38	12	50

The type & side of hernia in both the groups reveals that, **Right side Indirect inguinal hernia** more common.




Table 4: - Intraoperative & Postoperative complications in study groups

Complications	Open Hernioplasty	Laparoscopic TAPP Hernioplasty	TOTAL
Nil	30	10	40
Vas deference injury	1	0	1
Vascular injury	0	0	0
Seroma	2	1	3
Infection	2	0	2
Chronic wound Pain	3	1	4

Postoperative wound infection is 10.5% in open group and 0% in laparoscopic group. Postoperative chronic pain was observed in open group was 18.42% and in laparoscopic group 4.16%.




Table 5: - Operative Mean time based on type & site of hernioplasty

Operative time	Open hernioplasty	Laparoscopic hernioplasty
Direct	58.6 ± 4.2 minutes	63.4 ± 6.2minutes
Indirect	61.5 ± 4.5 minutes	65.9 ± 7.4 minutes
Unilateral	59.7 ± 4.3 minutes	62.1 ± 6.9 minutes
Bilateral	116.8 ± 9.1 minutes	122.9 ± 10.5 minutes

The mean operative time for Open hernioplasty (OH) group was 58.4 ± 2.4 minutes and that for laparoscopic group was 65.62 ± 6.23 minutes. During the initial period in laparoscopic procedure, mean time for surgery was 75.2 ± 4.8 minutes which gradually reduced to 65.62 ± 6.23 minutes. Mean time during initial laparoscopic procedure was higher which was gradually reduced due to experience in laparoscopic procedure.

The mean postoperative hospital stay after laparoscopic repair was 1.68 days (ranges, 1-3 days) & in open group was 2.40 days (ranges, 1-4 days). All the patients underwent laparoscopic repair were discharged within 24-48 hours.

Table 6: - Mean time to return daily activities and work

	Open hernioplasty	Laparoscopic hernioplasty
Mean time	22.7 ± 10.4 days	10.5 ± 6.3 days

The mean time to return to activity also was lower in the laparoscopic group compared to open group. Patients in laparoscopic group returned to their work early compared to open hernioplasty group.

DISCUSSION:-

Laparoscopic surgery has led to many changes in the management of inguinal hernia and significantly reduced the morbidity associated with open surgical procedures. At present, the laparoscopic repair of hernias gained clinical importance in patients with bilateral or recurrent hernias.^[4]

In my study, time taken the laparoscopic procedures was 65.62 ± 6.23 minutes & in Open hernia repair was 58.4 ± 2.4 minutes. In my study, the mean time during initial laparoscopic procedure was higher which was gradually reduced due to experience in laparoscopic procedure. It was compared with the study of Lal et al in which laparoscopic repair mean time was 75.72 ± 4.6 minutes & in open repair mean time was 54.00 ± 3.16 minutes. Time taken for laparoscopy procedure was more than open procedure.^[6]

In my study It was found that the return to daily activities and work in laparoscopy group was 10.5 ± 6.3 days and in open group was 22.7 ± 10.4 days. It was compared with the study of Lal in which mean time for return to daily activity in laparoscopy group was 12.8 ± 7.1 days & in open repair group was 19.3 ± 4.3 days. It was found that return to daily activities was earlier in the laparoscopic group than open group.^[6] This was due to minimal pain in the postoperative period and less chances of cord oedema.^[11-13]

Chronic wound related pain described as dull aching pain or sensation of mesh pricking in the groin. This is mainly related to nerve injuries which occur in Open hernioplasty due to cutting of nerves and also use of electrocautery around nerves. In this study the postoperative chronic pain was observed in open group was 18.42% and in laparoscopic group 4.16%. This finding was compared with the study of Winslow et al in which chronic wound pain was 6.8% in open group and 0 % in laparoscopic group. It was found that postoperative pain following laparoscopic surgery was lower compared to Open surgery group.^[11-13]

In laparoscopy there is smaller size of incision, no need of extra or bilateral incisions in case of bilateral hernias, minimal dissection and less handling of cord structures.^[11-13]

In study, the postoperative hospital stay in laparoscopic group was 1.68 days and in open group was 2.40 days. This was similar with study of Colak et al in which postoperative hospital stay in laparoscopy group was 1.80 days and in open group was 2.73 days. So, the postoperative hospital stay in laparoscopic group was shorter than open group.^[7]

In study, postoperative wound infection is 10.5% in open group and 0% in laparoscopic group. This was similar with study by Lal et al in which wound infection in laparoscopic group was 1.1% and in open group was 3.9%. The incidence of wound infection though slightly higher in open hernioplasty group.^[6]

CONCLUSION:-

Laparoscopic hernia repair is better and evolving surgical procedure compared to open repair. It has advantages like minimal postoperative wound pain, early return to daily activities, better cosmetic results. But it has limitations like longer operative time and long learning curves.

Funding:No funding sources

Conflict of interest:None declared

REFERENCES

1. Norman S Williams, P. Ronan O’Connell, Andrew Mccaskie in: Textbook of Bailey & Love’s Short Practice of Surgery. 27th edition. Abdominal wall, hernia & umbilicus. Chapter 60. 2018; 1028-1034
2. Lokesh Bathla, Robert J Fitzgibbons. Perspective on laparoscopic inguinal hernia. In: Maingot’s Abdominal operations. Chapter 8. 12th edition. Mc Graw Hill publication; 2013, 160-162
3. Amid PK. Lichtenstein’s Tension Free Hernioplasty. In: Mastery of Surgery chapter 176. 5th edition. Lippincott Williams and Wilkins’s publications; 2007: 1932-1939.
4. Liem MS, van Duyn EB, Van der Graaf Y, Van Vroonhoven TJ. Cola Trail Group; Recurrences after conventional anterior and laparoscopic inguinal hernia repair: a randomized comparison. Ann Surg. 2003;237(1):136-41
5. Neumayer L, Giobbie Hurder A, Jonasson O, Fitzgibbons R Jr, Dunlop D, Gibbs J, et al. Veterans Affairs Cooperative Studies Program 456 Investigators. Open mesh versus laparoscopic mesh repair of inguinal hernias. N Engl J Med. 2004;350(18):1819-27.
6. Lal P, Kajla RK, Chander J, Saha R, Ramteke V K. Randomized controlled study of laparoscopic total extra peritoneal versus open Lichtenstein’s inguinal hernia repair. Surg Endosc. 2003;17(6):850-6.
7. Colak T, Akca T, Kanik A, Aydin S. Randomized clinical trial comparing laparoscopic totally extra peritoneal approach with open mesh repair in inguinal hernia. Surg Laparosc Endosc Percut Tech. 2003;13(3):191-5.
8. Rattner D, Michael Van Noord. Totally Extra peritoneal Inguinal Hernia Repair. In: Mastery of Surgery, Chapter 1743. Fifth edition. Lippincott, Williams & Wilkins Publications; 2007:1905-1911
9. Jobe BA, Hunter JC. Minimally Invasive Surgery. In: Schwartz’s Principles of Surgery. 8th edition. Mc Graw publication; 2005:379-402.
10. Bringman S, Ramel S, Heikkinen TJ, Englund T, Westman B, Anderberg B. Tension free inguinal hernia repair: TEP versus mesh plug versus Lichtenstein: a prospective randomized controlled trial. Ann Surg. 2003;237(1):142-7.
11. Fujita F, Lahamann B, Otsuka K, Lyass S, Hiatt JR, Phillips EH. Quantification of pain and satisfaction following laparoscopic and open hernia repair. Arch Surg. 2004;139(6):596-602.
12. Winslow ER, Quasebarth M, Brunt LM. Perioperative outcomes and complications of open vs. laparoscopic extra peritoneal inguinal hernia repair in a mature surgical practice. Surg Endosc. 2004;18(2):221-7.
13. Pokorny H, Klinger A, Scheyer M, Fugger R, Bischof G. Postoperative pain and quality of life after laparoscopic and open inguinal hernia repair: results of a prospective randomized trial. Hernia. 2006;10(4):331-7.