



ORIGINAL RESEARCH PAPER

General Medicine

“STUDY OF URINARY TRACT INFECTIONS IN HEALTH CARE WORKER IN DEPARTMENT OF MEDICINE,,

KEY WORDS: UTI, urine culture, bacteriuria, Health care associated infections, UTI during pregnancy.

Dr. Amrita Kumari

Senior Resident, Department of General medicine PMCH Patna.

Dr. Pragati Prabhat

Senior Resident, Department of General medicine PMCH Patna.

Dr. Sunil Kumar*

Senior Resident, Department of General medicine PMCH Patna.
*Corresponding Author

ABSTRACT

Background: Urinary tract infection (UTI) is the most common infection experienced by humans after respiratory and gastro intestinal infections, and also the most common cause of both community acquired and nosocomial infections for patients admitted to hospitals. Injudicious use of antibiotics and better adaptation of organisms to the hospital environment contribute to increase in HAIs. There are additional patient safety concerns associated with urinary catheter insertion, like patient discomfort, activity restriction, discharge delays, and the potential development of a reservoir of multidrug-resistant organisms that can be spread to other patient.

Material and Methods: Patients admitted in wards of medicine department at Patna medical college and hospital Patna Bihar. Patients who were admitted for at least for past 48 hours, or readmitted in less than 14 days after their discharge from hospital. Sociodemographic data from all the included patients was collected from medical records. Clinical history was obtained from all the patients. Signs of HAI were observed and recorded. Bacteremia was defined as presence of bacteria in the blood confirmed by at least one positive blood culture.

Results: A total of 238 patients were included in the study who were admitted in the medicine wards. Out of 238 patients included in the study, 31 (13.02%) presented one or several signs suggestive of HAI and they received microbiological tests. 20 (8.40%) cases were confirmed or potential HAI. Of the 238 patients 109 (54.20%) were male and 129 (45.79%) were female. **Conclusion:** Prevalence of HAI in Urinary tract infection was 13.02% in this study, patients undergoing long term treatment are also vulnerable to the infection due to moist hospitalized conditions, can be used as an alternative to counteract the pathogen causing UTI.

INTRODUCTION

Urinary tract infection is a common contagion among men and women but the incidence is quite high among women due to their physiology, it can be referred as a condition which women will certainly encounter during the span of their life time and the prevalence is higher among women during pregnancy. Health care workers infections (HCAIs) or hospital acquired infections (HAIs) are infections that occur during hospitalization but they are neither present nor incubating before hospital admission. Health care-associated urinary tract infection (UTI) is an important cause of morbidity and increased health care costs in hospitals. More than 100 million indwelling urinary catheters e.g. Foley catheters are used annually in the world. Most of these catheters are used for short-term catheterization i.e. 30 days or less. Catheter-associated UTI (CAUTI), the most common of all health care-associated infections (HAIs) accounts for approximately 40% of all HAIs. The Centers for Disease Control and Prevention (CDC)

recommends to prevent these infections through appropriate antibiotic use and infection prevention practices¹⁷. Rapid advances in medical field, injudicious use of antibiotics and better adaptation of organisms to the hospital environment contribute to increase in HAI. Though the infection seems to be harmless in the initial stages, the patient shows a variety of symptoms as the stage progresses and can lead to death in severe circumstances, research studies have defined UTI as the most common form of bacterial infection.

It is shown that approximately half of all cases of HCAIs are associated with medical devices and implants. In majority of the cases there is improper insertion of the catheter and CAUTI risk increases considerably with duration of catheterisation. In HAUTI majority of causative agents are Gram-negative bacteria and emergence of antimicrobial resistance is of great concern with respect to urinary tract infections (UTIs). In a European multicentre study posted that the proportion of infected patients in intensive care units can

be as high as 51%; most of these are health care associated. There are additional patient safety concerns associated with urinary catheter insertion, like patient discomfort, activity restriction, discharge delays, and the potential development of a reservoir of multidrug-resistant organisms that can be spread to other patients. In a study, 42% of catheterized patients reported that their indwelling catheter was uncomfortable, 48% complained that it was painful, and 61% noted that it restricted their activities of daily living.

MATERIAL AND METHODS

The study was carried out in department of General Medicine at Patna medical college and Hospital Patna Bihar. Patients admitted in wards of medicine department were included in the present study. Informed written consent was obtained from all the patients.

INCLUSION CRITERIA:

Patients who were admitted for at least for past 48 hours, or readmitted in less than 14 days after their discharge from hospital.

EXCLUSION CRITERIA:

Patients who completed less than 48 hours of admission and those who received only outpatient care or had exclusively day hospital care were not included.

defined as infections seen after 48 hours of admission or within 14 days following discharge from the hospital. Confirmed HAI was defined as clinical signs and microbiological confirmation; Potential HAI: association of several clinical signs with no microbiological confirmation; Less potential HAI: a single clinical sign with no microbiological confirmation.

Sociodemographic data from all the included patients was collected from medical records. Clinical history was obtained from all the patients. Signs of HAI were observed and recorded. These signs were: hyperthermia Temperature >

38°C; Hypothermia Temperature ≤ 36°C; chills; urinary tract signs or lumbar pain, suprapubic, dysuria, urgency, or burning micturition. HIV status of all the patients, antibiotics administered during admission, site of HAI, existence of invasive device like urinary catheter, venous catheter, sensitivity to antibiotics, and evolution of healthcare associated infection was done.

Bacteremia was defined as presence of bacteria in the blood confirmed by at least one positive blood culture. However, blood culture must be confirmed by clinical signs such as fever (Temperature ≥ 38.5°C) or hypothermia (Temperature ≤ 36.5°C), chills or hypotension.

Statistical analysis was done out through SPSS 21.0 software, Continuous variables were expressed in form of mean values with their standard deviation, or medians with their interquartile ranges. Categorical variables were expressed in percentage. P value of <0.05 was considered as statistically significant.

RESULTS

A total of 238 patients were included in the study who were admitted in the medicine wards. Out of 238 patients included in the study, 31 (13.02%) presented one or several signs suggestive of HAI and they received microbiological tests. 20 (8.40%) cases were confirmed or potential HAI.

Table 1: Patients and HAI

HAI	N	%
One or several signs suggestive of HAI	31	13.02%
confirmed or potential HAI	20	8.40%

Table 2: Sociodemographic data

Variable	Total	Male (%)	Female (%)
Total patients	238	109 (54.20%)	129 (45.79%)
HAI patients	31	14 (45.16%)	17 (54.83%)
Mean Age (Total Patients)	46.52±16.74	46.32± 21.51	45.95 ± 14.25
Mean Age (HAI patients)	54.21± 17.41	52.41 ± 19.53	53.46 ± 16.77

Of the 238 patients 109 (54.20%) were male and 129 (45.79%) were female. 31 patients were diagnosed as HAI patients of which 14 (45.16%) were male and 17 (54.83%) were female. Mean age of HAI patients was 54.21 ± 17.41 while total mean of age was 46.52 ± 16.74. Mean age of male with HAI 52.41 ± 19.53 and female was 53.46 ± 16.77.

Table 3: Healthcare associated infection and urinary catheter

Health care associated infection (n=528)			
Urinary Catheter	Yes	No	Total
Yes	27 (11.34%)	178 (74.79%)	205 (86.13%)
No	4 (1.68%)	29 (12.18%)	33 (13.87%)
Total	31(13.03%)	207(86.97%)	238(100%)

Of the total 31 HAI cases 27(11.34%) had urinary catheter while 4 (1.68%) doesn't had catheter, over all prevalence of HAI was 31 (13.03%). Of the total 205 (86.13%) patients who had urinary catheter 178 (74.79%) doesn't show HAI. While 27 (11.34%) who doesn't had catheter 29 (12.18%) had no HAI and no catheter.

DISCUSSION:

The current review makes an attempt to put fourth the various that are responsible for the occurrence of the urinary tract infection and endeavors to confirm the significance of other factors like physiological changes during pregnancy, Virtually all healthcare-associated urinary tract infections are caused by instrumentation of the urinary tract, creating an

opportunity to prevent a large proportion of HAUTIs, including catheter- associated urinary tract infections (CAUTIs).HAI includes catheter-associated urinary tract infections (CAUTI), central-line-associated blood stream infections (CLABSI), and ventilator-associated pneumonias (VAP)^{xi}.The hospitals in developed countries generate their infection-control surveillance data from time to time as this is important for empirically treating infections especially in the intensive care unit setting. More than 20% of patients in Intensive Care Unit may be infected with various HCAI with mortality rate of >30%.

The 31 patients were diagnosed as HAI patients of which 14 (45.16%) were male and 17 (54.83%) were female. Mean age of HAI patients was 54.21 ± 17.41 while total mean of age was 46.52 ± 16.74. Mean age of male with HAI 52.41 ± 19.53 and female was 53.46 ± 16.77. Surveillance of HAI is an essential element so as to know the current prevalence of the condition and to identify potential risk factors. During hospitalization, various HAI directly related to different invasive procedures such infection, surgical site infection are often encountered.

In this study prevalence of HAI urinary tract infection was 31(13.03%). Agrawal R et al. observed 6.67% HAI prevalence. In a study by Laborde G et al., reported the infection rate to be 36.3% from 314 patients treated longer than 48 hours in neurosurgical ICU. Prevalence of HAI in a study by Dia N M. et al in Senegal 10.9%^{xvii}.

In the present study of the total 7 culture positive patients with HAI Escherichia coli was isolated in 3 cases, Klebsiella pneumonia in 2 caesa, Pseudomonas aeruginosa and Staphylococcus aureus were isolated in 1 cases each. Hospitalized patients are at high risk of colonization with health care-associated pathogens. 4.5% were colonized or infected with MRSA, 2.7% were colonized or infected with vancomycin-resistant enterococci (VRE), 1.4% were colonized or infected with *C. difficile*, 1.3% were colonized or infected with an extended-spectrum β-lactamase (ESBL)-producing organism, and 0.1% were colonized or infected with carbapenem-resistant *Enterobacteriaceae* (CRE). Mody et al observed that in indwelling devices 15% were colonized with multidrug-resistant *Acinetobacter baumannii*.

CONCLUSION

Urinary tract infection is a common contagion among both gender, Prevalence of HAI in Urinary tract infection was 13.02%. Preventive measures, periodic active surveillance over a longer period is required to reduce the rate of healthcare associated UTI.

REFERENCES

- Platt R, Polk BF, Murdoch B, Rosner B. Mortality associated with nosocomial urinary-tract infection. *N Engl J Med*. 1982 Sep 9;307(11):637-42.
- Saint S, Veenstra DL, Sullivan SD, Chenoweth C, Fendrick AM. The potential clinical and economic benefits of silver alloy urinary catheters in preventing urinary tract infection. *Arch Intern Med*. 2000 Sep 25; 160(17):2670-5.
- Jarvis WR. Selected aspects of the socioeconomic impact of nosocomial infections: morbidity, mortality, cost, and prevention. *Infect Control Hosp Epidemiol*. 1996 Aug; 17(8):552-7.
- Centers for Disease Control and Prevention. Antibiotic resistance threats in the United States, 2013. <http://www.cdc.gov/drugresistance/threat-report-2013/pdf/ar-threats-2013-508.pdf> as urinary tract infection, pneumonia, blood stream
- Deorukhkar S. C., Saini S., Mathew S. Non-albicans Candida infection: an emerging threat. *Interdisciplinary Perspectives on Infectious Diseases*. 2014;2014:7.
- Kojic E. M., Darouiche R. O. Candida infections of medical devices. *Clinical Microbiology Reviews*. 2004; 17(2):255-267. doi: 10.1128/cmr.17.2.255-267.2004.
- Gould CV, Umscheid CA, Agarwal RK, Kuntz G, Pegues DA. Healthcare Infection Control Practices Advisory Committee: Guideline for prevention of catheter- associated urinary tract infections 2009. *Infect Control Hosp Epidemiol*. 2010;31(4):319-326.
- Chen, L.F., Chiu, C.-T., Lo, J.-Y. et al. Clinical characteristics and antimicrobial susceptibility pattern of hospitalised patients with community- acquired urinary tract infections at a regional hospital in Taiwan. *Healthcare Infect*. 2014; 19:20-25
- World Health Organization. Health care-associated infections fact sheet. Available at: [http:// www.who.int/gpsc/country_work/gpsc_ccisc_fact_she](http://www.who.int/gpsc/country_work/gpsc_ccisc_fact_she)

- et_en.pdf, 2016
10. Saint S, Lipsky BA, Baker PD, McDonald LL, Ossenkop K. Urinary catheters: what type do men and their nurses prefer? *J Am Geriatr Soc.* 1999 Dec; 47(12):1453-7.
 11. Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control.* 2008 Jun; 36(5):309-32.
 12. Haley RW, Culver DH, White JW, Morgan WM, Emori TG, Munn VP, Hooton TM. The efficacy of infection surveillance and control programs in preventing nosocomial infections in US hospitals. *Am J Epidemiol.* 1985 Feb; 121(2):182-205.
 13. Orsi GB, Scorzolini L, Franchi C, Mondillo V, Rosa C, Venditti M. Hospital-acquired infection surveillance in a neurosurgical intensive care unit. *J Hosp Infect.* 2006 Sep; 64(1):23-9.
 14. Wa aszek M. The analysis of the occurrence of nosocomial infections in the neurosurgical ward in the District Hospital from 2003-2012. *Przegl Epidemiol.* 2015; 69(3):507-14, 619-23.
 15. Agarwal R, Mohapatra S, Rath GP, Kapil A. Active Surveillance of Health Care Associated Infections in Neurosurgical Patients. *J Clin Diagn Res.* 2017; 11 (7): DC01-DC04.
 16. Laborde G, Grosskopf U, Schmieder K, Harders A, Klimek L, Hardenack M, Gilsbach JM. Nosocomial infections in a neurosurgical intensive care unit. *Anaesthesist.* 1993 Oct; 42(10):724-31.
 17. Dia, N.M., Ka, R., Dieng, C., Diagne, M.L., Fortes, L., et al. (2008) Results of the Prevalence Survey of Healthcare-Associated Infections in the Fann CHNU (Dakar, Senegal). *Medicine and Infectious Diseases*, 38, 270-274.
 18. Williams V, Simor AE, Kiss A, McGeer A, Hirji Z, Larios Control—Canada. Is the prevalence of antibiotic-resistant organisms changing in Canadian hospitals? Comparison of point-prevalence survey results in 2010 and 2012. *Clin Microbiol Infect.* 2015 Jun; 21(6):553-9.
 19. Mody L, Gibson KE, Horcher A, Prenovost K, McNamara SE, Foxman B, Kaey KS, Bradley S. Prevalence of and risk factors for multidrug-resistant *Acinetobacter baumannii* colonization among high-risk nursing home residents. *Infect Control Hosp Epidemiol.* 2015 Oct; 36(10):1155-62.