



ORIGINAL RESEARCH PAPER

Microbiology

PHENOTYPIC DETECTION, BIOFILM PRODUCTION AND MOLECULAR CHARACTERIZATION OF E. COLI ISOLATES RECOVERED FROM URINARY TRACT INFECTIONS IN A TERTIARY CARE HOSPITAL

KEY WORDS: UTIs, Biofilm, E.coli, ESBL, blaTEM-1, blaSHV, blaCTXM, blaNDM-1, and blaVIM-2

Dr. Atul Khajuria

Ph.D., Assistant Professor, Department of Microbiology, World College of Medical Sciences and Research & Hospital, Jhajjar, Haryana-124103.

ABSTRACT

Objective: To examine Carbapenem resistance, & biofilm production in E.coli isolates recovered from UTIs.
Methods: A prospective, Cross-sectional study was conducted in a 650 bedded tertiary care centre in Jhajjar, Haryana India from July 2019 to April 2020. A total of 52 isolates were recovered from Urinary Tract Infections (UTIs). Carbapenem resistance, biofilm production was checked phenotypically and the presence of β -lactamase genes by PCR. Isolate relatedness was determined by REP PCR, and ERIC PCR.
Results: From July 2019 to April 2020, in this study a total of 52 E.coli isolates recovered from UTIs, 39 isolates were ESBL producers positive for TEM, SHV and CTX-M genes, 28 were carbapenemase producers and 23 were metallo beta lactamase producers positive for DDST, CDST and MBL (IP/IPI) E-test. Out of these 23 MBL producers 15 had blaNDM-1 gene, and 8 had blaVIM-2 gene respectively.
Conclusions: The significant presence of both metallo beta lactamase genes and biofilm co-existence in the same isolates is worrisome.

INTRODUCTION-

Urinary tract infections (UTIs) exclusively contribute to emergence of antimicrobial resistance. *Escherichia coli* is one of the major etiologic agents for urinary tract infection, affect both male and female of all ages but female are more vulnerable. Worldwide the antibiotic resistant uropathogens are alarming. Therapeutic options have become somewhat limited because of the emergence of organisms carrying extended-spectrum β -lactamases (ESBL) and plasmid-mediated AmpC β -lactamases. The carbapenems (imipenem, meropenem, and ertapenem) are sometimes the only effective agents for treatment of severe infection caused by ESBL-positive *E. coli*. Carbapenemases enzymes recognize almost all hydrolysable β -lactams, and most are resistant to inhibition by all commercially viable β -lactamase inhibitors. Ambler class A (including KPC and GES), A, and Ambler class D (CHDLs or OXA-48) beta-lactamases had serine at their active site, while Ambler class B (including IMP, VIM, SIM, and NDM) are all metalloenzymes with zinc in active-site. Biofilms are the microbial communities that are irreversibly associated with a surface and are enclosed in a self-produced extracellular polymeric matrix. Metalloenzymes producing uropathogenic *E. coli* within the biofilm are more resistant to antimicrobial agents leave patients with very few or no antimicrobial options.^{[1][2][3][4][5]} This study provides an insight into the acquisition and spread of the MBL genes along with Biofilm production.

MATERIALS AND METHODS

The Bacterial Isolates -A prospective, cross-sectional study was conducted in a 650 bedded tertiary care centre in Jhajjar, Haryana, India from July 2019 to April 2020. A total of 52 clinically significant, non-duplicate *E. coli* isolates were recovered from BSIs of hospitalized patients admitted to the Medical and Surgical intensive care units. Bacterial identification was performed by routine conventional microbial culture and biochemical tests using standard recommended techniques^{[3][6]} All the strains were preserved in 15% glycerol-supplemented Luria-Bertani medium at \square 80°C for molecular analysis.

Antimicrobial Susceptibility Testing

The antimicrobial susceptibility test was performed by the Kirby Bauer's disc diffusion technique on Mueller-Hinton agar, as per Clinical Laboratory Standard Institute (CLSI) guidelines.^[7] The antibiotics tested were as follows (potency in \square g/disc): Ampicillin (10), Cefuroxime (30), Cefpodoxime(30), Cefixime(5), Ceftazidime (30), Cefazolin(30), Cefoxitin(30), Cefepime (30), Cefotaxime (30), Piperacillin (100), Piperacillin-Tazobactam (100/10),

Aztreonam (30), Imipenem (10), Meropenem (10), Ertapenem (10), Doripenem(10), Norfloxacin (10), Colistin(10), Gentamicin (10), Tobramycin (10), Amikacin (30), Netilmicin (30), Ciprofloxacin (5), and Levofloxacin (5) (Hi Media Laboratories Pvt. Ltd., Mumbai, India). *Paeruginosa* ATCC 27853, *E. coli* ATCC 25922, *E. coli* ATCC 35218 and *K.pneumoniae* ATCC 700603 were used as quality control strains. The degree of susceptibility of the test isolate to each antibiotic was interpreted as sensitive (S), intermediate resistant (I) or resistant (R) by measuring the zone diameter of inhibition.^[7]

MIC Determination

Minimum inhibitory concentrations (MIC) of antibiotics were determined by Microbroth dilution Assays and further confirmed by the E-test (bioMérieux, Marcy l'Etoile), France for individual antibiotic.^{[3][6][7]}

Phenotypic Screening For ESBL Detection

Isolates with reduced susceptibility to **Ceftazidime (17 mm), Aztreonam (17 mm), Cefotaxime (22 mm) and Ceftriaxone (19mm)** as recommended by CLSI guidelines, were selected for confirmation of ESBL production. Isolates were tested for ESBL production by standard CLSI double-disc diffusion method and double disc synergy test and using E test (bioMérieux, Marcy l'Etoile, France) for detecting the MIC. These tests were checked for quality using standard control ESBL negative strain of *E. coli* ATCC 25922.^{[3][6][7]}

Phenotypic Screening For Carbapenemase Production

Isolates with reduced susceptibility to ertapenem, meropenem and imipenem (diameter of zones of inhibition \leq 18mm) by disc diffusion method were screened for the production of carbapenemase.^{[3][6][7]}

Phenotypic Screening For Metallo Beta Lactamase Production

MHT, DDST, CDST and MBL (IP/IPI) E-test was performed to detect Carbapenemase as well as Metallo-beta-lactamase production as described previously.^{[3][4][6][9][7]}

DNA Extraction And Molecular Detection

DNA was extracted from the bacterial isolates using the spin column method (QIAGEN; GmbH, Hilden, Germany) as per manufacturer's instructions. PCR-based detection of beta lactamase (ESBL) genes (*bla_{TEM}*, *bla_{SHV}*, *bla_{CTXM}* and *bla_{OXA}*), Ambler class B MBLs (*bla_{IMP}*, *bla_{VIM}*, *bla_{GIM}*, *bla_{SIM}*, *bla_{SPM}* and *bla_{NDM}*), Ambler class D (*bla_{OXA-23}*, *bla_{OXA-24}*, and *bla_{OXA-48}*) and serine carbapenemases (*bla_{KPC}*, *bla_{GES}*, and *bla_{NMC}*) were carried out on the isolates by using Gene Amp 9700 PCR System (Applied Biosystems, Singapore).^{[4][9]} PCR products were run on 1.5%

agarose gel, stained with ethidium bromide visualized under UV light and photographed. The amplicons were purified using QIAquick PCR purification kit (QIAGEN; GmbH, Hilden, Germany) DNA sequencing and sequence analysis Automated sequencing was performed on an ABI 3730XL DNA analyzer using the Big Dye system (Applied Biosystems Foster City, CA, USA). Sequences were compared with known sequences using the BLAST facility (<http://blast.ncbi.nlm.nih.gov>).^{[3][4][5][6]}

Detection Of Biofilm Production:

E. coli isolates producing biofilm were detected using Congo Red Agar method (qualitative) and Microtiter plate quantitative assay. The biofilm formation on microtiter plate were detected using 0.1% crystal violet and safranin separately and then the optical density (O.D) of each well were measured at 490 nm and 570 nm wavelength using automated (Lisa Scan EM Microplate Reader). The biofilm producers were differentiated as high, moderate, weak and non-adherent based upon the comparison with control.^{[4][5]}

Molecular Typing Of The Strains

Repetitive element-based PCR (REP-PCR) and Enterobacterial Repetitive Intergenic Consensus (ERIC – PCR) assays were performed, to rapidly characterize metalloenzymes producing *E. coli* strains which were recovered from the patients. Similarity clustering analysis was performed using unweighted pair group method with arithmetic mean and Dice coefficient. Clinical isolates with a similarity coefficient >85% were considered clonal.^{[4][5]}

RESULT AND DISCUSSION

Out of 52 isolates, 39 (75%) were found to be ESBL producers resistant to **Ceftazidime, Aztreonam, Cefotaxime and Ceftriaxone (MIC ≥ 16 µg/mL)**. Out of these 28 (53.8%) exhibited reduced susceptibility to **ertapenem, meropenem and imipenem (MIC ≥ 8 µg/mL)**. Among these 28 isolates, 23 had shown metallo beta lactamase enzyme production (44%). (Table -1) These 23 *E. coli* isolates were detected producing biofilm using Congo Red Agar method (qualitative) and Microtiter plate quantitative assay as compared with controls. Table -2 showing antibiograms of Metallo beta lactamase producing *E. coli*. Intensive care units provide an ideal environment for the dissemination of resistant determinant genes within the organisms. There are multiple risk factors associated with both environment and the patient that allow the development and spread of such pathogens. ICU patients often have frequent hospital admissions with their respective underlying medical conditions and there has been increase in the risk of colonization by multi-drug-resistant pathogens. Carbapenem resistant isolates are generally resistant to most other classes of antibiotics, while usually retaining susceptibility to Polymyxin B and colistin. PCR amplification for Ambler class D (*bla_{OXA-23}*, *bla_{OXA-24}*, and *bla_{OXA-48}*) and serine carbapenemases (*bla_{KPC}*, *bla_{GES}* and *bla_{NMC}*) was found to be negative for all isolates. Out of 23 MBL producing *E. coli*, 15(65%) had *bla_{NDM-1}*, while 8 (35%) had *bla_{VIM-2}*, *bla_{TEM-1}*, *bla_{SHV-12}*, *bla_{CTXM-15}* and *bla_{NDM-1}* gene were found in 9, Six had copresence *bla_{TEM-1}*, *bla_{SHV-28}*, *bla_{CTXM-14}* and *bla_{NDM-1}*, *bla_{TEM-1}*, *bla_{SHV-12}*, *bla_{CTXM-15}* and *bla_{VIM-2}* gene were present in 6 isolates, while *bla_{VIM-2}* in association with *bla_{TEM-1}*, *bla_{SHV-28}* and *bla_{CTXM-14}* was present in two isolates. The results of Microtiter plate assay for detection of quantitative biofilm formation were compared by staining with both 0.1% crystal violet and 0.1% safranin. Cut off value for Crystal violet had shown Strong = ≥ 0.72 O.D. while Safranin had ≥ 0.83 O.D. for strong biofilm. Strain molecular typing of 23 strains of *E. coli* by REP PCR generated 3 cluster with an average of 6 to 12 fragments while ERIC PCR produced 5 clonal clusters with an average of 5 to 16 fragments. This showed that ERIC -PCR was much better as compared to REP PCR.

CONCLUSION

This study identify the metalloenzymes producing *E. coli* and

their association with biofilm production. The study revealed that carbapenem resistant isolates are strong biofilm producer hence can be useful for clinical infection control purposes.

Declarations-

Financial Support And Sponsorship- Nil.

Conflict Of Interest-None to declare

Table -1 Showing Phenotypic Characterization Of *E. coli* Isolates Recovered From UTIs

Number of isolates	Carbapenem resistance by disc diffusion	ESBL	CDST	DDST	MBL-test	Biofilm
52	28	39	23	23	23	23

Table -2 Antibiograms Of Metallo Beta Lactamase Producing *E. coli* Along With Interpretive Criteria For Each Drug.

Antibiotic	Zone size in mm	MIC (µg/mL)
Ampicillin	≤13	≥32
Ampicillin sulbactam	≤11	≥32/16
Amoxicillin-Clavulanic Acid	≤13	≥32/16
Amikacin	≤16	≥64
Aztreonam	≤17	≥16
Cefazolin	≤19	≥8
Cefepime	≤18	≥16
Cefotaxime	≤22	≥4
Cefoxitin	≤14	≥32
Ceftazidime	≤17	≥16
Ceftriaxone	≤19	≥4
Cefuroxime	≤14	≥32
Cefixime	≤15	≥4
Ciprofloxacin	≤21	≥1
Colistin		≤2
Ertapenem	≤18	≥2
Gentamicin	≤12	≥16
Imipenem	≤19	≥4
Levofloxacin	≤16	≥2
Meropenem	≤19	≥4
Netilmicin	≤12	≥32
Norfloxacin	≤12	≥4
Piperacillin	≤17	≥128
Piperacillin/tazobactam	≤17	≥128/4
Doripenem	≤19	≥4
Tobramycin	≤12	≥16

REFERENCES –

- Critchley IA, Cotroneo N, Sulham K A, Melnick D, Mendes R. Resistance to Oral Antibiotics among Urinary Tract Infection Isolates of Escherichia coli from the United States and Europe in 2017. Poster #1465 presented at ID Week 2019, October 2-6, Washington, DC, USA
- Donnenberg M S. Enterobacteriaceae; In: Mandell G L, Bennett J E, Dolin R. (Eds.). Mandell, Douglas, and Bennett's principles and practice of infectious diseases. 7th edition. Philadelphia: Elsevier Churchill Livingstone, 2010; 2815-2833.
- Koneman EW, Allen SD, Janda WM, Schreckenberger PC, Winn WC. The Enterobacteriaceae. In Koneman EW, Allen SD, Janda WM, Schreckenberger PC, Winn WC (eds) Color Atlas and Textbook of Diagnostic Microbiology. 6th Edition. Lippincott Williams and Wilkins, 2006; 211–302.
- Khajuria A, Praharaj AK. Detection of carbapenem resistance genes in *E. coli* in a tertiary care hospital. World journal of pharmacy and pharmaceutical sciences. 2019; 8: 726-751.
- Khajuria A, Praharaj AK. Incidence and detection of metallo-beta-lactamase producing Klebsiella pneumoniae in a tertiary care hospital. European Journal of Pharmaceutical and Medical Research. 2019; 6: 324-340.
- Collee JG, Miles RS, Wan B. Tests for the identification of bacteria. In: Collee JG, Fraser AG, Marmion BP, Simmonds A editor. Mackie and Mc Cartney Practical Medical Microbiology. 14th ed. Edinburgh: Churchill Livingstone, 1996; 131–150.
- Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing: twenty-second informational supplement. Document M100-S30^{ed}. Wayne, PA: CLSI; 2020.