



**ORIGINAL RESEARCH PAPER**

**General Medicine**

**A STUDY OF ECG AND ECHOCARDIOGRAPHY FINDINGS IN PATIENTS WITH COPD**

**KEY WORDS:**

**Dr. Kartik Patel** M.D., General Medicine

**ABSTRACT**

**Background**- COPD is a very common preventable disease. Cardiac involvement is common in COPD patients. Early detection of cardiac involvement by ECG, Echocardiography can prevent morbidity and mortality in COPD patients. **Aim** - To study clinical profile and to assess cardiac dysfunction in COPD patients by ECG and Echocardiography. **Method** - An observational study was done in COPD patients age >30 years in civil hospital, navsari. ECG, Echocardiography were done .Responses were collected and analysed by microsoft excel. **Result**- ECG findings in copd patients - right axis deviation(44%), P pulmonale (37%), low voltage QRS complex (10%) poor R wave progression (39%), right ventricl hypertrophy (26%). Echocardiography findings were PAH (62%), corpulmonale (56%), RVD (45%), RAD (37%) right ventricle hypertrophy (27%), left ventricle systolic dysfunction(31%), left ventricle diastolic dysfunction(6%). **Conclusion** - COPD is a common and preventable disease. COPD patients should be assessed by ECG and echocardiography for better prognosis and to prevent complication.

**INTRODUCTION**

Chronic obstructive pulmonary disease is one of the most important cause of morbidity and mortality throughout the world<sup>1</sup>.The COPD burden is expected to rise in coming years because of continuous exposure to risk factors and aging of the population<sup>2</sup>.

COPD is very common, treatable & preventable disease. It is characterized by respiratory symptoms due to alveolar, airway abnormalities and exposure to noxious gases or particles .Dyspnoea, cough and sputum production are the most common symptoms. Important risk factor is smoking. Other environmental factors which contribute are airpollution & biomass fuel. Host factors that predispose to COPD includes abnormal lung development, genetic abnormalities and accelerated ageing<sup>3</sup>.

Cardiac involvement is common in patients with COPD particularly in those with low oxygen saturation<sup>3</sup>.It is essential to assess the extent of impairment of pulmonary function and the pulmonary hypertension caused by the same to establish the long-term prognosis of the disease . It affects the function of right ventricle leading to corpulmonale and has a poor prognosis<sup>3</sup>. so, early recognition of RV dysfunction and pulmonary hypertention may help in treatment and prolonging the survival of patients with corpulmonale.

The aim of this study is to study the clinical profile of patients and to assess ventricular dysfunction by utilizing ventricular parameters (obtained by clinical methods ,electrocardiography & echocardiography) and to correlate it with severity of COPD. Early recognition and treatment of ventricle dysfunction in COPD leads to prolong survival and improve quality of life.

**METHOD**

A prospective type of observational study was done in COPD patients above 30 years of age, attending the out patient clinic and who were admitted in medicine and pulmo medicine unit in navsari civil hospital were selected for this study.100 patients who fulfilled the inclusion & exclusion criteria were selected.Data has been collected over 2 year.

**Inclusion Criteria**

# COPD patients male & female age > 30 year with taking consent.

**Exclusion Criteria**

- # Patients have known respiratory condition other than COPD.
- # Patients having cardiac disease.
- # Patients in whom meaningful examination could not be performed were also excluded.

All patients were diagnosed clinically and satisfying the

spirometric criteria were included in this study. All patients were subjected to routine blood investigation, sputum analysis for AFB stain , gram stain , culture and drug sensitive test, ECG, chest X -ray ,Echo cardiography, pulmonary function test was done using spirometry. Responses were collected and analysed by microsoft excel.

**RESULT**

It was a prospective type observational study conducted in 100 cases of COPD. All cases were diagnosed clinically and confirmed by spirometry. Mean age of patient was 55.02 ±10 year. In this study 84% patients were male ; all males were smokers. 16 % were female; all females were exposed to biomass fuel. The mean duration of smoking observed was 26.8 ± 3.5 years. Majority of smokers (63%) had history of smoking more than 20 pack years. The mean duration of disease was 4.08 ± 3.2 years. Most of the patients had breathlessness (92%) ,cough with sputum (85%), swelling of feet (34%), fever (8.5%). The clinical signs at presentation were tachypnoea (75%), loud P2 (36%), parasternal heave (23%) ; evidence of congestive cardiac failure like raised JVP (34%) , pedal edema (24%) ,ascites (6%). A wide range of COPD severity was found. Although, the majority of patients had moderate to severe disease.

Disease severity (FEV1 of % Predicted)	No. of patients	Percentage
Mild	7	7%
Moderate	44	44%
Severe	38	38%
Very severe	11	11%

The chest X-ray findings were hyper inflation of lung (73%), increased bronchovascular marking (54%) ,evidence of pulmonary hypertention i.e.prominent right descending pulmonary artery (28%), cardiomegaly (13%) were observed.The ECG findings in copd patients were right axis deviation(44%), P pulmonale (37%), low voltage QRS complex (10%) poor R wave progression (39%), right ventricl hypertrophy (26%). Echocardiography findings were PAH (62%), corpulmonale (56%), RVD (45%), RAD (37%) right ventricle hypertrophy (27%), left ventricle systolic dysfunction(31%), left ventricle diastolic dysfunction(6%).

**DISCUSSION**

COPD is considered a worldwide cause of chronic morbidity and mortality <sup>4</sup>. COPD is a male dominant disease, due to higher prevalence of smoking and having more susceptibility than female <sup>3</sup>.This study is similar to manasa reddy et al <sup>5</sup>, bennita et al<sup>6</sup>, suma et al <sup>7</sup>,swathi talari et al <sup>8</sup>. In this study ECG changes were Right axis eviation (44%), P pulmonale (37%), low voltage QRS complex (10%), poor R wave progression (39%), RVH (26%). ECG changes observed in this study

correlates with manasa reddy et al<sup>5</sup>, sachin et al<sup>9</sup>, padmavati & raizada et al<sup>10</sup>, sekhar et al<sup>11</sup>. P pulmonale, RVH, right axis deviation in copd is due to change in haemodynamics of pulmonary vessels secondary to hypoxia<sup>12</sup>. Hyperinflation of lung in COPD causes poor R wave progression, low voltage QRS complex<sup>12</sup>. In this study most common echo finding was pulmonary arterial hypertension (82%). It appeared more in severe and very severe grade of copd than in mild/ moderate grades. There is significant structural changes occurs in patients with COPD. Alveolar hypoxemia and chronic ventilator insufficiency leads medial hypertrophy & intimal insufficiency which causes pulmonary vasoconstriction, pulmonary vessel bed destruction. Similar findings were seen in suma et al<sup>7</sup>, sekhar et al<sup>11</sup>, benita et al<sup>6</sup>, mansa reddy et al<sup>5</sup>, yaseer et al<sup>13</sup>. Rise in PAH causes increase after load which leads structural and functional changes in right ventricle i.e. right ventricular hypertrophy, right ventricle dilatation, right atrial dysfunction, corpulmonale. In this study right ventricular dilatation (45%), corpulmonale (56%), right atrial dilatation (37%), right ventricle hypertrophy (27%). Similar findings were seen at mansa reddy et al<sup>5</sup>, N.K. gupta et al<sup>14</sup>. In this study 31% patients had left ventricle systolic dysfunction while 6% had left ventricle diastolic dysfunction. In COPD patients hyper inflated lungs, reduced stroke volume and right ventricle dysfunction leads reduced left ventricle systolic dysfunction. Similar findings were seen in N.K.GUPTA et al<sup>14</sup>, yasir et al<sup>13</sup>, mansa reddy et al<sup>5</sup>.

obstructive pulmonary disease patient and its correlation with the severity of disease. Lung India. 2011 Apr-jun;28(2): 105-109. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3109831/>

### CONCLUSION

COPD is a common respiratory disease. This study shows presence of cardiac comorbidity in COPD patients. ECG, echocardiography can be used for screening of COPD. Echocardiography have a better modality for detection of cardiac abnormality than ECG. Echocardiography is less informative than pulmonary catheterization; being noninvasive and easily available echocardiography routinely recommends in copd patients. This study suggests early cardiac screening of all COPD patients should be done to assess prognosis and for better outcome.

### REFERENCES

1. Lozano R, Naghavi M, Foreman K, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380(989):209-128.
2. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006;3(11):e442.
3. 21st edition Harrison's principle of internal medicine Chapter no 292.
4. Global initiative for chronic obstructive pulmonary disease- global strategy for diagnosis, management and prevention of chronic obstructive pulmonary disease. <https://goldcopd.org/>
5. Manasa reddy musku, Abdul rafi shaik, Roopa reddy Mulamalla. A study of ECG and ECHOCARDIOGRAPHIC findings in chronic obstructive pulmonary disease patients in a tertiary care hospital. <https://uk.search.yahoo.com/search?ei=UTF-8&fr=crmas&p=mansara+reddy+musuku+study+of+ecg+and+echocardiography>
6. Benita mary.v et al. A study of ECG and echocardiographic findings in copd & correlation of right ventricular dysfunction with the severity of COPD. [https://uk.search.yahoo.com/search;\\_ylt=Awr9zRFWU99kfbMG.25B4iA5;to p&p=benitta+merry+a+study+of+ecg+and+echocardiography+finding+in+copd](https://uk.search.yahoo.com/search;_ylt=Awr9zRFWU99kfbMG.25B4iA5;to p&p=benitta+merry+a+study+of+ecg+and+echocardiography+finding+in+copd).
7. Suma et al. A study of Electrocardiographic and Echocardiographic changes in chronic obstructive pulmonary disease (COPD) of different grades of severity. *Journal of Evolution of medical and dental sciences* (vol.4)
8. Swathi talari et al. The study of electrocardiographic and echocardiographic changes in copd. January 2018. *Journal of Evolution of medical and dental sciences* 7 (02): 135-141. [https://www.researchgate.net/publication/323773580\\_THE\\_STUDY\\_OF\\_ELECTROCARDIOGRAPHIC\\_AND\\_ECHOCARDIOGRAPHIC\\_CHANGES\\_IN\\_COPD](https://www.researchgate.net/publication/323773580_THE_STUDY_OF_ELECTROCARDIOGRAPHIC_AND_ECHOCARDIOGRAPHIC_CHANGES_IN_COPD).
9. Sachin et al. A Study of ECG findings in copd patients in a tertiary care center. *European journal of molecular & clinical medicine*. Volume 10. 4004-4012. <https://ejmcm.com/issue-content/study-of-ecg-findings-in-copd-patients-in-a-tertiary-care-centre-1711>.
10. Padmavati and Raizada et al. Electrocardiogram in corpulmonale. *British heart journal* 1972 July 34(7) 658-667.
11. Sekhar et al. Electrocardiographic abnormalities in patients with acute exacerbation of chronic obstructive pulmonary disease. *Cureus*. 2021 Dec; 14 (12): e20820 published online. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC458516/?page=1>.
12. ECG in Chronic Obstructive pulmonary disease. <https://litfl.com/ecg-in-chronic-obstructive-pulmonary-disease/>
13. Yaseer et al. Cardiac evaluation of patients with chronic obstructive pulmonary disease using echocardiography. *The Egyptian journal of bronchology* 13, 12-16 (2019). [https://ejb.springeropen.com/articles/10.4103/ejb.ejb\\_2\\_18#citeas](https://ejb.springeropen.com/articles/10.4103/ejb.ejb_2_18#citeas)
14. N.K. Gupta et al. Electrocardiographic evaluation of heart in chronic