



ORIGINAL RESEARCH PAPER

Psychiatry

PREVALENCE AND ASSOCIATION OF COGNITIVE IMPAIRMENT IN PATIENTS WITH MODERATE TO SEVERE DEPRESSION: A CROSS SECTIONAL STUDY

KEY WORDS: cognitive impairment, depression, prevalence

Vighnesh K. Singh	Junior Resident Department Of Psychiatry, Career Institute Of Medical Sciences, Lucknow
Nirnay Sachdeva	Junior Resident Department Of Psychiatry, Career Institute Of Medical Sciences, Lucknow
Abbas Mehdi*	Professor & Head, Department Of Psychiatry, Career Institute Of Medical Sciences, Lucknow *Corresponding Author
Aditi Jain	Assistant Professor, Department Of Psychiatry, Career Institute Of Medical Sciences, Lucknow

ABSTRACT

Background: Almost all depressive disorder patients have cognitive impairments to a certain extent. Depression and cognitive dysfunction share a common neuropathological platform in cortical and sub-cortical brain areas. The current study was designed to explore the prevalence of cognitive impairment in severe depression and to examine the correlates of cognitive impairment and depression with other sociodemographic variables. **Methods:** A hospital based cross-sectional study on the prevalence of cognitive impairment in moderate to severe depression. Total 100 patients were evaluated using Hamilton rating scale for depression (HAM-D), Addenbrooke's cognitive examination (ACE-III). **Results:** The mean age of study sample was 31.92 ± 7.62 (SD) years. The mean Hamilton Depression Rating Scale score for severe depression was 21.70 ± 1.34 (SD: 1.34). 56(56.0%) patients suffering from depression found to have mild cognitive impairment followed by 28 (28.0%) having major neurocognitive disorder. The mean neurocognitive functions in Mild cognitive impairment was 66.84 ± 2.41 , and major neurocognitive disorder was 54.96 ± 4.83 . **Conclusion:** Cognitive impairment and depression were well known in elderly population. Present study suggests that more than one fourth of young age patients having severe depression reported cognitive impairment which is more prevalent in the females than in the males. Hence, psychiatrist should pay special attention for early detection and treatment of cognitive impairment.

INTRODUCTION

Cognitive dysfunction is characterised by impairment in the following areas: problem-solving, processing speed, visual and auditory processing, verbal and nonverbal learning, short-term and working memory, attention, and motor functioning. A major mediator of functional impairment in major depressive disorder (MDD) may be cognitive dysfunction^[1]. Individuals who appear with MDE usually exhibit self-reported measures of reduced concentration and attention as part of MDD. Furthermore, when treating MDD, it is frequently discovered that cognitive abnormalities continue even during times of symptomatic remission^[2], supporting the disconnect between emotional and functional improvement whereas cognitive impairment is a common often persistent, symptom of major depressive disorder that is disproportionately represented in patients who have not returned to full psychosocial functioning^[3].

Despite remission from depressive symptoms, cognitive impairments often linger, causing considerable functional burden. Its debilitating effects extend beyond emotional distress, disrupting work performance, social interactions, and overall quality of life. Recent research suggests a strong link between depression and cognitive decline, with depression potentially acting as a prodrome to dementia. Late-life depression, in particular, frequently presents with a constellation of cognitive difficulties, raising concerns about future cognitive deterioration.^[4]

It has also been suggested that depression represents a risk factor for the development of future dementia^[5]. Because MCI is conceptualized as a predementia state, it is likely that factors associated with dementia (including depression) are also linked to MCI. Despite this, comparatively little is known about the prevalence of depressive symptoms in MCI. The substantial overlap in key symptoms between depression and dementia complicates research in the area. While depression is a disorder of affect and dementia a disorder of cognition, in reality, the two conditions share a number of symptoms in common.

In light of these insights, it becomes clear that cognitive impairment should not be viewed as a mere consequence of depression. Instead, it emerges as a distinct dimension of the disorder, demanding targeted treatment alongside symptom management. Cognitive symptoms should therefore be regarded as a partially independent dimension of MDD, and an important target of any treatment that is initiated. So, we aim to study the cognitive impairment in patients with moderate to severe depression.

Aim of the study

Current study aimed to estimate the prevalence of cognitive impairment in moderate and severe depression and to examine the specific cognitive impairments and its correlation with different degree of depression.

MATERIAL AND METHODS-

Study design-

The present study design is a cross-sectional observational study was conducted in the Out Patient Department (OPD) and In Patient Department (IPD) of the Department of Psychiatry, Career institute of medical sciences and hospital, Ghaila, Lucknow during the study period from June 2022 to May 2023. All the patients who fulfilled the inclusion and exclusion criteria were enrolled in the study, after obtaining their written informed consent in Hindi/English.

Participants-

In this study, a total of 100 diagnosed cases of depression were selected from the patients visiting the inpatient and outpatient facility of the Department of Psychiatry of a tertiary care teaching hospital in North India. Semi-structured interview Proforma was used to collect the identification data, socio-demographic data and clinical data of the selected cases that included a socio-demographic questionnaire. The Hamilton Depression Rating Scale (HDRS) was applied to assess the severity of depression and the Addenbrooke's Cognitive Examination (ACE III) was administered to assess the level of cognitive impairment.

DISCUSSION-

Major Depressive Disorder (MDD) is one of the most common psychiatric disorders with an estimated lifetime prevalence of 16%.^[6] MDD, however, is not solely characterized by its high prevalence but also by its major global impact. Our study findings indicate that a significant number of patients having moderate to severe depression develop cognitive impairment, in which the majority of the moderate depressive patients develop mild cognitive impairment and almost half of the severely depressed patients develop mild cognitive impairment.

Our findings were consistent with the findings of **Ali M et al**^[7] who did a study on evaluation of cognitive impairment in patients with major depressive disorder in remission who reported that the mean age of patients was 33.0±8.2 years, 77.7% were female, the average age at onset of MDD was 23.3±6.3 years. The mean HAM-D score in moderate depression was 14.67±0.71 and in severe depression was 21.70±1.34 and the association of mean HAM-D score in moderate and severely depressed patients was found to be statistically significant (**p<0.05**).

In the present study, we noted that 56.0% of patients had mild cognitive impairment followed by 28.0% of patients had major cognitive impairment and the rest 16.0% of patients were normal. **Maramisa MM et al**^[8] reported that depression, cognitive impairment preceded depression, but it might be caused by anxiety-like behavior that occurred in early stimulation of chronic unpredictable mild stress. The mean ACE-III scores in normal were 74.88±2.33, in the cases of mild cognitive impairment it was 66.84±2.41 and in severe cognitive impairment was 54.96±4.83 and the association of mean ACE-III scores of normal, mild and major cognitive disorder patients was found to be highly significant (**p<0.01**).

Manit S et al^[9] reported that patients with mild-to-moderate depression reported perceived cognitive dysfunction and functional disability. **Douglas KM et al**^[10] reported the prevalence of cognitive impairment was highest for the inpatient depression sample and lowest for the outpatient depression sample. **Helmich LH et al**^[11] reported the burden of functional impairment in MDD and the importance of recognizing and managing cognitive symptoms in daily practice.

Our study findings clearly indicate that significant number of patients having moderate to severe depression develop cognitive impairment, in which majority of the moderate depressive patients develop mild cognitive impairment and almost half of the severely depressed patients develop mild cognitive impairment. Study revealed that major neuro cognitive disorder is strongly associated with the severity of depression.

The study revealed that major neurocognitive disorder is strongly associated with the severity of depression. Both depressive mood and cognitive impairment are associated with poor psychosocial functioning. As a result, we argue that treating cognitive impairment and alleviating depressive symptoms are both important in improving outcomes for patients with depression. In conclusion, this hospital-based cross-sectional, observational study shows that cognitive impairment is a core feature of depression that cannot be considered an epiphenomenon secondary to symptoms of low mood, and that it may be a valuable target for future interventions.

Limitations of the study

This is a cross-sectional study in which causal relationships between the variables cannot be determined. Future studies might consider alternate designs to examine how the relationship between clinical predictors and cognitive impairment changes throughout the course of depression.

CONCLUSION-

Our study findings indicate that a significant number of patients having moderate to severe depression develop cognitive impairment, in which the majority of the moderate depressive patients develop mild cognitive impairment and almost half of the severely depressed patients develop mild cognitive impairment. The study revealed that major neurocognitive disorder is strongly associated with the severity of depression.

Both depressive mood and cognitive impairment are associated with poor psychosocial functioning. As a result, we argue that treating cognitive impairment and alleviating depressive symptoms are both important in improving outcomes for patients with depression. In conclusion, this hospital-based cross-sectional, observational study shows that cognitive impairment is a core feature of depression that cannot be considered an epiphenomenon secondary to symptoms of low mood, and that it may be a valuable target for future interventions.

REFERENCES-

- McIntyre RS, Cha DS, Soczynska JK, Woldeyohannes HO, Gallagher LA, Kudlow P, et al. Cognitive deficits and functional outcomes in major depressive disorder: determinants, substrates, and treatment interventions. *Depress Anxiety*. 2013;30(6):515-527.
- McIntyre RS, Xiao HX, Syeda K, Vinberg M, Carvalho AF, Mansur RB, et al. The Prevalence, measurement, and treatment of the cognitive dimension/domain in major depressive disorder. *CNS Drugs*. 2015;29:577-89.
- Culpepper L, Lam RW, McIntyre RS. Cognitive Impairment in Patients With Depression: Awareness, Assessment, and Management. *J Clin Psychiatry*. 2017;78(9):1383-1394.
- Battaglia A, Dubini A, Mannheimer R, Pancheri P. Depression in the Italian community: epidemiology and socio-economic implications. *Int Clin Psychopharmacol*. 2004;19:135-142.
- Ownby RL, Crocco E, Acevedo A, John V, Loewenstein D. Depression and risk for Alzheimer disease: systematic review, meta-analysis, and meta-regression analysis. *Arch Gen Psychiatry*. 2006;63:530-538.
- Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, et al: The epidemiology of major depressive disorder: results from the national comorbidity survey replication (NCS-R). *Jama* 2003; 289:3095-3105
- Ali M, Fahmy M, Haggag W, El-Tantawy A and Hassan H. Evaluation of cognitive impairment in patients with major depressive disorder in remission. *Middle East Current Psychiatry* 2021;28:71
- Maramisa MM, Mahajudina MS, Khotib J. Impaired Cognitive Flexibility and Working Memory Precedes Depression: A Rat Model to Study Depression. *Neuropsychobiology*. 2021;80:225-233.
- Manit S, Yee Ming M, Yen Kuang Y, Herng-Nieng C, Constantine DD, Zuraida ZN et al. Cognitive Dysfunction in Asian Patients with Depression (CogDAD): A Cross-Sectional Study. *Clinical Practice & Epidemiology in Mental Health* 2017; 13(1): 185-199.
- Douglas KM, Gallagher P, Robinson LJ, Carter JD, McIntosh VV, Frampton CM, et al. Prevalence of cognitive impairment in major depression and bipolar disorder. *Bipolar Disord*. 2018;00:1-15.
- Helmich LH, Haro JM, Jonsson B, Melac AT, Nicola SD, Chollet J et al. Functional impairment in patients with major depressive disorder: the 2-year PERFORM study. *Neuropsychiatric Disease and Treatment* 2018;14:239-249.